Hemovigilance

Hans-Gert Heuft^a Behrouz Mansouri Taleghani^b

^a Institute for Transfusion Medicine, Hannover Medical School, Hanover, Germany;

^b University Department of Hematology, Inselspital, University of Bern, Bern, Switzerland

According to the International Haemovigilance Network, ‘hemovigilance’ is defined as ‘a set of surveillance procedures covering the whole transfusion chain (from the collection of blood and its components to the follow-up of recipients), intended to collect and assess information on unexpected or undesirable effects resulting from the therapeutic use of labile blood products, and to prevent their occurrence or recurrence’ [1].

Within a period of 4 years, TRANSFUSION MEDICINE AND HEMOTHERAPY again presents the topic ‘Hemovigilance’. In 2014 several papers considered the influence of blood product characteristics or modifications on product safety and transfusion efficacy [2–6]. The current issue deals with the nationwide implementation of pathogen inactivation (PI) for all platelet concentrates (PC) in Switzerland, as presented by Markus Jutzi and colleagues [7]. One of their major aims was to eliminate bacterial infections due to PC transfusion. Jutzi et al. [7] reported that PI was successful to reduce the PC bacterial infection rate in Switzerland since 2011 to zero. The focus of this issue, however, is the interaction between the Rhesus blood group system and the recipients of red blood cell (RBC) components and/or PC [8–11]. The key question is: Do RhD– patients suffer harm from the (sometimes inevitable) transfusion of RhD+ RBC or PC? To answer this question, three author groups analyzed the consequences of RhD unequal transfusion in different study settings. The trigger for this type of studies is the unavailability of a sufficient Rh antigen-compatible blood supply. Flommersfeld and colleagues [8] reported their experience with an emergency transfusion protocol applying blood group O RhD+ RBC as first line treatment in exsanguinating polytrauma patients. In their retrospective analysis that covered a period of 5 years they identified 62 patients (of whom 14 were RhD– finally) who received 259 potentially and 60 definitely RhD-different RBC transfusions. In contrast to this study, Villalba et al. [9] and Reckhaus et al. [10] presented data from RhD-different platelet (PLT) transfusions. Villalba et al. retrospectively studied the formation of anti-D in a group of 48 RhD– female leukemia patients below 55 years of age (roughly ‘childbearing age’ and

thus at risk for hemolytic disease of the newborn (HDN)) following the transfusion of 949 RhD+ PC. Their study covered a period of nearly 20 years and detected 2 cases with anti-D out of 48 patients (4%). Reckhaus et al. [10] investigated in depth hematological patients who formed new Rh antibodies without cognate antigen exposition during RBC transfusion, but with cognate antigen exposition during PLT transfusion. Over a study period of about 2 years, they found 11 patients with this kind of transfusion history: 6 patients with anti-D alone and 5 patients without anti-D, but with other Rh antibodies such as anti-E (2), anti-c/-E (2) and even anti-f (1). The patients had been transfused with a total of 156 Rh antigen-positive PLT concentrates. Ten out of 11 patients predominantly received buffy coat-derived PLT concentrates (BC-PC), either alone (n = 3) or combined with lower numbers of apheresis-derived PC (A-PC, n = 7). BC-PC were shown to contain a more than 20-fold higher RBC contamination than A-PC (median RBC content of $0.30 \times 10^9/l$ (BC-PC) vs. $0.014 \times 10^9/l$ (A-PC)). Taking the results of these three studies together, they transfer good and bad news. To start with the good news: Rh antigen-different transfusion did not contribute to immediate and dangerous harm. Neither RBC transfusion nor PLT recipients developed any hemolytic transfusion reaction. Moreover, A-PC might transfer a lower risk for RBC alloantibody induction than BC-PC. The bad news is: The posttransfusion Rh antibody formation might be higher than previously expected. Rh antibody formation could also be caused by PLT transfusion. To some extent the results of Villalba et al. [9] and Reckhaus et al. [10] are surprising as there is a relatively consistent body of literature that disputed or even denied the induction of RBC alloimmunization by PLT transfusion, in particular, if A-PC are exclusively applied [12–19]. Is there a resolution for these conflicting results? One relatively new explanation for a possible induction of Rh alloantibodies by antigen-positive PLT transfusions was offered by Kitazawa et al. [20]. They observed Rh alloantibody formation (anti-E, anti-C, anti-c) after PLT transfusion. Because RBC contamination in A-PC is probably insufficient to trigger alloimmunization [19, 21], Kitazawa et al.

[20] hypothesized that RBC microparticles could contribute to RBC alloimmunization. Microparticles can be phagocytosed and therefore could be as immunogenic as RBC. Moreover, both residual RBC and RBC microparticles could 'cooperate' to induce Rh alloantibody formation [22]. In view of these new hypotheses and the findings presented by Villalba et al. [9] and Reckhaus et al. [10] in this issue of TRANSFUSION MEDICINE AND HEMOTHERAPY, national and institutional guidelines for PC transfusion should be

critically reviewed. This is in particular true, if BC-PC are routinely applied in hematological patients or in patients at risk for HDN.

Disclosure Statement

The authors have no conflicts of interest to disclose.

References

- 1 International Haemovigilance Network: Definition of Haemovigilance. www.ihn-org.com/about/definition-of-haemovigilance/ (last accessed May 15, 2018)
- 2 Taleghani BM, Heuft HG: Hemovigilance. *Transfus Med Hemother* 2014;41:170-171.
- 3 Jimenez-Marco T, Ruiz-Alderton D, Bautista-Gili AM, Girona-Llobera E: Role of riboflavin- and UV light-treated plasma on prevention of transfusion-related acute lung injury. *Transfus Med Hemother* 2014;41:172-175.
- 4 Julmy F, Ammann RA, Fontana S, Mansouri Taleghani B, Hirt A, Leibundgut K: Transfusion efficacy of apheresis platelet concentrates irradiated at the day of transfusion is significantly superior compared to platelets irradiated in advance. *Transfus Med Hemother* 2014;41:176-181.
- 5 Fontana S, de la Cuadra C, Müller U, Schmid P, Perler M, Luginbühl M, Mansouri Taleghani B: A simple guideline reduces the need for red blood cell transfusions in Swiss hospitals: a prospective, multicentre, before-and-after study in elective hip and knee replacement. *Transfus Med Hemother* 2014;41:182-188.
- 6 Zhu M, Xu W, Wang B-L, Su H: Hemostatic function and transfusion efficacy of apheresis platelet concentrates treated with gamma irradiation in use for thrombocytopenic patients. *Transfus Med Hemother* 2014;41:189-196.
- 7 Jutzi M, Mansouri Taleghani B, Rueesch M, Amsler L, Buser A: Nationwide implementation of pathogen inactivation for all platelet concentrates in Switzerland. *Transfus Med Hemother* 2018;45: doi 10.1159/000489900.
- 8 Flommersfeld S, Mand C, Kühne CA, Bein G, Ruchholtz S, Sachs U: Unmatched type 0 Rh D+ red blood cells in multiple injured patients. *Transfus Med Hemother* 2018;45:DOI: 10.1159/000485388.
- 9 Villalba A, Santiago M, Freiria C, Montesinos P, Gomez I, Fuentes C, Rodriguez-Veiga R, Fernandez JM, Sanz G, Sanz MA, Carpio N, Solves P: Anti-D alloimmunization after RhD-positive platelet transfusion in RhD-negative women under 55 years diagnosed with acute leukemia: Results of a retrospective study. *Transfus Med Hemother* 2018;45:DOI: 10.1159/000488804.
- 10 Reckhaus J, Jutzi M, Fontana S, Bacher VU, Vogt M, Daslakis M, Mansouri Taleghani B: Platelet transfusion induces alloimmunization to D and non-D Rhesus antigens. *Transfus Med Hemother* 2018;45:167-172.
- 11 Kulkarni SS, Gogri H, Parchure D, Mishra G, Ghosh K, Rajadhyaksha S, Madkaikar M, Férec C, Fichou Y: RHD-positive alleles among D- C/E+ individuals from India. *Transfus Med Hemother* 2018;45: doi 10.1159/000479239.
- 12 Atoyebi W, Mundy N, Croxton T, Littlewood TJ, Murphy MF: Is it necessary to administer anti-D to prevent RhD immunization after the transfusion of RhD-positive platelet concentrates? *Br J Haematol* 2000;111:980-983.
- 13 Molnar R, Johnson R, Sweat LT, Geiger TL: Absence of alloimmunization in D- pediatric oncology patients receiving D-incompatible single-donor platelets. *Transfusion* 2002;42:177-182.
- 14 Cid J, Ortin X, Elies E, Castella D, Panades M, Martín-Vega C: Absence of anti-D alloimmunization in hematologic patients after D-incompatible platelet transfusions. *Transfusion* 2002;42:173-176.
- 15 Cid J, Carbassé G, Pereira A, Sanz C, Mazzara R, Escolar G, Lozano M: Platelet transfusions from D+ donors to D- patients: a 10-year follow-up study of 1014 patients. *Transfusion* 2011;51:1163-1169.
- 16 O'Brien JL, Haspel RL, Uhl L: Anti-D alloimmunization after D-incompatible platelet transfusions: a 14-year single-institution retrospective review. *Transfusion* 2014;54:650-654.
- 17 Weinstein R, Simard A, Ferschke J, et al: Prospective surveillance of D- recipients of D+ apheresis platelets: alloimmunisation against D is not detected. *Transfusion* 2015;55:1327-1330.
- 18 Cid J, Lozano M, Ziman A, West KA, O'Brien KL, Murphy MF, Wendel S, Vazquez A, Ortin X, Hervig TA, Delaney M, Flegel WA, Yazer MH: Low frequency of anti-D alloimmunization following D+ platelet transfusion: the anti-D alloimmunization after D-incompatible platelet transfusions (ADAPT) study. *Br J Haematol* 2015;168:598-603.
- 19 Cid J, Yazer MH, Lozano M: Platelet transfusion and respecting patient D type. *Curr Opin Hematol* 2015;2:540-546.
- 20 Kitazawa J, Nollet K, Morioka H, Tanaka K, Inomata M, Kubuki Y, Ohto H: Non-D Rh antibodies appearing after apheresis platelet transfusion: stimulation by red cells or microparticles? *Vox Sang* 2011;100:395-400.
- 21 Hendrickson JE, Eisenbarth SC, Tormey CA: Red blood cell alloimmunization: new findings at the bench and new recommendations for the bedside. *Curr Opin Hematol* 2016;23:543-549.
- 22 Rubin O, Canellini G, Delobel J, Lion N, Tissot JD: Red blood cell microparticles: clinical relevance. *Transfus Med Hemother* 2012;39:342-327.