Case Report

Case Report: Oxaliplatin-Induced Immune-Mediated Thrombocytopenia

Elizabeth Pan^a Eric Hsieh^a Caroline Piatek^b

^aDepartment of Medicine, Keck School of Medicine of USC, Los Angeles, CA, USA;

^bDivision of Hematology, Department of Medicine, Keck School of Medicine of USC, Los Angeles, CA, USA

Keywords

Adenocarcinoma · Oxaliplatin · Thrombocytopenia

Abstract

Thrombocytopenia is a frequent complication of cancer may be due to a variety of causes including malignancy itself, acute disease processes, or cancer therapy. Systemic cancer therapy is the most common cause of thrombocytopenia in cancer patients observed nearly two-thirds of patients with solid tumors. Thrombocytopenia with traditional chemotherapy agents is most frequently the result of megakaryocyte cytotoxicity. Oxaliplatin is a platinum derivative commonly used in gastrointestinal malignancies and is associated with drug-induced immune thrombocytopenia.

© 2018 The Author(s)
Published by S. Karger AG, Basel

Case Presentation

A 36 year-old woman with Her-2 negative metastatic gastric adenocarcinoma currently on capecitabine and oxaliplatin presented for oxaliplatin infusion. Laboratory testing one day prior showed a platelet count of 237×10^9 and hemoglobin of 10.9 g/dL with an MCV of 81.9 attributed to iron deficiency anemia. She had previously received 5-fluorouracil/leucovorin/oxaliplatin for 14 months and then was switched to single agent capecitabine for 2

months in the setting of stable disease. At progression of disease, oxaliplatin was added to capecitabine, which the patient had been on for the past 9 months.

Thirty minutes after the oxaliplatin infusion was started, the patient developed significant bleeding from her gums, epistaxis, and had 150 to 200 mL of bloody oral secretions. Laboratory testing showed a platelet count of $<5 \times 10^9$, white blood cell count of 2.6×10^9 , hemoglobin 6.2 g/dL, PT 16.4, INR 1.4, total bilirubin 0.5 mg/dL, LDH 211 U/L. Peripheral smear showed few platelets and no schistocytes. The patient was admitted and received intravenous immunoglobulin 1 g/kg, dexamethasone 40 mg IV for 4 days, and oral aminocaproic acid, with resolution of bleeding after one day. No platelet transfusions were given. She received 1 unit of packed red blood cells. The platelet count improved to 77×10^9 after 4 days and normalized to 341×10^9 one month later. Oxaliplatin was definitively discontinued.

Discussion

Although myelosuppression is the most common cause of thrombocytopenia in cancer patients receiving oxaliplatin, there are several other recognized mechanisms of oxaliplatin-induced thrombocytopenia [1–5]. With myelosuppression, thrombocytopenia is usually asymptomatic and is accompanied by anemia and neutropenia [6]. Management includes observation and the occasional need for dose reductions or delays [7]. Oxaliplatin-induced ITP is a well-recognized but uncommon etiology of thrombocytopenia. The mechanism is platelet destruction mediated by oxaliplatin-dependent antibodies to platelet antigens, leading to a sudden drop in platelet count to $<10 \times 10^9$ and bleeding manifestations within several hours of oxaliplatin infusion. This typically occurs after >12 cycles of oxaliplatin and may be preceded by hypersensitivity reaction [7, 8]. Management includes platelet transfusion, corticosteroid therapy, and IV immunoglobulin therapy with resolution of thrombocytopenia within approximately 2 weeks. Definitive discontinuation of oxaliplatin is recommended [1, 9, 10].

Statement of Ethics

The authors have no ethical conflicts to disclose

Disclosure Statement

The authors have no conflicts of interest to declare.

References

- 1 Jardim DL, Rodrigues CA, Novis YA, Rocha VG, Hoff PM. Oxaliplatin-related thrombocytopenia. *Ann Oncol*. 2012 Aug;23(8):1937–42.
- 2 Bautista MA, Stevens WT, Chen CS, Curtis BR, Aster RH, Hsueh CT. Hypersensitivity reaction and acute immune-mediated thrombocytopenia from oxaliplatin: two case reports and a review of the literature. *J Hematol Oncol*. 2010 Mar;3(1):12.
- 3 Taleghani BM, Meyer O, Fontana S, Ahrens N, Novak U, Borner MM, et al. Oxaliplatin-induced immune pancytopenia. *Transfusion*. 2005 May;45(5):704–8.
- 4 Curtis BR, Kaliszewski J, Marques MB, Saif MW, Nabelle L, Blank J, et al. Immune-mediated thrombocytopenia resulting from sensitivity to oxaliplatin. *Am J Hematol*. 2006 Mar;81(3):193–8.

- 5 Ohta S, Cho Y, Oshima S, Hosoya O, Juni K, Kojima H. Oxaliplatin-induced acute-onset thrombocytopenia and hemorrhage: case report and review of the literature. *Oncol Lett*. 2012 Jun;3(6):1297–300.
- 6 Suh SE, Jang MJ, Chong SY, Aster RH, Curtis BR, Oh D. A case of oxaliplatin-induced immune-mediated thrombocytopenia. *Blood Res*. 2014 Mar;49(1):61–4.
- 7 Wu Y, Aravind S, Ranganathan G, Martin A, Nalysnyk L. Anemia and thrombocytopenia in patients undergoing chemotherapy for solid tumors: a descriptive study of a large outpatient oncology practice database, 2000–2007. *Clin Ther*. 2009;31(Pt 2):2416–32.
- 8 Dold FG, Mitchell EP. Sudden-onset thrombocytopenia with oxaliplatin. *Ann Intern Med*. 2003 Jul;139(2):E156.
- 9 Piatek CI, Liebman HA. Thrombocytopenia in patients with solid tumors or hematologic malignancies. *Platelets in Thrombotic and Non-Thrombotic Disorders*. DOI: 10.1007/978-3-319-47462-5_56.
- 10 Liebman HA. Thrombocytopenia in cancer patients. *Thromb Res*. 2014 May;133 Suppl 2:S63–9.