Neonatal Autopsy: A 21st Century Approach?

Susan C. Shelmerdine^{a, b} Willemijn Klein^c Owen J. Arthurs^{a, b}

^aDepartment of Clinical Radiology, Great Ormond Street Hospital for Children, London, UK; ^bUCL Great Ormond Street Institute of Child Health, Great Ormond Street Hospital for Children, London, UK; ^cDepartment of Radiology and Nuclear Medicine, Radboud University Medical Center, Nijmegen, The Netherlands

Dear Editor,

We read with interest the article in *Neonatology* by de Sévaux et al., entitled “The Value of Autopsy in Neonates in the 21st Century” [1]. We agree that autopsy consent rates are declining, and offering parents an accurate, noninvasive method alongside or instead of autopsy is to be encouraged [2]. However, we have significant reservations about the study and manuscript in terms of its scientific rigor, methodology, and the conclusions drawn from them.

The authors retrospectively analyzed clinical notes and autopsy findings for neonatal deaths on a Dutch NICU, and they concluded that autopsy has an important role, but over half of the findings would be unlikely detected with postmortem MRI [1]. Of the reported 298 deaths in 2,785 infants on their NICU between 2008 and 2015, only 117 underwent autopsy (of which 100 were eligible for the study). It is likely that patients who underwent autopsy were those in whom the cause of death was unclear, or whose parents were counselled directly towards the benefits of autopsy, as they form less than 40% of the target population. There is no information provided regarding changes in clinical practice or sample distribution over this time. Clinician

bias is a significant factor in directing consent for autopsy [3, 4] and thus, a small sample of approximately 12 deaths per year over an 8-year period is unlikely to be an accurate reflection of local or national clinical practice, where the true yield in an unselected NICU population is likely to be much lower.

We were also extremely surprised by the speculative and retrograde nature of postmortem MRI assessment in this study, in the absence of any imaging performed. The assessment was presented as a subjective opinion of a single radiologist on the basis of a retrospective case-based review [1], which is unusually simplistic and highly unscientific when considered against a background of published evidence of actual postmortem MRI experience [5, 6]. The most rigorous scientific studies will be double-blinded clinical trials, with imaging reported blinded to clinical findings, double reported, or consensus read, in order to minimise subjectivity, training bias, retrospective assumptions, and subjective impressions.

We are also concerned by the “vast” but unquantified experience of the radiologist, particularly as “postmortem MRI is not a standard procedure” in their unit [1]. Our

experience is that training radiologists in postmortem imaging is fundamental to accurate image interpretation and typically requires 200 cases or more [6]. They do not cite their own experience in this field; neither (to the best of our knowledge) does the group participate in national or international postmortem imaging guideline groups [7, 8]. Crucial data is also missing, such as full disclosure regarding what diagnoses were deemed potentially diagnosable (or not). This level of detail is precisely what is required to counsel parents appropriately regarding the likelihood of imaging being useful on an individual case basis. Some of their examples listed as “unlikely detectable” were atrial septal defect (ASD), pneumonia, and hypoplasia of the ductus venosus. Data published in 2015 demonstrate that ASDs are commonly diagnosed [9, 10], but we agree that sepsis is challenging [11], although certain infections (e.g., CMV) may have maternal or placental sequelae that do not require invasive autopsy.

We conclude that this study does not stand up to scientific scrutiny regarding the utility of postmortem MRI. Its results cannot form the basis of wider implications or be used to found generalisations: this study only reports from a small, potentially bi-

ased population who did not in fact undergo the imaging that it argues would not assist the diagnosis. There is growing scientific evidence to support postmortem MRI, openly available for discussion and contribution across international forums [7, 8]. We use this in our routine clinical practice to counsel all parents appropriately during bereavement [6, 12]. We actively encourage others to consider the same approach and participate in multicentre trials, international expert discussion, and educational opportunities such that we can move forwards as a multidisciplinary profession.

Disclosure Statement

The authors have no conflicts of interest to declare.

Funding Sources

S.C.S. is supported by a Research Councils United Kingdom and United Kingdom Research and Innovation Fellowship, and Medical Research Council Clinical Research Training Fellowship (Grant Ref: MR/R00218/1). This award is jointly fund-

ed by the Royal College of Radiologists. O.J.A. is funded by a National Institute for Health Research Career Development Fellowship (NIHR-CDF-2017-10-037). This article presents independent research funded by the Medical Research Council, Royal College of Radiologists, National Institute for Health Research, and the views expressed are those of the author(s) and not necessarily those of the aforementioned institutions, National Health Service, or the Department of Health.

References

- 1 de Sévaux JL, Nikkels PG, Lequin MH, Groenendaal F. The Value of Autopsy in Neonates in the 21st Century. *Neonatology*. 2018 Oct;115(1):89–93.
- 2 Arthurs OJ, Bevan C, Sebire NJ. Less invasive investigation of perinatal death. *BMJ*. 2015 Jul;351:h3598.
- 3 Lewis C, Hill M, Arthurs OJ, Hutchinson JC, Chitty LS, Sebire N. Health professionals' and coroners' views on less invasive perinatal and paediatric autopsy: a qualitative study. *Arch Dis Child*. 2018 Jun;103(6):572–8.
- 4 Lewis C, Hill M, Arthurs OJ, Hutchinson C, Chitty LS, Sebire NJ. Factors affecting uptake of postmortem examination in the prenatal, perinatal and paediatric setting. *BJOG*. 2018 Jan;125(2):172–81.
- 5 Thayyil S, Sebire NJ, Chitty LS, Wade A, Chong W, Olsen O, et al.; MARIAS collaborative group. Post-mortem MRI versus conventional autopsy in fetuses and children: a prospective validation study. *Lancet*. 2013 Jul;382(9888):223–33.
- 6 Ashwin C, Hutchinson JC, Kang X, Langan D, Jones R, Norman W, et al. Learning effect on perinatal post-mortem magnetic resonance imaging reporting: single reporter diagnostic accuracy of 200 cases. *Prenat Diagn*. 2017 Jun;37(6):566–74.
- 7 Arthurs OJ, van Rijn RR, Whitby EH, Johnson K, Miller E, Stenzel M, et al. ESPR post-mortem imaging task force: where we begin. *Pediatr Radiol*. 2016 Aug;46(9):1363–9.
- 8 Sonnemans LJ, Vester ME, Kolsteren EE, Erwich JJ, Nikkels PG, Kint PA, et al.; Dutch post-mortem imaging guideline group. Dutch guideline for clinical foetal-neonatal and paediatric post-mortem radiology, including a review of literature. *Eur J Pediatr*. 2018 Jun;177(6):791–803.
- 9 Taylor AM, Arthurs OJ, Sebire NJ. Postmortem cardiac imaging in fetuses and children. *Pediatr Radiol*. 2015 Apr;45(4):549–55.
- 10 Taylor AM, Sebire NJ, Ashworth MT, Schiavano S, Scott RJ, Wade A, et al.; Magnetic Resonance Imaging Autopsy Study Collaborative Group. Postmortem cardiovascular magnetic resonance imaging in fetuses and children: a masked comparison study with conventional autopsy. *Circulation*. 2014 May;129(19):1937–44.
- 11 Arthurs OJ, Thayyil S, Olsen OE, Addison S, Wade A, Jones R, et al.; Magnetic Resonance Imaging Autopsy Study (MaRIAS) Collaborative Group. Diagnostic accuracy of post-mortem MRI for thoracic abnormalities in fetuses and children. *Eur Radiol*. 2014 Nov;24(11):2876–84.
- 12 Cannie M, Votino C, Moerman P, Vanheste R, Segers V, Van Berkel K, et al. Acceptance, reliability and confidence of diagnosis of fetal and neonatal virtuopsy compared with conventional autopsy: a prospective study. *Ultrasound Obstet Gynecol*. 2012 Jun;39(6):659–65.