Single Case

Flagellate Dermatitis by Shiitake Mushroom

Camilla Santos Ribeiro Celso Tavares Sodré Marcia Ramos-e-Silva

Sector of Dermatology and Post-Graduation Course in Dermatology, University Hospital and School of Medicine, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil

Keywords

Dermatitis · Flagellate dermatitis · Shiitake

Abstract

The authors present a case of flagellate dermatitis in a previously healthy young male, triggered by the consumption of shiitake mushrooms. Complete remission without complications was obtained within 7 days, simply by the application of dexamethasone cream 0.1%. Considering the growing use of this originally Asian fungus in cookery worldwide, it is likely that we will see increased occurrence of this benign and self-limited condition also in the Western world.

© 2019 The Author(s)
Published by S. Karger AG, Basel

Introduction

Lentinula edodes or shiitake is a widely consumed fungus in Asian cooking and currently the second most consumed mushroom worldwide [1] (Fig. 1). Shiitake's pharmacological properties are well-known in Oriental medicine, such as anticarcinogenic and antihypertensive characteristics and of the reduction of serum cholesterol levels. However, the consumption of raw or rare cooked shiitake may cause linear eruptions [2] forming small papules, papular vesicular eruptions, or intensely infiltrated pruriginous plaques called flagellate dermatitis (FD) [3].

Nakamura [4] described the first case of skin lesions after the ingestion of shiitake in 1977, followed by some other authors. The lesions typically appear within a period of 24–48 h after ingestion. The eruption preferably affects the trunk, limbs and neck, regressing spontaneously after a period of few days to up to 2 weeks [3]. The name “flagellate dermatitis” was chosen because of the association with the practice of self-flagellation by the early Christians, whipping each other as an act of purification to pay for their sins. FD lesions are linear and erythematous, resembling injuries caused by whiplashes [3].

Case Presentation

The patient was a previously healthy 23-year-old Caucasian male who complained of the sudden onset of linear rash on his trunk evolving for 2 days. The patient denied other symptoms, such as pruritus, arthralgia, mucosal abnormalities, and fever. When asked about consumption of shiitake, he confirmed that he had eaten those mushrooms 4 days before he sought medical assistance and stated that he thought the shiitakes had been well cooked. The patient denied the intake of other medications or changes of personal hygiene products.

At physical examination, the patient presented with small, grouped papules forming a linear design, spreading on the trunk and extending to the proximal part of the upper limbs (Fig. 2, 3, 4). No alterations on the face, scalp, or mucosa were observed. Dexamethasone ointment was applied for 5 days in the area of the lesions, and complete remission was observed after 7 days.

Discussion

The physiopathology of flagellate dermatitis is not yet well understood. The most accepted theory is that the development of skin lesions is caused by toxicity from the lentinan polysaccharide present in the mycelium of the fungus. This substance promotes vessel dilatation and inflammation caused by means of the interleukin protein [1, 4, 5, 6]. Lentinan is thermolabile, and at temperatures between 130 and 145°C, its molecular structure can be irreversibly changed, which explains why cooked shiitake usually does not cause flagellate dermatitis [1]. Patients treated for gastric and intestinal adenocarcinoma with fungus extract have already developed similar reactions to those caused by shiitake [4].

Discussions also suggest a possible involvement of late sensitization mediated by T cells [3]. Although contact tests for dermatitis by shiitake are negative, allergic reactions such as asthma, contact dermatitis, and alveolitis may occur after the first episode [7].

The histopathological findings are usually nonspecific. Epidermal spongiosis and lymphocyte infiltrate in the dermis can sometimes be noticed [3].

Dermatitis induced by bleomycin must be considered in the differential diagnosis. Clinical findings are similar to FD due to other causes, and histopathological examination is nonspecific; however, there may be mucosal involvement, and, as opposed to what occurs in “shiitake-induced dermatitis,” it may lead to hyperpigmentation [4]. Bleomycin is a cytostatic antibiotic of the glycopeptide class [3] produced by the fermentation of *Streptococcus verticillus* that exerts antitumor activity against cervix squamous cell carcinoma, testicle tumors, and lymphomas [8]. There is also a structural similarity between the sulfur component present in shiitake fungi and bleomycin, which may play a triggering role for FD [1]. Flagellate erythema associated with other drugs such as peplomycin (a bleomycin derivative) and docetaxel and

with adult Still's diseases, dermatomyositis, and HIV infection must be listed as differential diagnosis [9, 10].

The disease disappears spontaneously within up to 2 weeks [11]. Symptoms can be treated with antihistamines, topical corticosteroids, and eventually systemic corticosteroids [10]. It is advisable to avoid further exposure to the fungus to avert the occurrence of new episodes [1].

In this case, we did not perform a histopathological examination, considering that the physical examination and history of shiitake consumption were sufficient for the diagnosis of FD. A complete remission of the lesions was seen within 7 days after the use of dexamethasone ointment in the affected area for 5 days.

Conclusion

With the expansion of Oriental cuisine to the West, there is an increased possibility for new cases of flagellate dermatitis induced by shiitake consumption. The dermatologist, in presence of linear eruptions on the trunk resembling whip lashes, should always ask the patient if shiitake was consumed, considering that the patient would probably fail in associating mushroom ingestion with flagellate dermatitis.

Statement of Ethics

The authors have no ethical conflicts to disclose.

Disclosure Statement

The authors have no conflicts of interest to declare.

There were no financial interests (stocks, patents, employment, honoraria, or royalties) or nonfinancial relationships (political, personal, or professional) that may be interpreted as having influenced the writing of this paper.

Funding Sources

Our article received no funding.

Author Contributions

All authors made substantial contributions to the conception and interpretation of the data for this work. All helped drafting the article and revising it critically for important intellectual content; gave their final approval of the submitted version; and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

References

- 1 Netchiporouk E, Pehr K, Ben-Shoshan M, Billick RC, Sasseville D, Singer M. Pustular flagellate dermatitis after consumption of shiitake mushrooms. *JAAD Case Rep*. 2015 May;1(3):117–9.
- 2 Poppe LM, Anders D, Kneitz H, Bröcker EB, Benoit S. Flagellate dermatitis caused by shiitake mushrooms. *An Bras Dermatol*. 2012 May-Jun;87(3):463–5.
- 3 Czarnecka AB, Kreft B, Marsch WC. Flagellate dermatitis after consumption of Shiitake mushrooms. *Postepy Dermatol Alergol*. 2014 Jun;31(3):187–90.
- 4 Nakamura T. Shiitake (*Lentinus edodes*) dermatitis. *Contact Dermatitis*. 1992 Aug;27(2):65–70.
- 5 Corazza M, Zauli S, Ricci M, Borghi A, Pedriali M, Mantovani L, et al. Shiitake dermatitis: toxic or allergic reaction? *J Eur Acad Dermatol Venereol*. 2015 Jul;29(7):1449–51.
- 6 Tan Q, Tan C. Log-grown shiitake is perhaps the real cause for Shiitake dermatitis. *J Eur Acad Dermatol Venereol*. 2016 Jan;30(1):197–8.
- 7 Aalto-Korte K, Susitaival P, Kaminska R, Mäkinen-Kiljunen S. Occupational protein contact dermatitis from shiitake mushroom and demonstration of shiitake-specific immunoglobulin E. *Contact Dermat*. 2005 Oct;53(4):211–3.
- 8 Grynszpan R, Niemeyer-Corbellini JP, Lopes MS, Ramos-e-Silva M. Bleomycin-induced flagellate dermatitis. *BMJ Case Rep*. 2013 Jun;2013 bcr2013009764.
- 9 Hamer S, Rabindranathnambi R. A wide-spread flagellate dermatitis. *BMJ Case Rep*. 2013 Jan;2013 bcr2012007682.
- 10 Loo HV, Oon HH. Flagellate dermatitis following consumption of shiitake mushroom. *Dermatol Rep*. 2011 Oct;3(2):e21.
- 11 Adriano AR, Acosta ML, Azulay DR, Quiroz CD, Talarico SR. Shiitake dermatitis: the first case reported in Brazil. *An Bras Dermatol*. 2013 May-Jun;88(3):417–9.

Fig. 1. Shiitake mushrooms.

Fig. 2. Linear lesions formed by grouped papules on the back.

Fig. 3. Lesions on the lateral trunk.

Fig. 4. Close-up of the lesions with better evidence of the grouped papules.