Prevention and Management of Chemotherapy-Induced Polyneuropathy

Berit Jordan^a Franziska Jahn^b Sandra Sauer^c Karin Jordan^c

^aDepartment of Neurology, University of Heidelberg, Heidelberg, Germany; ^bDepartment of Hematology and Oncology, University of Halle-Wittenberg, Halle (Saale), Germany; ^cDepartment of Hematology, Oncology and Rheumatology, University of Heidelberg, Heidelberg, Germany

Keywords

Chemotherapy-induced peripheral neurotoxicity · Neuropathic pain · Side effects · Duloxetine · Neurotoxicity

Abstract

Chemotherapy-induced peripheral neurotoxicity (CIPN) is a severe and common side effect caused by a variety of anti-neoplastic agents. Approximately 30–40% of patients treated with agents such as taxanes, vinca alkaloids, or platinum derivatives will develop CIPN. CIPN presents predominantly as a sensory axonal neuro(n)opathy with occasional motor and autonomic dysfunction exhibiting considerable variability of clinical symptoms ranging from mild tingling sensation to severe neuropathic pain. Typical symptoms include numbness (“minus symptom”), weakness, and abnormal gait as well as paresthesia and pain (“positive symptoms”). As CIPN symptoms potentially lead to long-term morbidity and can even aggravate after cessation of therapy, patients’ quality of life can be tremendously affected. In view of improved breast cancer survival outcomes, the late effects of CIPN are an unmet need in these patients. Therefore, early detection and assessment of first symptoms is important to effectively prevent severe CIPN. Therapeutic options for patients with CIPN are still limited, and pharmacological treatment focuses primarily on reduction or relief of neuropathic pain. CIPN is usually acutely managed by dose reduction or discontinuation of causative chemotherapy, potentially compromising treatment outcome. Currently, there is no causative proven therapy for the prevention of CIPN.

© 2019 S. Karger AG, Basel

Introduction

Peripheral neurotoxicity due to antineoplastic therapy is a common and often dose-limiting side effect. The risk of potential short- and long-term impairment of quality of life must be balanced against the benefits of cancer treatment. Over the years the term “CIPN” has been well established to describe primarily chemotherapy-induced peripheral neuropathy but should also be used to describe neurotoxicity deriving from newer targeted agents. Putative targets of neurotoxicity therefore include dorsal root ganglia (leading to so-called neuronopathy), axon and axonal components (myelin, microtubules, mitochondria, vascular network) as well as distal nerve terminals [1]. CIPN in breast cancer is often caused by taxanes, whereas nanoparticle albumin bound (nab)-paclitaxel, carboplatin, eribulin, and vinorelbine may also lead to significant CIPN. Faced with improved survival outcomes of early and metastatic breast cancer, research into improving the quality of survivorship – notably CIPN – is still an unmet need in these patients. This review gives an overview of CIPN and summarizes the latest recommendations on diagnosis, prevention, and treatment of peripheral neuropathy.

Incidence and Risk Factors

Incidence and severity of CIPN vary according to antineoplastic agents, cumulative dose, duration of exposure, scheduling, and combination of different agents.

Table 1. Incidence and therapy-associated risk factors for selected antineoplastic agents [1, 6]

Agent	Therapy-associated risk factors			
Cisplatin	Increased risk of neurotoxicity >300 mg/m ²			
Incidence	Single dose ¹	Cumulative dose ²	Infusion time	Duration of treatment
Grade 1–2 (14–63%)	Relevant	Relevant	Not relevant	Not relevant
Grade 3–4 (7–21%)				
Clinical symptoms	Predominantly sensory neuropathy			
Taxanes +/- carboplatin	Increased risk of neurotoxicity (docetaxel: >100 mg/m ² ; paclitaxel: 250 mg/m ²)			
Incidence	Single dose ¹	Cumulative dose ²	Infusion time ³	Duration of treatment
Grade 1–2 (20–50%)	Relevant	Relevant	Relevant	Not relevant
Grade 3–4 (6–20%)				
Clinical symptoms	Predominantly sensory neuropathy; in higher doses myalgia and myopathies possible			
Eribulin				
Incidence	Single dose	Cumulative dose	Infusion time	Duration of treatment
Grade 1–4 (~35%)	No data	No data	No data	No data
Grade 3–4 (~8%)				
Clinical symptoms	Sensory neuropathy, myalgia			

¹ Single dose: higher single doses carry the risk of increased neurotoxicity. ² Cumulative doses: higher cumulative doses increase the risk of neurotoxicity. ³ Prolonged infusion time seems to reduce the risk of neurotoxicity.

Remarkably, paclitaxel and carboplatin combination treatment is generally associated with similar neurotoxicity to paclitaxel alone [2]. Examples are shown in Table 1. Clinical symptoms of CIPN reflect a predominantly sensory axonal neuro(no)pathy with occasional motor and autonomic involvement in a predominant “glove and stocking” distribution. Typically, primarily sensory fibres are affected but some cytostatic agents also cause a sensorimotor pattern. Typical symptoms include numbness, paresthesia, lancinating pain, abnormal gait, and motor weakness. It is important to consider that CIPN may persist for many years beyond antineoplastic therapy and is associated with an increased risk of falls [3]. After completion of paclitaxel chemotherapy, approximately one-half of patients improve over a period of 4–6 months [4]. In this study, up to 80% of patients still had neuropathic symptoms up to 2 years following completion of treatment. Of these patients, 25% still had severe symptoms such as numbness in hands and feet.

Strong independent individual risk factors for developing CIPN are diabetes mellitus, age [5], and concurrent exposure to other neurotoxic agents as well as pre-existing neuropathy. Furthermore, alcohol abuse, metabolic diseases such as renal insufficiency, hypothyroidism, vitamin deficiency (e.g., B₁, B₆, B₁₂), and pre-existing hereditary neuropathies like CMT1A are considered to be risk factors [6]. The search for genetic markers carrying a high risk for developing CIPN has been unsuccessful for use in clinical practice so far [7].

Assessment and Diagnosis of CIPN

Symptoms of CIPN typically occur during the first 2 months of treatment, progress during active antineoplastic treatment, and then usually stabilize soon after treatment is completed. However, worsening of neuropathic symptoms after cessation of therapy called “coasting phenomenon” must be considered (e.g., paclitaxel). Coasting phenomenon is explained by drugs persisting in nerve axons after finishing therapy that lead to ongoing toxicity [8]. It should be acknowledged that there is considerable variability of symptoms depending on the agent used and the individual risk factors of the patient.

Typical clinical symptoms are predominantly sensory and usually include acral pain and paresthesia (tingling like pins and needles), accompanied by allodynia and hyperalgesia (“plus symptoms”). Sensory loss appears in a “glove and stocking type” distribution and leads to “minus” symptoms like numbness in hands and feet including impaired perception of light touch, vibration sense, and proprioception (tuning fork test) in clinical examination.

Vinca alkaloids and taxanes cause small fibre neuropathy, affecting the nerve terminals. As a result, patients experience the typical burning feet and hands and even lancinating pain. Decreased pain perception, hyperalgesia, and reduced temperature sensation are typical symptoms in clinical examination. Small fibre damage typically also shows autonomic involvement which may include abdominal pain, constipation, postural hypotension, disturbances of bladder, delayed gastric emptying, and reduced variability of heart rate.

Neurophysiological Examination

Nerve conduction studies and electromyography may be helpful in characterizing neuropathies (e.g., axonal vs. demyelinating) and provide additional information. However, diagnosis of CIPN is preferentially based on patients' complaints (patient-reported outcomes and grading scales) and neurological examination (functional measures). A progressive reduction in sensory nerve action potential amplitude (compound muscle action potential if motor fibres are affected) indicates axonal degeneration and is followed by impairment of nerve conduction velocity. Again, patients' symptoms and severity of CIPN during therapy cannot be adequately reflected using conventional nerve conduction parameters. Of note, in small fibre neuropathies all findings based on standard neurophysiological techniques are normal and only skin biopsy (gold standard) may demonstrate degeneration of small C (heat) and A δ (cold) fibres.

Assessment Tools Including Patient-Reported Outcome Measurements

Various tools are available, some for clinical trials and some for daily clinical practice. Domains included in the assessment tools are often narrow but deliver important information [9]. Many trials confirmed that patient-reported outcome measurements identified a higher incidence and severity of treatment-related toxicities than did clinician-reported outcome measurements [10].

The typical example for a clinician-reported outcome measurement is represented by the National Cancer Institute Common Terminology Criteria for Adverse Events (NCI-CTCAE). This assessment still remains the most commonly utilized tool. Cavaletti's Total Neuropathy Score clinical version (TNSc) can be used with good inter-rater reliability. As such, the TNS can detect very early changes occurring in the peripheral nervous system even at subclinical level.

Patient-reported outcome measurement assessment includes for example, the EORTC CIPN20 and the Fact-GOG-NTX12.

EORTC CIPN20. The CIPN20 is a 20-item quality of life questionnaire, which has been developed to elicit patients' experience of symptoms and functional limitations related to CIPN. The CIPN20 has three subscales: a sensory, motor, and autonomic subscale.

Fact-GOG-NT. This 12-question tool requests the user to circle or mark one number per line to indicate the response as it applies to the past 7 days. The forms can be filled out by either patient or caregiver while at the clinic.

This type of tool also gives caregivers a chance to report their observations if the patient is not able to converse easily or does not want to report something that may affect the therapy [1].

In order to address symptomatic differences between chemotherapeutic agents (e.g., cisplatin and taxanes), various chemotherapy-specific scales have also been developed [11]. FACT-Taxane consists of both FACT-General and an added taxane subscale. The subscale has 16 items, including an 11-item neurotoxicity subscale and 5 additional taxane-specific questions related to the effects of arthralgia, myalgia, and skin changes.

Prevention of CIPN

Pharmacological Intervention

To date, no suitable drug has been found to effectively prevent CIPN pharmacologically. Use of acetyl-L-carnitine, acetylcystein, α -lipoic acid, amifostine, amitriptyline, calcium/magnesium, carbamazepine, diethyldithiocarbamate, glutathione, gosha-jinki-gan, minocycline, nimodipine, omega-3 fatty acids, vitamin B, and also vitamin E has been investigated in the prevention of peripheral neuropathy in clinical trials. However, none of these has yet led to positive results [12–24]. Results, however, must be interpreted with caution: trials in this area are often underpowered and heterogeneous.

Nurse-Led Intervention

The efficacy of a nurse-led care program in improving quality of life outcomes in patients receiving adjuvant chemotherapy for breast cancer was investigated in a randomized trial (routine hospital care vs. both hospital and nurse-led care). The intervention included regular phone consultations every 1–2 weeks. The primary endpoint of difference in FACT-General score was surprisingly not reached. However various important secondary endpoints were reached. In summary, this study showed the complexity of factors that contribute to CIPN, and underline the potential role of non-pharmacological interventions such as modulation of pain perception of CIPN [25].

Cryotherapy/Compression Therapy

For non-pharmacological prevention of CIPN, cryotherapy with frozen socks or gloves seems to be effective, although evidence is still scarce. In a pilot trial with 20 breast cancer patients receiving weekly paclitaxel one limb was cooled, and the contralateral limb acted as a control. The hypothermia process reduces skin temperature by $1.5 \pm 0.7^\circ\text{C}$. The amount of skin cooling was significantly associated with preservation of motor nerve amplitude after 6 months ($p < 0.0005$) [26]. Another study with 40 patients confirms these observa-

tions: when wearing frozen gloves and socks on the dominant side for 90 min during paclitaxel infusion, patients had diminished objective and subjective symptoms of CIPN as compared to the non-dominant, untreated side [27]. Larger studies still need to confirm these findings, and in particular practicability of the treatment as frozen socks and gloves can also cause significant discomfort.

Another protective effect may be noted for compression therapy using surgical gloves. In a small study with 42 patients receiving nab-paclitaxel, patients acted as their own control wearing surgical gloves on one hand and leaving the other hand ungloved. The gloved hand was compressed using gloves a size smaller. Less neuropathy was seen in the gloved versus the ungloved hand [28]. Compression therapy pursues the same approach as hypothermia by reducing blood flow in cold tissue as the fingertips of the glove-protected hand had a significant lower temperature than the ungloved hand.

Exercise

Early studies suggest a protective effect of exercise and functional training on CIPN, therefore it is advisable to encourage patients – if there are no contraindications – to strengthen their muscles and improve sensorimotor function [6].

Acupuncture (Prevention and Therapy)

Studies investigating the effectiveness of acupuncture to prevent or treat CIPN are often underpowered and without a control group and therefore not sufficient to support or refute its use. In a phase IIa single-arm clinical trial, acupuncture was useful for 27 breast cancer patients, who developed grade 2 CIPN due to weekly paclitaxel to prevent escalation of CIPN to higher grades (NCI-CTCAE, Version 4.0) [29]. Another single-arm observational study with only 10 patients with breast cancer showed effectiveness in reducing taxane-induced peripheral neuropathic pain [30]. In both studies, acupuncture was well tolerated, and patients reported no adverse events. However, study results are inconclusive (also in other patient populations) and do not allow a positive recommendation.

Treatment of CIPN

Efficacious therapeutic options for patients with established CIPN are still limited. Special attention should therefore be paid to the onset and severity of the symptoms during the antineoplastic treatment in order to reduce the dose if necessary or to change antineoplastic therapy. If symptoms and functional limitations have occurred, detailed patient advice is necessary and effective.

Behaviourally Based Techniques

To reduce pain deriving from CIPN, the self-guided online cognitive and behaviourally based pain management intervention PROSPECT (Proactive Self-Management Program for Effects of Cancer Treatment) was successful in a pilot trial with 60 patients (mean age >60 years) [31]. In a randomized setting, patients received PROSPECT or standard care over 8 weeks. Patients in the intervention (PROSPECT) group had greater improvements in “worst” pain than patients with standard care alone.

Exercise

Physical exercise is often applied in multimodal settings; therefore, many patients receive this as part of their standard treatment. Owing to this fact, there is a lack of strong scientific evidence about its specific usefulness in the treatment of CIPN. Nevertheless, the number of studies showing the effectiveness of physical exercise and functional training (e.g., vibration training) is increasing [32, 33]. To improve coordination, sensorimotor function, and fine motor function the training should begin with the onset of manifest CIPN at the latest. It would be advisable to initiate exercise at the time potentially neurotoxic cancer treatment is initiated [6]. Improvement of physical function is particularly important in the prevention of malnutrition and improvement of physical function in order to prevent disability and falls, especially in elderly patients [3].

Pharmacological Treatment

As there are no pharmacological interventions to reduce numbness, paresthesia, or motor weakness, treatment for patients with chronic CIPN focuses on reduction or relief of neuropathic pain (so-called plus symptoms).

In a cross-over study by Smith et al. [34], duloxetine, a selective serotonin reuptake inhibitor, showed moderate clinical benefits in patients with painful CIPN. A total of 231 patients treated with platinum derivatives or taxanes with grade ≥ 1 neuropathy (according to NCI-CTCAE scale) and at least 4 points on a neuropathic pain scale of 0–10 points received duloxetine or placebo. More patients with duloxetine reported a significant decrease in neuropathic pain (59 vs. 38%), noting that the effect of duloxetine on platinum-induced neuropathic pain was greater than the effect of taxane-induced neuropathic pain.

If duloxetine treatment has failed or is not indicated, anticonvulsants or tricyclic antidepressants may have the potential for symptom control in patients with CIPN. However, evidence specifically in CIPN is still limited. As such, the treatment approach is analogous to usage of these drugs in other forms of neuropathic pain.

Mishra et al. [35] investigated the efficacy of amitriptyline, gabapentin, pregabalin or placebo in 120 cancer patients with neuropathic pain. In all 4 groups a signifi-

cant reduction of the pain level (VAS 100 mm) was achieved after the intervention. The most pronounced improvement of neuropathic pain was obtained in the pregabalin arm (compared to amitriptyline $p = 0.003$, gabapentin $p = 0.042$, and placebo $p = 0.042$).

In a study with 131 patients receiving lamotrigine or placebo after or during treatment with taxanes, vinca alkaloids, or platinum derivatives, no improvement in neuropathic pain could be shown. However, the lamotrigine arm detected a noticeably high dropout rate [36]. Opioids may be also useful to relieve neuropathic pain, but this has only been proven for neuropathic pain from causes other than antineoplastic therapies [37, 38].

Topical local interventions may be helpful, particularly in small fibre affection of peripheral nerves. Substantial pain relief in 31 out of 38 evaluable patients was shown in a phase II trial using 1% menthol cream to the affected area [39]. Since menthol cream has no noted side effects and is inexpensive, it could be worth investigating its use further, although randomized data are lacking. Non-CIPN studies showed benefit using 8% capsaicin-containing patches in patients with painful neuropathies [40] and attempts were made to try this also in patients with CIPN [38]. A topical 4% amitriptyline/2% ketamine preparation was studied in 462 patients in an RCT showing that neuropathic pain, numbness, or tingling could not be alleviated [41].

It must be acknowledged, however, that perception of neuropathic pain may be aggravated by sleep disturbance, anxiety, depression, and central sensitization of pain, a holistic treatment concept is therefore mandatory.

Conclusion

Considering the increasing prevalence of cancer, wider scope of chemotherapy drugs, and long-term side effects, patients with polyneuropathy will present with many issues and not only physical findings. In view of the large scale of patient numbers and prevalence, polyneuropathy as an important side effect of antineoplastic treatments must be addressed as a so far unmet need. Quality of life issues and rehabilitation concepts for long-term functional deficits will be future directions for research.

Acknowledgements

The authors would like to thank Dr. Camilla Leithold and PD Dr. Alexandra Jensen for the editorial assistance during the preparation of the manuscript.

Disclosure Statement

Berit Jordan: advisory board and or honoraria for presentations for Alexion Pharmaceuticals, Temmler (Hormosan Pharma), Novartis, BioMarin, Merz Pharmaceuticals, Ipsen Pharma, and Allergan Anita Margulies. No financial disclosures.

Franziska Jahn: advisory board and or honoraria for presentations for MSD, Amgen, Riemser, Tesaro, Pomme-med, and Gyn-Onko Update. No financial disclosures.

Sandra Sauer: no financial disclosures.

Karin Jordan: financial interests, honoraria for speaker, consultancy or advisory role, royalties, direct research funding: MSD, Merck, Amgen, Hexal, Riemser, Helsinn, Tesaro, Kreussler, Voluntas, Pfizer, Pomme-med, Pharma Mar, Prime Oncology, Onko Update, Annals of Oncology, and UpToDate.

References

- 1 Park SB, Goldstein D, Krishnan AV, Lin CS, Friedlander ML, Cassidy J, et al. Chemotherapy-induced peripheral neurotoxicity: a critical analysis. *CA Cancer J Clin*. 2013 Nov-Dec; 63(6):419–37.
- 2 Kandula T, Farrar MA, Kiernan MC, Krishnan AV, Goldstein D, Horvath L, et al. Neuropathological and clinical outcomes in chemotherapy-induced neuropathy in cancer. *Clin Neurophysiol*. 2017 Jul;128(7):1166–75.
- 3 Winters-Stone KM, Horak F, Jacobs PG, Trubowitz P, Dieckmann NF, Stoyles S, et al. Falls, Functioning, and Disability Among Women With Persistent Symptoms of Chemotherapy-Induced Peripheral Neuropathy. *J Clin Oncol*. 2017 Aug;35(23):2604–12.
- 4 Hershman DL, Weimer LH, Wang A, Krawinkel G, Brafman L, Fuentes D, et al. Association between patient reported outcomes and quantitative sensory tests for measuring long-term neurotoxicity in breast cancer survivors treated with adjuvant paclitaxel chemotherapy. *Breast Cancer Res Treat*. 2011 Feb; 125(3):767–74.
- 5 Hershman DL, Till C, Wright JD, Awad D, Ramsey SD, Barlow WE, et al. Comorbidities and Risk of Chemotherapy-Induced Peripheral Neuropathy Among Participants 65 Years or Older in Southwest Oncology Group Clinical Trials. *Comorbidities and Risk of Chemotherapy-Induced Peripheral Neuropathy Among Participants 65 Years or Older in Southwest Oncology Group Clinical Trials*. *J Clin Oncol*. 2016 Sep;34(25): 3014–22.
- 6 Jordan K, Feyer P, Höller U, Link H, Wörmann B, Jahn F. Supportive Treatments for Patients with Cancer. *Dtsch Arztebl Int*. 2017 Jul;114(27-28):481–7.
- 7 Argyriou AA, Bruna J, Genazzani AA, Cavaletti G. Chemotherapy-induced peripheral neurotoxicity: management informed by pharmacogenetics. *Nat Rev Neurol*. 2017 Aug;13(8):492–504.
- 8 Staff NP, Grisold A, Grisold W, Windebank AJ. Chemotherapy-induced peripheral neuropathy: A current review. *Ann Neurol*. 2017 Jun;81(6):772–81.
- 9 Paice JA. Clinical challenges: chemotherapy-induced peripheral neuropathy. *Semin Oncol Nurs*. 2009 May;25(2 Suppl 1):S8–19.
- 10 Jordan K, Aapro M, Kaasa S, Ripamonti CI, Scotté F, Strasser F, et al. European Society for Medical Oncology (ESMO) position paper on supportive and palliative care. *Ann Oncol*. 2018 Jan;29(1):36–43.
- 11 Cella D, Peterman A, Hudgens S, Webster K, Socinski MA. Measuring the side effects of taxane therapy in oncology: the functional assesment of cancer therapy-taxane (FACT-taxane). *Cancer*. 2003 Aug;98(4): 822–31.
- 12 Hershman DL, Lacchetti C, Dworkin RH, Lavoie Smith EM, Bleeker J, Cavaletti G, et al.; American Society of Clinical Oncology. Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: american Society of Clinical Oncology clinical practice guideline. *J Clin Oncol*. 2014 Jun;32(18):1941–67.

- 13 Guo Y, Jones D, Palmer JL, Forman A, Dakhil SR, Velasco MR, et al. Oral alpha-lipoic acid to prevent chemotherapy-induced peripheral neuropathy: a randomized, double-blind, placebo-controlled trial. *Support Care Cancer*. 2014 May;22(5):1223–31.
- 14 von Delius S, Eckel F, Wagenpfeil S, Mayr M, Stock K, Kullmann F, et al. Carbamazepine for prevention of oxaliplatin-related neurotoxicity in patients with advanced colorectal cancer: final results of a randomised, controlled, multicenter phase II study. *Invest New Drugs*. 2007 Apr;25(2):173–80.
- 15 Gandara DR, Nahhas WA, Adelson MD, Lichtman SM, Podczaski ES, Yanovich S, et al. Randomized placebo-controlled multicenter evaluation of diethyldithiocarbamate for chemoprotection against cisplatin-induced toxicities. *J Clin Oncol*. 1995 Feb;13(2):490–6.
- 16 Leal AD, Qin R, Atherton PJ, Haluska P, Behrens RJ, Tiber CH, et al.; Alliance for Clinical Trials in Oncology. North Central Cancer Treatment Group/Alliance trial N08CA—the use of glutathione for prevention of paclitaxel/carboplatin-induced peripheral neuropathy: a phase 3 randomized, double-blind, placebo-controlled study. *Cancer*. 2014 Jun;120(12):1890–7.
- 17 Schloss JM, Colosimo M, Airey C, Masci P, Linnane AW, Vitetta L. A randomised, placebo-controlled trial assessing the efficacy of an oral B group vitamin in preventing the development of chemotherapy-induced peripheral neuropathy (CIPN). *Support Care Cancer*. 2017 Jan;25(1):195–204.
- 18 Hershman DL, Unger JM, Crew KD, Till C, Greenlee H, Minasian LM, et al. Two-Year Trends of Taxane-Induced Neuropathy in Women Enrolled in a Randomized Trial of Acetyl-L-Carnitine (SWOG S0715). *J Natl Cancer Inst*. 2018 Jun;110(6):669–76.
- 19 Albers JW, Chaudhry V, Cavaletti G, Donehower RC. Interventions for preventing neuropathy caused by cisplatin and related compounds. *Cochrane Database Syst Rev*. 2014 Mar;(3):CD005228.
- 20 Kautio AL, Haanpää M, Leminen A, Kalso E, Kautiainen H, Saarto T. Amitriptyline in the prevention of chemotherapy-induced neuropathic symptoms. *Anticancer Res*. 2009 Jul;29(7):2601–6.
- 21 Jordan B, Jahn F, Beckmann J, Unverzagt S, Müller-Tidow C, Jordan K. Calcium and Magnesium Infusions for the Prevention of Oxaliplatin-Induced Peripheral Neurotoxicity: A Systematic Review. *Oncology*. 2016;90(6):299–306.
- 22 Kuriyama A, Endo K. Goshajinkigan for prevention of chemotherapy-induced peripheral neuropathy: a systematic review and meta-analysis. *Support Care Cancer*. 2018 Apr;26(4):1051–9.
- 23 Pachman DR, Dockter T, Zekan PJ, Fruth B, Ruddy KJ, Ta LE, et al. A pilot study of minocycline for the prevention of paclitaxel-associated neuropathy: ACCRU study RU221408I. *Support Care Cancer*. 2017 Nov;25(11):3407–16.
- 24 Ghoreishi Z, Esfahani A, Djazayeri A, Djalali M, Golestan B, Ayromlou H, et al. Omega-3 fatty acids are protective against paclitaxel-induced peripheral neuropathy: a randomized double-blind placebo controlled trial. *BMC Cancer*. 2012 Aug;12(1):355.
- 25 Lai XB, Ching SS, Wong FK, Leung CW, Lee LH, Wong JS, et al. A Nurse-Led Care Program for Breast Cancer Patients in a Chemotherapy Day Center: A Randomized Controlled Trial. *Cancer Nurs*. 2019 Jan/Feb;42(1):20–34.
- 26 Sundar R, Bandla A, Tan SS, Liao LD, Kumarakulasinghe NB, Jeyasekharan AD, et al. Limb Hypothermia for Preventing Paclitaxel-Induced Peripheral Neuropathy in Breast Cancer Patients: A Pilot Study. *Front Oncol*. 2017 Jan;6:274.
- 27 Hanai A, Ishiguro H, Sozu T, Tsuda M, Yano I, Nakagawa T, et al. Effects of Cryotherapy on Objective and Subjective Symptoms of Paclitaxel-Induced Neuropathy: Prospective Self-Controlled Trial. *J Natl Cancer Inst*. 2018 Feb;110(2):141–8.
- 28 Tsuyuki S, Senda N, Kanng Y, Yamaguchi A, Yoshibayashi H, Kikawa Y, et al. Evaluation of the effect of compression therapy using surgical gloves on nanoparticle albumin-bound paclitaxel-induced peripheral neuropathy: a phase II multicenter study by the Kamigata Breast Cancer Study Group. *Breast Cancer Res Treat*. 2016 Nov;160(1):61–7.
- 29 Bao T, Seidman AD, Piulson L, Vertosick E, Chen X, Vickers AJ, et al. A phase IIA trial of acupuncture to reduce chemotherapy-induced peripheral neuropathy severity during neoadjuvant or adjuvant weekly paclitaxel chemotherapy in breast cancer patients. *Eur J Cancer*. 2018 Sep;101:12–9.
- 30 Jeong YJ, Kwak MA, Seo JC, Park SH, Bong JG, Shin IH, et al. Acupuncture for the Treatment of Taxane-Induced Peripheral Neuropathy in Breast Cancer Patients: A Pilot Trial. *Evid Based Complement Alternat Med*. 2018 Oct;2018:5367014.
- 31 Knoerl R, Smith EM, Barton DL, Williams DA, Holden JE, Krauss JC, et al. Self-Guided Online Cognitive Behavioral Strategies for Chemotherapy-Induced Peripheral Neuropathy: A Multicenter, Pilot, Randomized, Wait-List Controlled Trial. *J Pain*. 2018 Apr;19(4):382–94.
- 32 Kleckner IR, Kamen C, Gewandter JS, Mohile NA, Heckler CE, Culakova E, et al. Effects of exercise during chemotherapy on chemotherapy-induced peripheral neuropathy: a multicenter, randomized controlled trial. *Support Care Cancer*. 2018 Apr;26(4):1019–28.
- 33 Schönsteiner SS, Bauder Mißbach H, Benner A, Mack S, Hamel T, Orth M, et al. A randomized exploratory phase 2 study in patients with chemotherapy-related peripheral neuropathy evaluating whole-body vibration training as adjunct to an integrated program including massage, passive mobilization and physical exercises. *Exp Hematol Oncol*. 2017 Feb;6(1):5.
- 34 Smith EM, Pang H, Cirrincione C, Fleishman S, Paskett ED, Ahles T, et al.; Alliance for Clinical Trials in Oncology. Effect of duloxetine on pain, function, and quality of life among patients with chemotherapy-induced painful peripheral neuropathy: a randomized clinical trial. *JAMA*. 2013 Apr;309(13):1359–67.
- 35 Mishra S, Bhatnagar S, Goyal GN, Rana SP, Upadhyay SP. A comparative efficacy of amitriptyline, gabapentin, and pregabalin in neuropathic cancer pain: a prospective randomized double-blind placebo-controlled study. *Am J Hosp Palliat Care*. 2012 May;29(3):177–82.
- 36 Rao RD, Flynn PJ, Sloan JA, Wong GY, Novotny P, Johnson DB, et al. Efficacy of lamotrigine in the management of chemotherapy-induced peripheral neuropathy: a phase 3 randomized, double-blind, placebo-controlled trial, N01C3. *Cancer*. 2008 Jun;112(12):2802–8.
- 37 Finnerup NB, Attal N, Haroutounian S, McNicol E, Baron R, Dworkin RH, et al. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol*. 2015 Feb;14(2):162–73.
- 38 Cavaletti G, Marmiroli P. Pharmacotherapy options for managing chemotherapy-induced peripheral neurotoxicity. *Expert Opin Pharmacother*. 2018 Feb;19(2):113–21.
- 39 Fallon MT, Storey DJ, Krishan A, Weir CJ, Mitchell R, Fleetwood-Walker SM, et al. Cancer treatment-related neuropathic pain: proof of concept study with menthol—a TRPM8 agonist. *Support Care Cancer*. 2015 Sep;23(9):2769–77.
- 40 Simpson DM, Robinson-Papp J, Van J, Stoker M, Jacobs H, Snijder RJ, et al. Capsaicin 8% Patch in Painful Diabetic Peripheral Neuropathy: A Randomized, Double-Blind, Placebo-Controlled Study. *J Pain*. 2017 Jan;18(1):42–53.
- 41 Gewandter JS, Mohile SG, Heckler CE, Ryan JL, Kirshner JJ, Flynn PJ, et al. A phase III randomized, placebo-controlled study of topical amitriptyline and ketamine for chemotherapy-induced peripheral neuropathy (CIPN): a University of Rochester CCOP study of 462 cancer survivors. *Support Care Cancer*. 2014 Jul;22(7):1807–14.