Pathophysiology of Diabetic Retinopathy: Contribution and Limitations of Laboratory Research

Timothy S. Kern^{a, b} David A. Antonetti^c Lois E.H. Smith^d

^aGavin Herbert Eye Institute, University of California Irvine, Irvine, CA, USA; ^bVeterans Administration Medical Center Research Service 151, Cleveland, OH, USA; ^cKellogg Eye Center, Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, MI, USA; ^dDepartment of Ophthalmology, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA

Keywords

Diabetic retinopathy · Laboratory research · Preclinical models

Abstract

Preclinical models of diabetic retinopathy are indispensable in the drug discovery and development of new therapies. They are, however, imperfect facsimiles of diabetic retinopathy in humans. This chapter discusses the advantages, limitations, and physiological and pathological relevance of preclinical models of diabetic retinopathy. The judicious interpretation and extrapolation of data derived from these models to humans and a correspondingly greater emphasis placed on translational medical research in early-stage clinical trials are essential to more successfully inhibit the development and progression of diabetic retinopathy in the future.

© 2019 S. Karger AG, Basel

It is generally accepted that animal models do not perfectly represent human conditions, and this is true also for diabetic retinopathy. Thus, a reasonable question is how closely laboratory models of diabetic retinopathy

mimic human disease and how that influences our willingness to use and trust data from such animal models. Animal models do have important advantages in providing mechanistic insight and when assessing potential new treatments with unknown long-term adverse effects. However, when a proposed treatment is known to be essentially free of safety concerns or adverse effects (such as an off-label use of an FDA approved drug), it can be validly questioned as to why laboratory animal models should be used.

Specifically with respect to diabetic retinopathy, the deficiencies of most species of small animals used for laboratory research include the apparent failure to develop diabetes-induced preretinal neovascularization and clinically meaningful stages of retinopathy, the lack of a macula, and the apparent failure of most models to develop diabetes-induced retinal/macular thickening and edema. Other potential weaknesses compared to humans include the fundamental difference that most rodents are nocturnal, have a short life span (which affects the development of disease severity), have a very small retina, lack many of the lifestyle factors that can adversely affect human disease, and are unable to communicate about what they are or are not seeing (visual function).

Fig. 1. Vascular histopathology of diabetic retinopathy progression is similar between species, differing mainly in the severity of retinopathy that they develop (likely due at least in part to differences in life span). Diabetic patients can develop substantial microvascular pathology, including microaneurysms and degenerate capillaries, and in some cases, preretinal neovascularization. Similar lesions develop in large animal models of diabetic retinopathy, except that they have not been observed to develop preretinal neo-

vascularization in the 5+ years of diabetes that they have been studied. Rodents develop predominantly only degenerate capillaries (large arrow) and pericyte ghosts (small arrow) and basement membrane thickening (not shown) in their limited lifetime. The duration of time needed for the various stages of the retinopathy in the different species are indicated in parentheses. PDR, proliferative diabetic retinopathy.

Animal models do have important advantages, however. They require much less time and effort to obtain results than human clinical trials and are considerably less expensive to conduct. Those potential advantages are not meaningful, however, unless the disease processes leading to retinopathy in animals are generally similar to those in diabetic patients.

The available evidence suggests that the steps along the progression of diabetic retinopathy in the animal species tested are similar to humans, except that the animals do not progress as far along those steps as the longer-lived humans (Fig. 1). Capillary degeneration, loss of pericytes, thickening of vascular basement membranes, as well as functional changes including changes in blood flow, visual function, and vascular permeability develop in dia-

betic rodents, just as they do in longer-lived diabetic dogs and even longer-lived diabetic primates and patients. Larger and longer-lived species (dogs [1–3], cats [4], primates [5, 6], and humans) progress beyond those early lesions and also develop microaneurysms, cotton-wool spots, IRMA, and at least in diabetic dogs [7] and primates [8, 9], also retinal edema and/or intraretinal neovascularization. To date, only humans have been found to develop preretinal neovascularization due to diabetes, typically after durations of diabetes that are considerably longer than can be studied in any animal species.

Have animal studies of diabetic retinopathy been reliable at predicting the response of diabetic patients to disease or therapy? Animal models have been studied using similar therapies as diabetic patients in only a limited

number of cases (in large part due to the extraordinary cost of clinical trials related to diabetic retinopathy), but some comparisons can be made. A good example of animal studies predicting the response of diabetic patients to therapy is the demonstration of the beneficial effect of glycemic control on diabetic retinopathy in 5-year studies of diabetic dogs, which predated the later demonstration of a similar benefit of glycemic control in diabetic patients by more than a decade [10]. Studies in dogs [11, 12] and then also rats [13, 14] demonstrated a resistance of retinopathy to arrest with treatment once retinopathy had been initiated (later known as metabolic memory) that was found also in humans [15]. Studies in diabetic dogs [16] as well as diabetic patients [17] also showed no beneficial effect of an aldose reductase inhibitor to inhibit initiation or progression of the microaneurysms and capillary degeneration that characterize diabetic retinopathy. However, results related to aldose reductase and retinopathy differed from human findings in diabetic mice [18, 19] or some [20–22] (but not all [16, 23]) galactose-feeding studies. Administration of aspirin was reported to inhibit the development or progression of diabetic retinopathy in dogs and rats [24, 25] and in one study of diabetic patients [26], but not in another 2-year clinical study (Early Treatment of Diabetic Retinopathy; ETDRS) [27]. This difference might have been due to experimental design, because the animal studies showing a beneficial effect of the aspirin used higher (potentially anti-inflammatory) doses of the drug than the ETDRS study, which used only a lower (antiplatelet) dose.

All-in-all, the evidence suggests that at least dogs and perhaps other long-lived mammals have shown value in terms of predicting the outcome of similar studies in diabetic patients, suggesting that the pathogenesis of retinopathy in those species is similar to that in human patients.

The evidence that rodents show the same predictive ability with respect to the effect of a therapy on the development of retinal histopathology is less compelling, but there is evidence that diabetes induces abnormalities in retinal vascular permeability and visual function like those seen in diabetic patients. Also, molecular abnormalities in the retina are shared in common between diabetic patients and diabetic rodents. Moreover, diabetes-induced alterations in visual function (which has not been measured to date in large animals) becomes impaired in diabetic rodents [28, 29], as it does in diabetic patients, providing some evidence that rodent animal models also offer a meaningful opportunity to investigate the pathogenesis and treatment of retinopathy that develops in diabetic patients.

A significant value of animal models is their use to clarify molecular mechanisms by which diseases progress and to provide insight into molecular mechanisms underlying effects of a therapy. Accordingly, what have animal studies revealed about the pathogenesis and pathophysiology of diabetic retinopathy? Initial studies using experimental hyperglycemia (such as with fructose [30, 31] or galactose feeding [16, 32–36]) clearly showed that hyperglycemia (as opposed to insulin resistance or deficiency) plays a major role in the pathogenesis of retinopathy in a way that could not be demonstrated in diabetic patients. Beyond that, many biochemical abnormalities have been detected in retinas of diabetic animals, presumably developing as a result of abnormalities in substrate availability (glucose and lipids) and hormonal abnormalities (insulin, growth hormone) acting systemically as well as on the retina [37–42], and animal models clearly are central in elucidating the potential importance of these abnormalities in the development of retinopathy. In addition, the development of retinal vasculature after birth in the mouse combined with the development of endothelial specific Cre recombinases has provided an unprecedented opportunity to explore retinal vascular development and differentiation, thus providing the mechanistic framework from which therapies targeting vessel growth and remodeling have been made. Additional studies into the mechanisms of retinal function, signal transduction, and differentiation continue to provide novel insight.

Probably the most impressive success story in the past decades with regard to diabetic retinopathy has been the remarkable effects of anti-VEGF (vascular endothelial growth factor) therapies to inhibit retinal neovascularization and reverse macular edema in diabetes and other diseases. These therapies currently are the only approved intervention for existing advanced diabetic retinopathy. Clinical research played a large role in identifying how best to use this therapeutic approach, but animal research played a huge role in identifying and characterizing VEGF and demonstrating the relationship between VEGF and ischemia [43–48]. These therapies were developed largely as anticancer drugs but were found to be effective in models of neovascularization in mice and were supported by years of angiogenesis research including gene deletion studies and the use of VEGF-binding antibodies and soluble receptors [49].

A previously mentioned proof-of-principle study was the demonstration of the beneficial effect of good glycemic control on the development of diabetic retinopathy prior to comparable convincing data in humans. A good

example of animal studies demonstrating the molecular mechanisms underlying a therapy is the work that was done to understand the unexpected finding that diabetic retinopathy was significantly inhibited in patients treated with fibrates [50, 51]. The mechanism of effect was judged not to be via lowering of lipids (as originally assumed), and subsequent animal studies demonstrated a variety of actions of fibrates on metabolism and transcription factors as well as antioxidant and anti-inflammatory actions [52–56]. Animal models are providing insight also into the role of photoreceptors in the pathogenesis of both the early and late stages of diabetic retinopathy (in part via release of soluble factors [57–61]), which follows from the clinical recognition that diabetic patients with retinitis pigmentosa seemed protected from the development of diabetic retinopathy [62].

For many years, many therapeutic studies had been conducted in animal models with the assumption that a single (unidentified) molecular defect or pathway occurring in the retina in diabetes was a major cause of retinopathy and that inhibition of that abnormality or pathway would lead to inhibition of retinopathy [16, 24, 25, 63–77]. Indeed, a large number of animal studies using systemic pharmacologic therapies or genetically modified animals have demonstrated impressive inhibition of diabetes-induced abnormalities in retinal function or capillary permeability or degeneration at a particular duration of diabetes.

Should confidence in animal models be reduced by the fact that many experimental therapies administered to animal models reportedly show a beneficial effect on lesions of retinopathy compared to the paucity of beneficial therapies clinically to inhibit early stages of retinopathy? Caution is obviously warranted since most studies neither tested the longevity of the beneficial effect nor the effect of the therapy on the spectrum of other lesions or abnormalities that are part of clinically defined “diabetic retinopathy.” On the other hand, perhaps this can be explained by the possibility that preclinical researchers now are testing therapies that are closer to the actual molecular mechanism by which retinopathy can be inhibited than we were previously. An example of this is the strong research focus on anti-inflammatory approaches to inhibit retinopathy, which is supported by the clearly beneficial effect on DME by corticosteroids in diabetic patients.

Very important insight has been provided recently by increasing evidence that essentially all retinal cell types become abnormal in diabetes and that many of those cells participate in the development of diabetic retinopathy. For example, deletion of VEGF only from Muller cells

markedly inhibits diabetes-induced abnormalities in retinal vascular function [78] and permeability [79]. Likewise, inhibiting phototransduction in photoreceptors [60, 80] or slowing visual cycle activity by inhibiting RPE65 activity in retinal pigment epithelium [81] significantly inhibits the diabetes-induced increase in capillary leakage and degeneration. Even cells that are not part of the retina, such as leukocytes [82, 83] and stem/progenitor cells [84], now are known to participate in the development of the vascular complications of diabetic retinopathy or at least to fail to repair lesions that are developing due to diabetes. Evidence also suggests that multiple pathogenic processes are involved in the pathogenesis of retinopathy, and those signaling pathways can persist, converge, and change as glycemia fluctuates and probably also change during the different stages of retinopathy.

Considering the myriad of different functions of those various cell types and the different signaling pathways that are active in those different cells, the concept that the inhibition of a single pathway should be adequate to inhibit the development of retinopathy seems much less promising than it once did. Proteomic and transcriptomic studies have indicated that single therapies administered to diabetic mice corrected only a fraction of the molecular defects in the retina due to diabetes [85]. Thus, combinations of drugs that therapeutically modulate multiple mechanistic nodes within multiple cells or stabilize networks between different signaling pathways may enable better outcomes than treatment with a single drug. Such approaches now are being tested for diabetic macular edema, but there is a strong rationale also for extending this to preclinical studies attempting to inhibit the earlier stages of retinopathy [29].

Modern “-omic” profiling can enhance the preclinical development of therapies for the treatment of diabetic retinopathy or other diseases by providing a global assessment of the alterations in signaling systems caused by the disease and by providing a quantitative and unbiased assessment of how well a therapy turns that abnormal cell landscape to normal. The presumption of such an approach is that the therapies that best restore the transcriptome or other systems to normal are likely to be the best inhibitors of retinopathy. Such an unbiased approach to therapy does not require prior knowledge (or bias) about which pathway or cell type(s) are involved or dominant in the progression of disease.

Animal models have provided tremendous insight into the pathogenesis of diabetic retinopathy to date and are expected to continue to do so in the future. Nevertheless, there are steps that can be taken to make animal pre-

clinical studies more relevant to clinical research and care and thus, more useful.

One problem that impairs the clinical translation of results from animal studies of diabetic retinopathy (and other diseases) is that the endpoints used in animal studies are not the same as those used in clinical studies. Thus, it has been difficult or impossible to verify that the effects of therapy in preclinical animal studies are (or are not) occurring also in similar clinical studies. For example, both clinicians and basic researchers agree that degeneration of retinal capillaries in diabetes is a powerful stimulus for retinal ischemia and eventual development of preretinal neovascularization. However, most animal studies evaluating the effect of a therapy on that retinal histopathology evaluate and quantitate capillary degeneration microscopically at a very high resolution after removal of the neural retina ("trypsin digest" method), whereas clinical evaluation of capillary degeneration in patients has been graded at low resolution using fluorescein angiography until now. New techniques are now available that can provide the same information in animals and in patients, and incorporation of optical coherence tomography (OCT), OCT-angiography (OCTA), 2-photon microscopy, and magnetic resonance imaging should allow direct comparison between the animal and patient studies using common techniques.

Another important difference between clinical and animal studies of diabetic retinopathy is that clinical studies place a priority on preserving or regaining vision in diabetes, but this has not been incorporated in many animal studies. Since there now is recognition that both vascular

and neural abnormalities participate in the spectrum of abnormalities that make up diabetic retinopathy, animal studies need to measure both retinal histopathology and visual function, not one or the other. Finally, animal studies demonstrating reproducibility across different laboratories and in various species, multiple interventions targeting a common pathway yielding similar results, and genetic data combined with therapeutic intervention all provide the best possibility of producing significant and reproducible results in humans.

Future animal studies also might follow up on clinical observations that might provide novel insight into the pathogenesis of retinopathy. For example, how does high myopia [86–88] inhibit the development of diabetic retinopathy? What are the mechanisms behind the development of diabetic-like retinopathy following radiation [89, 90]?

Acknowledgements

This work was supported by grants EY022938 and R24EY024864 from NIH and BX003604 from the Department of Veterans Affairs to T.S.K., R01 EY012021 and EY029349 to D.A.A., and R24EY024868, EY017017, and BCH IDDCRC (1U54HD090255) to L.E.H.S.

Disclosure Statement

The authors have nothing to disclose.

References

- Engerman RL, Bloodworth JM Jr. Experimental diabetic retinopathy in dogs. *Arch Ophthalmol*. 1965 Feb;73(2):205–10.
- Engerman RL. Animal models of diabetic retinopathy. *Trans Sect Ophthalmol Am Acad Ophthalmol Otolaryngol*. 1976 Jul-Aug;81(4 Pt 1):OP710–5.
- Zheng L, Kern T. In vivo animal models of diabetic retinopathy; in HP H, M P (eds): *Experimental Approaches to Diabetic Retinopathy*. Basel, Karger, 2010, pp 42–60.
- Hatchell DL, Braun RD, Luttly GA, McLeod DS, Toth CA. Progression of diabetic retinopathy in a cat. *Invest Ophthalmol Vis Sci*. 1995;36:S1067.
- Kim SY, Johnson MA, McLeod DS, Alexander T, Hansen BC, Luttly GA. Neutrophils are associated with capillary closure in spontaneously diabetic monkey retinas. *Diabetes*. 2005 May;54(5):1534–42.
- Johnson MA, Luttly GA, McLeod DS, Otsuji T, Flower RW, Sandagar G, et al. Ocular structure and function in an aged monkey with spontaneous diabetes mellitus. *Exp Eye Res*. 2005 Jan;80(1):37–42.
- Engerman RL, Wallow IH. Retinopathy in nonhereditary diabetes. In: Podolsky S, Viswanathan M, editors. *Secondary Diabetes*. New York: Raven Press; 1980. pp. 503–19.
- Kim SY, Johnson MA, McLeod DS, Alexander T, Otsuji T, Steidl SM, et al. Retinopathy in monkeys with spontaneous type 2 diabetes. *Invest Ophthalmol Vis Sci*. 2004 Dec;45(12):4543–53.
- Kern TS, Tang J, Pan L, Li Z, Shen Y, Zeng W, et al. Assessment of macular thickening in spontaneously diabetic rhesus monkeys [ARVO abstract]. *Invest Ophthalmol Vis Sci*. 2018;59:3581.
- Engerman R, Bloodworth JM Jr, Nelson S. Relationship of microvascular disease in diabetes to metabolic control. *Diabetes*. 1977 Aug;26(8):760–9.
- Engerman RL, Kern TS. Progression of incipient diabetic retinopathy during good glyce-
- mic control. *Diabetes*. 1987 Jul;36(7):808–12.
- Engerman RL, Kern TS. Retinopathy in galactosemic dogs continues to progress after cessation of galactosemia. *Arch Ophthalmol*. 1995 Mar;113(3):355–8.
- Kowluru RA, Kanwar M, Kennedy A. Metabolic memory phenomenon and accumulation of peroxynitrite in retinal capillaries. *Exp Diabetes Res*. 2007;2007:21976.
- Chan PS, Kanwar M, Kowluru RA. Resistance of retinal inflammatory mediators to suppress after reinstitution of good glycemic control: novel mechanism for metabolic memory. *J Diabetes Complications*. 2010 Jan-Feb;24(1):55–63.
- Nathan DM, Genuth S, Lachin J, Cleary P, Crofford O, Davis M, et al.; Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med*. 1993 Sep;329(14):977–86.

- 16 Engerman RL, Kern TS. Aldose reductase inhibition fails to prevent retinopathy in diabetic and galactosemic dogs. *Diabetes*. 1993 Jun; 42(6):820–5.
- 17 Sorbinil Retinopathy Trial Research Group. A randomized trial of sorbinil, an aldose reductase inhibitor, in diabetic retinopathy. *Arch Ophthalmol*. 1990 Sep;108(9):1234–44.
- 18 Chung SS, Chung SK. Aldose reductase in diabetic microvascular complications. *Curr Drug Targets*. 2005 Jun;6(4):475–86.
- 19 Tang J, Du Y, Petrash JM, Sheibani N, Kern TS. Deletion of aldose reductase from mice inhibits diabetes-induced retinal capillary degeneration and superoxide generation. *PLoS One*. 2013 Apr;8(4):e62081.
- 20 Kador PF, Akagi Y, Takahashi Y, Ikebe H, Wyman M, Kinoshita JH. Prevention of retinal vessel changes associated with diabetic retinopathy in galactose-fed dogs by aldose reductase inhibitors. *Arch Ophthalmol*. 1990 Sep;108(9):1301–9.
- 21 Robison WG Jr, Laver NM, Jacot JL, Glover JP. Sorbinil prevention of diabetic-like retinopathy in the galactose-fed rat model. *Invest Ophthalmol Vis Sci*. 1995 Nov;36(12):2368–80.
- 22 Neuenschwander H, Takahashi Y, Kador PF. Dose-dependent reduction of retinal vessel changes associated with diabetic retinopathy in galactose-fed dogs by the aldose reductase inhibitor M79175. *J Ocul Pharmacol Ther*. 1997 Dec;13(6):517–28.
- 23 Kern TS, Engerman RL. Galactose-induced retinal microangiopathy in rats. *Invest Ophthalmol Vis Sci*. 1995 Feb;36(2):490–6.
- 24 Kern TS, Engerman RL. Pharmacological inhibition of diabetic retinopathy: aminoguanidine and aspirin. *Diabetes*. 2001 Jul;50(7):1636–42.
- 25 Zheng L, Howell SJ, Hatala DA, Huang K, Kern TS. Salicylate-based anti-inflammatory drugs inhibit the early lesion of diabetic retinopathy. *Diabetes*. 2007 Feb;56(2):337–45.
- 26 The DAMAD Study Group. Effect of aspirin alone and aspirin plus dipyridamole in early diabetic retinopathy. A multicenter randomized controlled clinical trial. *Diabetes*. 1989 Apr;38(4):491–8.
- 27 Group ET; Early Treatment Diabetic Retinopathy Study Research Group. Effects of aspirin treatment on diabetic retinopathy. ET-DRS report number 8. *Ophthalmology*. 1991 May;98(5 Suppl):757–65.
- 28 Cheng Y, Du Y, Liu H, Tang J, Veenstra A, Kern TS. Photobiodulation Inhibits Long-Term Structural and Functional Lesions of Diabetic Retinopathy. *Diabetes*. 2017.
- 29 Du Y, Cramer M, Lee CA, Tang J, Muthusamy A, Antonetti DA, et al. Adrenergic and serotonin receptors affect retinal superoxide generation in diabetic mice: relationship to capillary degeneration and permeability. *FASEB J*. 2015 May;29(5):2194–204.
- 30 Boot-Handford R, Heath H. Identification of fructose as the retinopathic agent associated with the ingestion of sucrose-rich diets in the rat. *Metabolism*. 1980 Dec;29(12):1247–52.
- 31 Tilton RG, Chang K, Nyengaard JR, Van den Enden M, Ido Y, Williamson JR. Inhibition of sorbitol dehydrogenase. Effects on vascular and neural dysfunction in streptozocin-induced diabetic rats. *Diabetes*. 1995 Feb;44(2):234–42.
- 32 Engerman RL, Kern TS. Experimental galactosemia produces diabetic-like retinopathy. *Diabetes*. 1984 Jan;33(1):97–100.
- 33 Engerman RL, Kern TS. Diabetic retinopathy: is it a consequence of hyperglycaemia? *Diabet Med*. 1985 May;2(3):200–3.
- 34 Kador PF, Akagi Y, Terubayashi H, Wyman M, Kinoshita JH. Prevention of pericyte ghost formation in retinal capillaries of galactose-fed dogs by aldose reductase inhibitors. *Arch Ophthalmol*. 1988 Aug;106(8):1099–102.
- 35 Kern TS, Engerman RL. Galactose-induced retinal microangiopathy in rats. *Invest Ophthalmol Vis Sci*. 1995 Feb;36(2):490–6.
- 36 Kern TS, Engerman RL. A mouse model of diabetic retinopathy. *Arch Ophthalmol*. 1996 Aug;114(8):986–90.
- 37 Barber AJ, Nakamura M, Wolpert EB, Reiter CE, Seigel GM, Antonetti DA, et al. Insulin rescues retinal neurons from apoptosis by a phosphatidylinositol 3-kinase/Akt-mediated mechanism that reduces the activation of caspase-3. *J Biol Chem*. 2001 Aug;276(35):32814–21.
- 38 Imai H, Misra GP, Wu L, Janagam DR, Gardner TW, Lowe TL. Subconjunctivally Implanted Hydrogels for Sustained Insulin Release to Reduce Retinal Cell Apoptosis in Diabetic Rats. *Invest Ophthalmol Vis Sci*. 2015 Dec;56(13):7839–46.
- 39 Reiter CE, Gardner TW. Functions of insulin and insulin receptor signaling in retina: possible implications for diabetic retinopathy. *Prog Retin Eye Res*. 2003 Jul;22(4):545–62.
- 40 Reiter CE, Sandrasegarane L, Wolpert EB, Klinger M, Simpson IA, Barber AJ, et al. Characterization of insulin signaling in rat retina in vivo and ex vivo. *Am J Physiol Endocrinol Metab*. 2003 Oct;285(4):E763–74.
- 41 Grant MB, Caballero S Jr. The potential role of octreotide in the treatment of diabetic retinopathy. *Treat Endocrinol*. 2005;4(4):199–203.
- 42 Thounaojam MC, Powell FL, Patel S, Gutsaeva DR, Tawfik A, Smith SB, et al. Protective effects of agonists of growth hormone-releasing hormone (GHRH) in early experimental diabetic retinopathy. *Proc Natl Acad Sci USA*. 2017 Dec;114(50):13248–53.
- 43 Adamis AP, Miller JW, Bernal MT, D'Amico DJ, Folkman J, Yeo TK, et al. Increased vascular endothelial growth factor levels in the vitreous of eyes with proliferative diabetic retinopathy. *Am J Ophthalmol*. 1994 Oct;118(4):445–50.
- 44 Miller JW, Adamis AP, Shima DT, D'Amore PA, Moulton RS, O'Reilly MS, et al. Vascular endothelial growth factor/vascular permeability factor is temporally and spatially correlated with ocular angiogenesis in a primate model. *Am J Pathol*. 1994 Sep;145(3):574–84.
- 45 Aiello LP, Bursell SE, Clermont A, Duh E, Ishii H, Takagi C, et al. Vascular endothelial growth factor-induced retinal permeability is mediated by protein kinase C in vivo and suppressed by an orally effective beta-isoform-selective inhibitor. *Diabetes*. 1997 Sep;46(9):1473–80.
- 46 Pierce EA, Foley ED, Smith LE. Regulation of vascular endothelial growth factor by oxygen in a model of retinopathy of prematurity. *Arch Ophthalmol*. 1996 Oct;114(10):1219–28.
- 47 Aiello LP, Pierce EA, Foley ED, Takagi H, Chen H, Riddle L, et al. Suppression of retinal neovascularization in vivo by inhibition of vascular endothelial growth factor (VEGF) using soluble VEGF-receptor chimeric proteins. *Proc Natl Acad Sci USA*. 1995 Nov; 92(23):10457–61.
- 48 Pierce EA, Avery RL, Foley ED, Aiello LP, Smith LE. Vascular endothelial growth factor/vascular permeability factor expression in a mouse model of retinal neovascularization. *Proc Natl Acad Sci USA*. 1995 Jan;92(3):905–9.
- 49 Titchener PM, Antonetti DA. Using the past to inform the future: anti-VEGF therapy as a road map to develop novel therapies for diabetic retinopathy. *Diabetes*. 2013 Jun;62(6):1808–15.
- 50 Keech AC, Mitchell P, Summanen PA, O'Day J, Davis TM, Moffitt MS, et al.; FIELD study investigators. Effect of fenofibrate on the need for laser treatment for diabetic retinopathy (FIELD study): a randomised controlled trial. *Lancet*. 2007 Nov;370(9600):1687–97.
- 51 Chew EY, Ambrosius WT, Davis MD, Danis RP, Gangaputra S, Greven CM, et al.; ACCORD Eye Study Group. Effects of medical therapies on retinopathy progression in type 2 diabetes. *N Engl J Med*. 2010 Jul;363(3):233–44.
- 52 Garcia-Ramírez M, Hernández C, Palomer X, Vázquez-Carrera M, Simó R. Fenofibrate prevents the disruption of the outer blood retinal barrier through downregulation of NF-κB activity. *Acta Diabetol*. 2016 Feb;53(1):109–18.
- 53 Gong Y, Shao Z, Fu Z, Edin ML, Sun Y, Liegl RG, et al. Fenofibrate Inhibits Cytochrome P450 Epoxygenase 2C Activity to Suppress Pathological Ocular Angiogenesis. *EBioMedicine*. 2016 Nov;13:201–11.
- 54 Noonan JE, Jenkins AJ, Ma JX, Keech AC, Wang JJ, Lamoureux EL. An update on the molecular actions of fenofibrate and its clinical effects on diabetic retinopathy and other microvascular end points in patients with diabetes. *Diabetes*. 2013 Dec;62(12):3968–75.
- 55 Simó R, Roy S, Behar-Cohen F, Keech A, Mitchell P, Wong TY. Fenofibrate: a new treatment for diabetic retinopathy. Molecular mechanisms and future perspectives. *Curr Med Chem*. 2013;20(26):3258–66.
- 56 Trudeau K, Roy S, Guo W, Hernández C, Villarreal M, Simó R, et al. Fenofibric acid reduces fibronectin and collagen type IV overexpression in human retinal pigment epithelial cells grown in conditions mimicking the diabetic milieu: functional implications in retinal permeability. *Invest Ophthalmol Vis Sci*. 2011 Aug;52(9):6348–54.

- 57 de Gooyer TE, Stevenson KA, Humphries P, Simpson DA, Gardiner TA, Stitt AW. Retinopathy is reduced during experimental diabetes in a mouse model of outer retinal degeneration. *Invest Ophthalmol Vis Sci*. 2006 Dec; 47(12):5561–8.
- 58 de Gooyer TE, Stevenson KA, Humphries P, Simpson DA, Curtis TM, Gardiner TA, et al. Rod photoreceptor loss in Rho-/- mice reduces retinal hypoxia and hypoxia-regulated gene expression. *Invest Ophthalmol Vis Sci*. 2006 Dec; 47(12):5553–60.
- 59 Du Y, Veenstra A, Palczewski K, Kern TS. Photoreceptor cells are major contributors to diabetes-induced oxidative stress and local inflammation in the retina. *Proc Natl Acad Sci USA*. 2013 Oct; 110(41):16586–91.
- 60 Liu H, Tang J, Du Y, Saadane A, Tonade D, Samuels I, et al. Photoreceptor Cells Influence Retinal Vascular Degeneration in Mouse Models of Retinal Degeneration and Diabetes. *Invest Ophthalmol Vis Sci*. 2016 Aug; 57(10):4272–81.
- 61 Tonade D, Liu H, Palczewski K, Kern TS. Photoreceptor cells produce inflammatory products that contribute to retinal vascular permeability in a mouse model of diabetes. *Diabetologia*. 2017 Oct; 60(10):2111–20.
- 62 Arden GB. The absence of diabetic retinopathy in patients with retinitis pigmentosa: implications for pathophysiology and possible treatment. *Br J Ophthalmol*. 2001 Mar; 85(3): 366–70.
- 63 Hammes HP, Martin S, Federlin K, Geisen K, Brownlee M. Aminoguanidine treatment inhibits the development of experimental diabetic retinopathy. *Proc Natl Acad Sci USA*. 1991 Dec; 88(24):11555–8.
- 64 Ishii H, Jirousek MR, Koya D, Takagi C, Xia P, Clermont A, et al. Amelioration of vascular dysfunctions in diabetic rats by an oral PKC beta inhibitor. *Science*. 1996 May; 272(5262): 728–31.
- 65 Kowluru RA, Jirousek MR, Stramm L, Farid N, Engerman RL, Kern TS. Abnormalities of retinal metabolism in diabetes or experimental galactosemia: V. Relationship between protein kinase C and ATPases. *Diabetes*. 1998 Mar; 47(3):464–9.
- 66 Kowluru RA, Engerman RL, Kern TS. Abnormalities of retinal metabolism in diabetes or experimental galactosemia. VI. Comparison of retinal and cerebral cortex metabolism, and effects of antioxidant therapy. *Free Radic Biol Med*. 1999 Feb; 26(3–4):371–8.
- 67 Joussen AM, Poulaki V, Mitsiades N, Kirchhof B, Koizumi K, Döhmen S, et al. Nonsteroidal anti-inflammatory drugs prevent early diabetic retinopathy via TNF-alpha suppression. *FASEB J*. 2002 Mar; 16(3):438–40.
- 68 Hammes HP, Du X, Edelstein D, Taguchi T, Matsumura T, Ju Q, et al. Benfotiamine blocks three major pathways of hyperglycemic damage and prevents experimental diabetic retinopathy. *Nat Med*. 2003 Mar; 9(3):294–9.
- 69 Zheng L, Szabó C, Kern TS. Poly(ADP-ribose) polymerase is involved in the development of diabetic retinopathy via regulation of nuclear factor-kappaB. *Diabetes*. 2004 Nov; 53(11):2960–7.
- 70 Hoffmann J, Alt A, Lin J, Lochnit G, Schubert U, Schleicher E, et al. Tenilsetam prevents early diabetic retinopathy without correcting pericyte loss. *Thromb Haemost*. 2006 Apr; 95(4):689–95.
- 71 Kanwar M, Chan PS, Kern TS, Kowluru RA. Oxidative damage in the retinal mitochondria of diabetic mice: possible protection by superoxide dismutase. *Invest Ophthalmol Vis Sci*. 2007 Aug; 48(8):3805–11.
- 72 Zhang JZ, Xi X, Gao L, Kern TS. Captopril inhibits capillary degeneration in the early stages of diabetic retinopathy. *Curr Eye Res*. 2007 Oct; 32(10):883–9.
- 73 Gubitosi-Klug RA, Talahalli R, Du Y, Nadler JL, Kern TS. 5-Lipoxygenase, but not 12/15-lipoxygenase, contributes to degeneration of retinal capillaries in a mouse model of diabetic retinopathy. *Diabetes*. 2008 May; 57(5): 1387–93.
- 74 Du Y, Tang J, Li G, Berti-Mattera L, Lee CA, Bartkowski D, et al. Effects of p38 MAPK inhibition on early stages of diabetic retinopathy and sensory nerve function. *Invest Ophthalmol Vis Sci*. 2010 Apr; 51(4):2158–64.
- 75 Li G, Tang J, Du Y, Lee CA, Kern TS. Beneficial effects of a novel RAGE inhibitor on early diabetic retinopathy and tactile allodynia. *Mol Vis*. 2011; 17:3156–65.
- 76 Wang Q, Gorbey S, Pfister F, Höger S, Dorn-Beineke A, Krügel K, et al. Long-term treatment with suberythropoietic Epo is vaso- and neuroprotective in experimental diabetic retinopathy. *Cell Physiol Biochem*. 2011; 27(6): 769–82.
- 77 Opreanu M, Tikhonenko M, Bozack S, Lydic TA, Reid GE, McSorley KM, et al. The unconventional role of acid sphingomyelinase in regulation of retinal microangiopathy in diabetic human and animal models. *Diabetes*. 2011 Sep; 60(9):2370–8.
- 78 Le YZ. VEGF production and signaling in Müller glia are critical to modulating vascular function and neuronal integrity in diabetic retinopathy and hypoxic retinal vascular diseases. *Vision Res*. 2017 Oct; 139:108–14.
- 79 Wang J, Xu X, Elliott MH, Zhu M, Le YZ. Müller cell-derived VEGF is essential for diabetes-induced retinal inflammation and vascular leakage. *Diabetes*. 2010 Sep; 59(9):2297–305.
- 80 Liu H, Tang J, Du Y, Saadane A, Samuels I, Veenstra A, et al. Transducin1, phototransduction and the development of early diabetic retinopathy. *Invest Ophthalmol Vis Sci*. Accepted for publication 2019.
- 81 Liu H, Tang J, Du Y, Lee CA, Golczak M, Muthusamy A, et al. Retinylamine Benefits Early Diabetic Retinopathy in Mice. *J Biol Chem*. 2015 Aug; 290(35):21568–79.
- 82 Talahalli R, Zarini S, Tang J, Li G, Murphy R, Kern TS, et al. Leukocytes regulate retinal capillary degeneration in the diabetic mouse via generation of leukotrienes. *J Leukoc Biol*. 2013 Jan; 93(1):135–43.
- 83 Li G, Veenstra AA, Talahalli RR, Wang X, Gubitosi-Klug RA, Sheibani N, et al. Marrow-derived cells regulate the development of early diabetic retinopathy and tactile allodynia in mice. *Diabetes*. 2012 Dec; 61(12):3294–303.
- 84 Bhatwadekar AD, Duan Y, Korah M, Thinschmidt JS, Hu P, Leley SP, et al. Hematopoietic stem/progenitor involvement in retinal microvascular repair during diabetes: implications for bone marrow rejuvenation. *Vision Res*. 2017 Oct; 139:211–20.
- 85 Kandpal RP, Rajasimha HK, Brooks MJ, Nellissery J, Wan J, Qian J, et al. Transcriptome analysis using next generation sequencing reveals molecular signatures of diabetic retinopathy and efficacy of candidate drugs. *Mol Vis*. 2012; 18:1123–46.
- 86 Lim LS, Lamoureux E, Saw SM, Tay WT, Mitchell P, Wong TY. Are myopic eyes less likely to have diabetic retinopathy? *Ophthalmology*. 2010 Mar; 117(3):524–30.
- 87 Chao DL, Lin SC, Chen R, Lin SC. Myopia is Inversely Associated With the Prevalence of Diabetic Retinopathy in the South Korean Population. *Am J Ophthalmol*. 2016 Dec; 172: 39–44.
- 88 Bazzazi N, Akbarzadeh S, Yavarikia M, Poorolajal J, Fouladi DF. High myopia and diabetic retinopathy: A contralateral eye study in diabetic patients with high myopic anisometropia. *Retina*. 2017 Jul; 37(7):1270–6.
- 89 Kinyoun JL. Long-term visual acuity results of treated and untreated radiation retinopathy (an AOS thesis). *Trans Am Ophthalmol Soc*. 2008; 106:325–35.
- 90 Ramos MS, Echegaray JJ, Kuhn-Asif S, Wilkinson A, Yuan A, Singh AD, et al. Animal models of radiation retinopathy - From teletherapy to brachytherapy. *Exp Eye Res*. 2019 Apr; 181:240–51.