Endoscopic-Assisted Powered Adenoidectomy versus Conventional Adenoidectomy: A Randomized Study

Saroo Singh B. Vageesh Padiyar Nishi Sharma

Department of Otorhinolaryngology, Dr. Ram Manohar Lohia Hospital, New Delhi, India

Keywords

Adenoidectomy · Conventional adenoidectomy · Endoscopic-assisted powered adenoidectomy

Abstract

Background: Adenoidectomy is one of the most common surgical procedures performed in children. Conventional curettage adenoidectomy has the advantage of being cost-effective and continues to be a commonly used method, especially in developing countries. **Objective:** To compare conventional curettage adenoidectomy and endoscopic-assisted powered adenoidectomy using a microdebrider. **Methods:** A total of 60 patients were randomly divided into two groups. Group I underwent conventional curettage adenoidectomy while group II underwent endoscopic-assisted powered adenoidectomy. Duration of surgery, amount of intraoperative bleeding, adequateness of removal, and damage to adjacent structures were assessed and compared between the two groups. All patients were followed for 24 months. **Results:** In our study, we observed approximately three times more mean blood loss and total operating time in the endoscopic-assisted powered procedure compared to the conventional technique (150 and 56 mL, respectively; 63 and 27 min, respectively). Nine (30%) cases in group I had more than 50% residual adenoid tissue while 20–50% of residual adenoid tissue was documented among 7 patients

(23%). Postoperative pain was found to be significantly higher in group I compared to group II. In both groups, recovery time ranged from 24 to 48 h with a mean of 33.6 h for group I and 36 h for group II; 23 patients (77%) in group II presented with residual disease in the 3-month follow-up period.

Conclusion: The endoscopic-assisted powered adenoidectomy technique is very efficient in complete removal of adenoid tissue, thus preventing patients from undergoing a second surgery and its possible associated risks. The precision of the microdebrider prevents damage to adjacent vital structures. However, on the other hand, conventional adenoidectomy scored better in terms of shorter duration of surgery and less intraoperative bleeding.

© 2019 The Author(s)

Published by S. Karger AG, Basel

Introduction

Wilhelm Meyer, in 1885, first described conventional adenoidectomy and since then it has been one of the most frequently performed surgical procedures in the pediatric age group [1, 2]. Indications for adenoidectomy include adenoid hypertrophy causing nasopharyngeal obstruction with subsequent sleep disordered breathing, otitis media with effusion, recurrent otitis media, and recurrent rhinosinusitis [3–6].

Over time, adenoidectomy has been performed using many new techniques which include an electronic molecular resonance, suction diathermy, a microdebrider, endoscopy, and laser [7–11]. Each method has advantages and disadvantages in terms of outcome, complications, operative time, and cost.

However, despite improvements in the techniques, complications related to the procedure are often inevitable. Major complications include primary and secondary hemorrhages while minor complications include fever, soreness, neck stiffness, and postoperative pain.

Currently, with the increasing trend towards day care surgery, it is essential for the surgeon to use the most optimal technique with the least postoperative morbidity. The classical surgical technique with adenoid curette has now evolved into a safer and more controlled removal of adenoids with the introduction of the endoscope and powered instruments such as the microdebrider.

Over the years, since its introduction in the late 1990s, endoscopic-assisted powered adenoidectomy has been described widely in the literature. However, considering it as a choice over the conventional curettage technique has been widely debated. To this objective, we compared the two techniques with respect to duration of surgery, amount of intraoperative bleeding, recovery time, adequateness of removal, damage to adjacent structures, and postoperative pain.

Materials and Methods

This randomized, single-blinded comparative study was performed in the Department of Otorhinolaryngology at Dr. Ram Manohar Lohia Hospital from August 2015 to July 2017. The study comprised 60 patients of either sex aged 6–12 years who were planned for adenoidectomy. The subjects were randomly divided into two groups. Randomization was performed using a table of random numbers for 30 subjects each. Group I included numbers 1–30 while group II included numbers 31–60. Each group had randomly distributed 15 even and 15 odd numbers. The study subjects were then allocated numbers according to the randomization. Those allocated with even numbers (group I) underwent conventional curettage adenoidectomy while those in group II (odd numbers) underwent endoscopic-assisted powered adenoidectomy. Patients with significant deviated nasal septum and patients with cleft palate were excluded from this study.

On enrolment, all patients underwent baseline evaluation which included a diagnostic nasal endoscopy. The grade of adenoid hypertrophy was assessed using the scale described by Clemens et al. [12].

Both surgical techniques were performed by a single surgeon and the observations were documented by a single resident. In the conventional technique, adenoidectomy was done using the St. Clair Thomson adenoid curette while in the endoscopic technique,

Table 1. Age and gender distribution

a Age distribution of all subjects ($n = 60$)

Age, years	
Mean $\pm$ SD	8.7 $\pm$ 2.3
Range	6–12
Median	8.5

b Groupwise age distribution of subjects

Technique	n	Mean age, years	p value
Conventional	30	9.30 $\pm$ 2.200	0.076
Endoscopic	30	8.23 $\pm$ 2.373	

c Gender distribution of subjects in the two groups

	Technique		Total
	conventional	endoscopic	
Gender, n (%)			
Female	8 (27%)	5 (17%)	13
Male	22 (73%)	25 (83%)	47
Total, n	30	30	60

the endoscope was used along with a microdebrider in the oscillating mode with saline irrigation. Curettage of the adenoid tissue was performed using 45-rad adenoidectomy blades with the oscillating speed up to 2,400 rpm. The procedure was visualized using a 2.7-mm or 4-mm rigid endoscope. An angled 45–70° scope was introduced through the oral cavity to visualize the working end of the microdebrider when it was not possible to introduce the scope from the opposite side.

The intraoperative parameters included total operative time, amount of bleeding, completeness of removal of adenoid tissue, and collateral damage to adjacent structures. Postoperative recovery time was recorded in all patients. The time period between the initial introduction and the final removal of the mouth gag was considered as the total operative time.

For the conventional adenoidectomy group, each 3-inch² soaked gauze was assumed to correspond to blood loss of 10 mL. However, in the endoscopic method, the blood loss was assessed by whatever came into the suction canister minus the irrigation solution.

The completeness of adenoid removal was assessed by nasal endoscopy at the end of the procedure in both groups; 20% or less residual adenoid was regarded as complete removal, 20–50% was regarded as partial, while more than 50% residual was considered suboptimal removal. Recovery time was defined as the total number of days taken to return to normal activity as gauged by the patient/guardian during the routine postoperative follow-up visit on the seventh day.

Statistical analysis was done using statistical software package SPSS v22.0. Data are represented as mean $\pm$ SD. Continuous variables were compared using the t test while nonparametric data

Table 2. Comparison of bleeding volume (mean $\pm$ SD) between the two groups

Technique	<i>n</i>	Bleeding volume, mL	<i>p</i> value
Primary bleeding			
Conventional	30	56.67 $\pm$ 18.952	<0.001
Endoscopic	30	149.33 $\pm$ 20.833	

Table 3. Comparison of total operative time (mean $\pm$ SD) between the two groups

Technique	<i>n</i>	Operative time, min	<i>p</i> value
Conventional	30	26.83 $\pm$ 5.796	<0.001
Endoscopic	30	62.67 $\pm$ 6.915	

Table 4. Comparison of recovery time (mean $\pm$ SD) between the two groups

Technique	<i>n</i>	Recovery time, days	<i>p</i> value
Conventional	30	33.60 $\pm$ 11.959	0.445
Endoscopic	30	36.00 $\pm$ 12.205	

were compared using the Mann-Whitney U test. The χ^2 test was used to compare the nominal data. A *p* value <0.05 was considered statistically significant.

Results

The mean age of all 60 patients aged 6–12 years was 8.7 ± 2.3 years. Group I comprised 8 males (26%) and 22 females (73.3%) while group II comprised 5 males (17%) and 25 females (83%). The mean age of patients was 9.3 years in group I and 8.2 years in group II. There was no statistical difference between the two groups with respect to age and gender distribution between the two groups (*p* = 0.076 and 0.347, respectively) (Table 1a–c).

The majority of the two groups showed grade III–IV adenoid hypertrophy (77% in group I and 87% in group II). Mann-Whitney U analysis was used to compare the mean rank of adenoid grades between the two groups.

There was no significant statistical difference between the two groups (*p* = 0.449).

In group I, blood loss ranged from 30 to 100 mL with mean blood loss of 56.6 mL. However, in group II, bleeding ranged from 100 to 180 mL with mean blood loss of 149.3 mL. This difference was statistically significant (*p* < 0.05) (Table 2).

Total operative times in groups I and II were 26.8 min (range 15–40 min) and 62.6 min (range 50–80 min), respectively. The difference in total operating time between the two groups was statistically significant (*p* < 0.05) (Table 3).

Postprocedural endoscopy done to look for residual adenoid tissue showed that resection was invariably complete by the endoscopic-assisted powered technique. Contrary to this, 9 (30%) cases in group I had more than 50% residual adenoid tissue while 20–50% of residual adenoid tissue was noticed among 7 cases (23%). Statistical comparison showed better completeness of removal with the endoscopic technique compared to the conventional procedure (*p* < 0.001).

Postoperative pain was found to be significantly higher in group I compared to group II, most likely due to injury to adjacent structures. In both groups, recovery time ranged from 24 to 48 h with a mean of 33.6 h for group I and 36 h for group II. There was no significant difference in recovery time following the two techniques (*p* > 0.05) (Table 4).

At the 3-month follow-up, no residual disease was found in group II. However, in group I, 23 patients (77%) presented with residual disease causing nasopharyngeal symptoms and sleep-disordered breathing. It was hence observed that chances of residual disease were significantly higher with the conventional technique compared to the endoscopic procedure (*p* < 0.001).

Discussion

Although conventional curettage adenoidectomy is a quick and simple procedure, it has its own associated complications owing to the blind technique. Besides injury to the eustachian tube orifice and pharyngeal musculature, various studies have reported a high percentage of residual tissue following this technique [13, 14]. Residual tissue usually leads to a sequence of potential problems which include peritubal obstruction, hyperplasia of remnant tissue, and a nidus for bacterial reservoirs. To overcome these drawbacks, the need for endoscopic-assisted microdebrider adenoidectomy came into existence.

In this regard, our study planned to compare the functional outcome of the two techniques considering certain specific parameters.

In all patients, we used the microdebrider transorally, since we felt it was suitable for children, especially those with narrow nasal passages. The safety and precision of the transoral curved microdebrider for adenoidectomy has been well documented in the literature [15–17].

In our study, we observed approximately three times more mean blood loss and total operating time in the endoscopic-assisted powered procedure compared to the conventional technique (150 and 56 mL, respectively; 63 and 27 min, respectively). Our observation was in contrast to studies performed by Koltai et al. [15] (2002), Murray et al. [16] (2002), Rodriguez et al. [17] (2002), and Heras and Koltai [18] (1998), who reported less total operative time and blood loss with the endoscopic-assisted powered technique.

All patients belonging to group II had complete removal of the adenoid tissue by the endoscopic method. However, in group I, 9 (30%) cases had more than 50% residual tissue while 20–50% of adenoid tissue was left in 7 patients (23%). The evidence of residual adenoid tissue postoperatively in patients undergoing conventional adenoid curettage has been reported by Havas and Lowinger [14], Stanislaw et al. [19], Datta et al. [20], Ezzat [21], and Hussein and Al-Juboori [22], with an incidence of 39, 39, 30, 14.5, and 20%, respectively.

Group II patients has more postoperative pain compared to those in group I. This was in agreement with the study performed by Lister et al. [23], who reported significant less postoperative pain in the debrider group.

The recovery period with the debrider-assisted adenoidectomy was shorter than that with conventional adenoidectomy and this difference was statistically significant. Similarly, a study by Somani et al. [24] showed less

morbidity after endoscopic-assisted powered adenoidectomy.

Conclusion

Although nasal endoscopes are fast becoming basic tools, powered instrumentation like microdebriders are not common. The need for special equipment and the cost of the procedure have to be borne in mind.

The newer method of endoscopic-assisted powered adenoidectomy was found to be a safe and useful tool for adenoidectomy. The advantages of this technique include completeness of resection, accurate removal, less damage to adjacent structures, less postoperative pain, and faster recovery.

However, in light of certain drawbacks such as increased total operative time, increased blood loss, need for special equipment, and cost of procedure limit its use. The use of powered adenoidectomy is technically demanding in the pediatric age group due to the relative difficulty in simultaneously passing both the scope and the debrider blade through the nose.

Statement of Ethics

The subjects (or their parents or guardians) gave their written informed consent. The study protocol was approved by the research institute's committee on human research.

Disclosure Statement

The authors have no conflicts of interest to declare. No funding was received for this study.

References

- 1 Thornval A. Wilhelm Meyer and the adenoids. *Arch Otolaryngol*. 1969 Sep;90(3): 383–6.
- 2 Al-Mazrou KA, Al-Qahtani A, Al-Fayez AI. Effectiveness of transnasal endoscopic powered adenoidectomy in patients with choanal adenoids. *Int J Pediatr Otorhinolaryngol*. 2009 Dec;73(12):1650–2.
- 3 Costantini F, Salamanca F, Amaina T, Zibordi F. Videoendoscopic adenoidectomy with microdebrider. *Acta Otorhinolaryngol Ital*. 2008 Feb;28(1):26–9.
- 4 Mitchell RB. Adenoidectomy techniques for sleep disordered breathing. *Oper Tech Otolaryngol*. 2005;16(4):229–31.
- 5 Wang DY, Clement PA, Kaufman L, Derde MP. Chronic nasal obstruction in children. A fiberoptic study. *Rhinology*. 1995 Mar;33(1): 4–6.
- 6 Ungkanont K, Damrongsak S. Effect of adenoidectomy in children with complex problems of rhinosinusitis and associated diseases. *Int J Pediatr Otorhinolaryngol*. 2004 Apr; 68(4):447–51.
- 7 Tarantino V, D'Agostino R, Melagrana A, Porcu A, Stura M, Vallarino R, et al. Safety of electronic molecular resonance adenoidectomy. *Int J Pediatr Otorhinolaryngol*. 2004 Dec; 68(12):1519–23.
- 8 Walker P. Pediatric adenoidectomy under vision using suction-diathermy ablation. *Laryngoscope*. 2001 Dec;111(12):2173–7.
- 9 Sorin A, Bent JP, April MM, Ward RF. Complications of microdebrider-assisted powered intracapsular tonsillectomy and adenoidectomy. *Laryngoscope*. 2004 Feb;114(2):297–300.

- 10 Shin JJ, Hartnick CJ. Pediatric endoscopic transnasal adenoid ablation. *Ann Otol Rhinol Laryngol*. 2003 Jun;112(6):511–4.
- 11 Özkiriş M, Karaçavuş S, Kapusuz Z, Saydam L. Comparison of two different adenoidectomy techniques with special emphasize on postoperative nasal mucociliary clearance rates: coblation technique vs. cold curettage. *Int J Pediatr Otorhinolaryngol*. 2013 Mar; 77(3):389–93.
- 12 Clemens J, McMurray JS, Willging JP. Electrocautery versus curette adenoidectomy: comparison of postoperative results. *Int J Pediatr Otorhinolaryngol*. 1998 Mar;43(2):115–22.
- 13 Cannon CR, Replogle WH, Schenk MP. Endoscopic-assisted adenoidectomy. *Otolaryngol Head Neck Surg*. 1999 Dec;121(6):740–4.
- 14 Havas T, Lowinger D. Obstructive adenoid tissue: an indication for powered-shaver adenoidectomy. *Arch Otolaryngol Head Neck Surg*. 2002 Jul;128(7):789–91.
- 15 Koltai PJ, Chan J, Younes A. Power-assisted adenoidectomy: total and partial resection. *Laryngoscope*. 2002 Aug;112(8 Pt 2 Suppl 100): 29–31.
- 16 Murray N, Fitzpatrick P, Guarisco JL. Powered partial adenoidectomy. *Arch Otolaryngol Head Neck Surg*. 2002 Jul;128(7):792–6.
- 17 Rodriguez K, Murray N, Guarisco JL. Power-assisted partial adenoidectomy. *Laryngoscope*. 2002 Aug;112(8 Pt 2 Suppl 100):26–8.
- 18 Heras HA, Koltai PJ. Safety of powered instrumentation for adenoidectomy. *Int J Pediatr Otorhinolaryngol*. 1998 Jul;44(2):149–53.
- 19 Stanislaw P Jr, Koltai PJ, Feustel PJ. Comparison of power-assisted adenoidectomy vs adenoid curette adenoidectomy. *Arch Otolaryngol Head Neck Surg*. 2000 Jul;126(7):845–9.
- 20 Datta R, Singh VP, Deshpal. Conventional versus Endoscopic Powered Adenoidectomy: a Comparative Study. *Med J Armed Forces India*. 2009 Oct;65:308–12.
- 21 Ezzat WF. Role of endoscopic nasal examination in reduction of nasopharyngeal adenoid recurrence rates. *Int J Pediatr Otorhinolaryngol*. 2010 Apr;74(4):404–6.
- 22 Hussein IA, Al-Juboori S. Conventional versus endoscopic-assisted adenoidectomy: a comparative study. *Med J Babylon*. 2012;9(3): 570–82.
- 23 Lister MT, Cunningham MJ, Benjamin B, Williams M, Tirrell A, Schaumberg DA, et al. Microdebrider tonsillotomy vs electrosurgical tonsillectomy: a randomized, double-blind, paired control study of postoperative pain. *Arch Otolaryngol Head Neck Surg*. 2006 Jun;132(6):599–604.
- 24 Somani SS, Naik CS, Bangad SV. Endoscopic adenoidectomy with microdebrider. *Indian J Otolaryngol Head Neck Surg*. 2010 Oct;62(4): 427–31.