Tamoxifen Exposure during Pregnancy: A Systematic Review and Three More Cases

Barbara Buonomo^{a,b} Antonella Brunello^c Stefania Noli^{b,d} Loredana Miglietta^e
Lucia Del Mastro^{f,g} Matteo Lambertini^{g,h} Fedro Alessandro Peccatori^b

^aGynecology and Obstetrics Unit, Department of Neuroscience, Reproductive Sciences and Dentistry, School of Medicine, University of Naples Federico II, Naples, Italy; ^bFertility and Procreation Unit, Division of Gynecologic Oncology, European Institute of Oncology IRCCS, Milan, Italy; ^cMedical Oncology 1 Unit, Department of Oncology, Istituto Oncologico Veneto IOV – IRCCS, Padua, Italy; ^dDepartment of Clinical Sciences and Community Health, Università degli Studi di Milano, Milan, Italy; ^eU.O.C. Oncologia Medica 2, Department of Medical Oncology, Ospedale Policlinico San Martino, Genoa, Italy; ^fS.S. Sviluppo Terapie Innovative, Department of Medical Oncology, Ospedale Policlinico San Martino, Genoa, Italy; ^gDepartment of Internal Medicine and Medical Specialties (DIMI), School of Medicine, University of Genoa, Genoa, Italy; ^hClinica di Oncologia Medica, Department of Medical Oncology, Ospedale Policlinico San Martino – IST, Genoa, Italy

Keywords

Breast carcinoma · Breast cancer · Pregnancy · Tamoxifen · Teratogenic effects

Abstract

Tamoxifen is frequently used as adjuvant treatment in premenopausal patients with hormone receptor-positive early breast cancer. According to guidelines, the use of nonhormonal barrier contraception is recommended during tamoxifen treatment and up to 3 months after its interruption prior to attempting conception. Nevertheless, when conception occurs inadvertently during tamoxifen treatment, the effects on the fetus and on the course of pregnancy are still not completely known. Here, we report 3 cases of young women who accidentally became pregnant while taking tamoxifen and perform a systematic review of the literature to provide more elements for better and clear multidisciplinary counselling of women facing this challenging situation.

© 2019 S. Karger AG, Basel

Introduction

Breast cancer represents the most common tumor diagnosed in women and the most frequent malignancy in patients of reproductive age [1–3]. In premenopausal patients with hormone receptor-positive early breast cancer, adjuvant endocrine therapy is indicated and may include the administration of tamoxifen for 5–10 years [4–9]. To avoid the risk of fetal malformation, pregnancy is contraindicated during tamoxifen treatment and up to 3 months after its interruption. Thus, barrier contraception is recommended, even if its uptake may be considered suboptimal [10–15]. Tamoxifen may also be secreted in breast milk. Hence, women taking tamoxifen should not breastfeed [16, 17].

Here, we report 3 cases of young women who accidentally became pregnant while taking tamoxifen and perform a systematic literature review to provide more elements for better counselling of women facing this challenging situation.

Case 1

A 40-year-old lady underwent quadrantectomy and sentinel lymph node biopsy in May 2011 for a moderately differentiated ductal invasive carcinoma (pT2 pN0(sn),

estrogen receptor [ER] 90%, progesterone receptor [PgR] 90%, Ki-67 15%, HER2 negative). Staging showed no evidence of distant metastases. In September 2011, the patient started treatment with tamoxifen 20 mg daily. In June 2012, after 9 months of tamoxifen treatment, a pelvic ultrasound showed a viable 12-week fetus. The patient decided to stop tamoxifen and to continue the pregnancy, which ended with a term vaginal delivery on December 20, 2012. The female newborn weighed 2,785 g and was 49.5 cm in length, appropriate for her gestational age. She subsequently attained all developmental milestones as assessed by her parents. The patient decided not to breast-feed and resumed tamoxifen treatment in February 2013. At present, she is doing well on tamoxifen, and has no evidence of disease 6 years after diagnosis.

Case 2

A 34-year-old lady presented with a left breast lump and underwent radical mastectomy and sentinel lymph node biopsy in May 2011. Histology revealed a moderately differentiated infiltrating ductal carcinoma (pT2 pN0(sn), ER 99%, PgR 90%, Ki-67 36%, HER2 negative). The patient received adjuvant chemotherapy with cyclophosphamide, epirubicin, and 5-fluorouracil for 3 cycles, followed by docetaxel for another 3 cycles. At completion of chemotherapy, tamoxifen 20 mg daily was initiated. A gonadotropin-releasing hormone agonist was given during chemotherapy and continued until June 2014 as part of adjuvant endocrine therapy. In November 2014, still under tamoxifen treatment, the patient became pregnant. According to ultrasound examination and the date of the last menstrual period, gestational age was 8 weeks. The patient decided to stop tamoxifen treatment and to continue her pregnancy. In July 2015, a healthy male infant was delivered at term by cesarean section with Apgar scores of 9/10 at 1 min and 10/10 at 5 min. The baby weighed 3,160 g and no malformation was described. Tamoxifen was resumed after delivery and was stopped in June 2017. At present, both the baby and the mother are well.

Case 3

In November 2007, a 38-year-old lady became pregnant while receiving adjuvant tamoxifen for a previously diagnosed low-grade ductal carcinoma in situ treated with conservative surgery. Tamoxifen was stopped at 7 weeks of gestation and the pregnancy was uneventful until 27 weeks, when a breast lump was discovered. She underwent left breast biopsy, revealing a new low-grade ductal carcinoma in situ. She delivered a healthy male infant by cesarean section at 34 weeks of gestation with normal weight and length according to gestational age (birthweight 2,800 g, length 46 cm). Thereafter, a left radical mastectomy was performed, demonstrating invasive ductal carcinoma (ER 55%, PgR 20%, Ki-67 14%, HER2 negative); adjuvant en-

docrine therapy with an aromatase inhibitor plus gonadotropin-releasing hormone agonist was prescribed. The boy is now 10 years old and he has attained the developmental milestones for age, except for a problem of dysorthography; the mother is well and free of disease.

Literature Review

To assess the obstetrical and fetal impact of prenatal exposure to tamoxifen, we performed a systematic review of the literature. We searched on PubMed for relevant papers published up to May 5, 2019, with restriction to publications in English, using the following keywords and Boolean operators: breast AND (carcinoma OR cancer OR neoplasm), AND (pregnancy OR pregnant OR gestation), AND (tamoxifen) AND (teratogenic effects). The work was done and reported according to the PRISMA guidelines for reporting of systematic reviews [18]. Guidelines of the American Society of Clinical Oncology (ASCO), European Society for Medical Oncology (ESMO), National Comprehensive Cancer Network (NCCN), and International Federation of Gynecology and Obstetrics (FIGO) were also consulted as sources of relevant references. Both series and cases of exposure during pregnancy were considered, dissecting possible confounders such as other concomitant treatment for which a teratogenic effect is described.

Results

The 3 patients reported on here had spontaneous pregnancies while taking tamoxifen during the 1st trimester, they all stopped the medication within the 12th week, and all had uneventful pregnancies, with normal newborns.

To contextualize our reports, we searched the literature for data on teratogenicity of tamoxifen in animals and conducted a systematic review on the effects of tamoxifen during pregnancy in humans, focusing on fetal wellbeing and the malformation rate.

The systematic research revealed 723 papers; of these, 718 papers were found on PubMed and 5 from other relevant sources. A total of 32 full-text articles and 1 abstract were deemed eligible and were included in this review; 12 papers reported data on teratogenic effects of tamoxifen in animals [8, 11, 19–28], and 21 were case reports, case series, and reviews about tamoxifen exposure during human pregnancies [8, 29–48] (Fig. 1).

The animal studies showed an increase in the incidence of abnormalities in the reproductive tracts of the offspring [11, 19–21] and irregularly ossified ribs in rat pups (“kinky ribs” and “wavy rib syndrome”) [11, 22]. Tamoxifen produced epithelial changes similar to those caused by diethylstilbestrol (DES) or clomiphene citrate [8], known teratogenic agents. Tamoxifen exposure resulted in metaplastic changes of the uterine epithelium of fetal rats and vaginal adenosis in newborn mice. Cunha et al. [23] examined the effect of tamoxifen in 54 genital

Fig. 1. PRISMA flowchart summarizing the process for the identification of eligible articles.

tracts isolated from 4- to 19-week-old human female fetuses and grown for 1–2 months in untreated athymic nude mice or host mice. They showed that tamoxifen in human fetal genital tracts resulted in delayed proliferation and maturation of the vaginal epithelium. Prenatal treatment of guinea pigs with tamoxifen showed delayed vaginal opening in female offspring, which was similar to DES-exposed female guinea pigs [24]. The long-term effects of tamoxifen use and whether it may increase gynecological cancers in daughters (as DES) are still unknown; in pregnant rats, tamoxifen exposure during gestation has been associated with breast cancer in the female offspring [25, 26]. Yamasaki et al. [27] performed an in utero and lactational exposure assay using tamoxifen at doses of 0.12, 0.6, or 3 $\mu\text{g}/\text{kg}/\text{day}$ in rats. A delay in preputial separation and hypospadias were detected at the doses of 0.6 and 3 $\mu\text{g}/\text{kg}/\text{day}$. No negative consequences for male and female fertility were found. Nobakht et al. [28] showed that tamoxifen treatment of pregnant rats could affect hippocampus formation in prenatal and postnatal rats.

The effects of tamoxifen during human pregnancy have been described in 17 case reports [29–45], in a trial on breast cancer prevention [46], in the Lareb and INCIP databases [47], in the records provided by AstraZeneca [8], and in a prospective analysis of pregnancies having occurred during and after treatment with trastuzumab and/or lapatinib in patients with early breast cancer [48]. Our systematic review of the literature identified a total of 249 fetuses exposed to tamoxifen during the 1st trimester of pregnancy or thereafter, including our 3 cases (Table 1). Among the 249 cases cited, there were 68 live births. Among the live births, 1 case showed genital tract abnormalities (ambiguous genitalia with clitoromegaly and labial fusion) in a female fetus exposed to tamoxifen for 20 weeks [29]. In another case, Goldenhar syndrome was reported, including right-sided microtia (underdevelopment of the exterior ear) and hemifacial microsomia (underdevelopment of the lower half of one side of the face). This infant, who also had preauricular skin tags, had been exposed to tamoxifen during the 1st and 2nd

Table 1. Summary of the reported cases with exposure to tamoxifen during pregnancy

Study [Ref.], year	Cases, <i>n</i>	Tamoxifen indication	Tamoxifen exposure	Complications of pregnancy	GA at delivery, weeks	Way of delivery	Fetal outcome	Weight at delivery, g	Long-term neonatal outcome	Confounders (other exposures during pregnancy)
Koizumi and Aono [34], 1986	2	Hyperprolactinemia and infertility due to pituitary adenoma	1st trimester	NR	Case 1: 41; Case 2: 37	Vaginal	Healthy at birth; no malformations	Case 1: 3,240; Case 2: 2,600	NR	Bromocriptine
Clark [46], 1993	85	Prevention of breast cancer	Unknown				No fetal anomalies			No information about duration of tamoxifen exposure and pregnancy outcome
Cullins et al. [30], 1994	1	Adjuvant hormone therapy for node-negative breast cancer	1st and 2nd trimesters	Preterm labor and chorioamnionitis	26	CS	Normal karyotype; Goldenhar syndrome	896	NR	X-ray (bone scan using ^{99m} Tc) and drugs (cocaine and marijuana)
DiPaola et al. [44], 1997	1	Multiple chemotherapy regimens for metastatic melanoma	2nd and 3rd trimesters	CS due to progression of the tumor in the mother	30	CS	Healthy at birth; no malformations	1,520	Normal milestones up to 15 months	Carmustine, dacarbazine, cisplatin
Tewari et al. [29], 1997	1	Hormone therapy for metastatic breast cancer	1st and 2nd trimesters	Induction of labor due to deteriorating condition of the patient	29	Vaginal	Normal karyotype; ambiguous genitalia	1,360	NR	NR
Isaacs et al. [32], 2001	1	Primary endocrine therapy for metastatic breast cancer	During whole pregnancy	CS due to deteriorating condition of the patient	31	CS	Normal karyotype; healthy at birth	1,940	Normal up to 2 years of age	Radiation and radiotherapy
Öksüzoglu and Güler [35], 2002	1	Adjuvant hormone therapy for node-negative breast cancer	1st trimester	NR	NR	NR	Healthy at birth; no malformations	NR	Normal milestones at 27 months	None
Andreadis et al. [41], 2004	1	Hormone therapy for metastatic breast cancer	During whole pregnancy	Preterm labor	35	CS	Healthy at birth; no malformations	2,070	Normal milestones at 12 months	Biphosphonate, chemotherapy (FEC) and radiotherapy exposure
Li et al. [33], 2007	1	Multiple chemotherapy regimens for metastatic melanoma	1st and 2nd trimesters	NO	34	CS	Healthy at birth; no malformations	2,750	NR	Multiple chemotherapy regimens (carmustine, dacarbazine, cisplatin) during 1st and 2nd trimesters
Berger and Clericuzio [31], 2008	1	Adjuvant hormone therapy for breast cancer	1st trimester	3rd trimester gestational diabetes; preeclampsia	32	Vaginal	Normal karyotype; microretrognathia, cleft of the secondary palate, glossop-tosis; Pierre Robin sequence + left acetabular and sacral dysplasia	1,983	Unknown	3rd trimester gestational diabetes; preeclampsia; use of ramelteon
Beale et al. [42], 2009	1	Adjuvant hormone therapy for breast cancer	1st and 2nd trimesters	Oligohydramnios; preterm premature rupture of membranes at 31 ⁺² weeks	31 ⁺⁶	CS	Twin A: no malformations; renal failure at 12 weeks of age Twin B: no malformations	Twin A: 1,590 Twin B: 1,705	Twin A: death at 13 weeks of age after respiratory arrest Twin B: unknown	Methadone; trastuzumab during 1st and 2nd trimesters; cigarette smoking; nifedipine tocolysis; preterm delivery
Warraich and Smith [43], 2009	1	Adjuvant hormone therapy for breast cancer	Until 28 weeks	Anhydramnios	37	CS	No malformations; severe pulmonary hypoplasia and atelectasis	2,690	Death a few days after birth	Herceptin and goserelin assumption; exposure to radioactive scans
Grandvillain et al. [45], 2009	1	NR	Until 16 weeks	NR	Medical abortion during 2nd trimester for maternal reason	NR	Female fetus with an enlarged clitoris	NR	NR	NR
Koca et al. [36], 2010	1	Adjuvant hormone therapy for breast cancer	1st trimester	Polyhydramnios	39	CS	Healthy at birth; no malformations	3,150	Healthy in her 66th month	Radiotherapy and chemotherapy (FEC) exposure
Koca et al. [37], 2013	3	Case 1: adjuvant endocrine therapy for metastatic breast cancer Case 2: adjuvant hormone therapy for breast cancer + LHRH Case 3: adjuvant hormone therapy	1st trimester	NR	NR	NR	Healthy at birth in cases 1 and 2 Case 3: voluntary medical abortion at 6 weeks	NR	Case 1: normal up to 2 years of age Case 2: NR	Case 1: chemotherapy (paclitaxel, carboplatin, doxorubicin)
Ishizuka and Satou [38], 2016	1	Adjuvant hormone therapy for breast cancer	1st and 2nd trimesters	NR	At term	CS	Healthy at birth; no malformations	2,544	Normal up to 5 years of age	Goserelin while pregnancy unknown

Table 1 (continued)

Study [Ref.], year	Cases, n	Tamoxifen indication	Tamoxifen exposure	Complications of pregnancy	GA at delivery, weeks	Way of delivery	Fetal outcome	Weight at delivery, g	Long-term neonatal outcome	Confounders (other exposures during pregnancy)
Jyoti et al. [39], 2016	1	Adjuvant hormone therapy for breast cancer	1st trimester	NO	39	CS	Healthy at birth; no malformations	Normal but not specified	NR	NR
Mohamed and Mirghani [40], 2017	1	Adjuvant hormone therapy for node-negative breast cancer	1st and 2nd trimesters	NO	At term	NR	Healthy at birth; no malformations	NR	NR	NR
AstraZeneca safety database	37	NR	1st trimester	NR	NR	NR	2 live births with congenital anomalies: 1 girl delivered at 29 weeks with XXX chromosomes and also a phallus-like clitoris and huge labia, and 1 with idiopathic chylothorax; 2 elective terminations with fetal defects; 6 spontaneous abortions; 6 live births without congenital anomalies; 4 elective terminations (no fetal defects or unknown); 17 unknown	NR	NR	NR
AstraZeneca safety database	15		After 1st trimester				1 live birth with congenital anomaly: congenital hand malformation; 1 elective termination with fetal defects; 9 live births without congenital anomaly; 1 elective termination (no fetal defects or unknown); 3 unknown			
AstraZeneca safety database	10		During whole pregnancy				1 live birth with congenital anomaly: Goldenhar syndrome (Cullins' report); 8 live births without congenital anomalies; 1 elective termination (no fetal defects or unknown)			
AstraZeneca safety database	74		Unknown				6 live births with congenital anomaly: 1 with cleft palate, 1 with ear malformation, 1 with trisomy 21, 1 with a small degree of labial fusion, 1 with craniofacial defects, and 1 with slight clitoral hypertrophy; 1 stillbirth with fetal defects; 3 elective terminations with fetal defects; 1 stillbirth without fetal defects; 5 spontaneous abortions; 1 ectopic pregnancy; 11 live births without congenital anomaly; 10 elective terminations (no fetal defects or unknown); 36 unknown			
Lambertini et al. [48], 2019	1	Hormone therapy for HER2-positive early breast cancer	1st trimester	NR	40	CS	Healthy at birth; no malformations	3,145	NR	Lapatinib + trastuzumab exposure
Lareb database	2	NR	1st trimester	NR	Case 1: at term Case 2: induced abortion	NR	Case 1: no congenital anomalies Case 2: induced abortion	NR	NR	NR
INCIP database	2	NR	1st trimester	NR	Case 1: spontaneous abortion at 10 weeks Case 2: at term	NR	Case 1: spontaneous abortion Case 2: no congenital anomalies	NR	NR	NR
Current clinical case 1	1	Adjuvant hormone therapy for breast cancer	1st trimester	NO	40	Vaginal	Healthy at birth; no malformations	NR	Normal up to 6 years of age	NR
Current clinical case 2	1	Adjuvant hormone therapy for breast cancer	1st trimester	NO	At term	CS	Healthy at birth; no malformations	3,160	Regular	NR
Current clinical case 3	1	Adjuvant hormone therapy for breast cancer	1st trimester	NO	34	CS	Healthy at birth; no malformations	2,800	Regular until the last follow-up visit	NR
GA, gestational age; NR, not reported; CS, cesarean section; LHRH, luteinizing hormone-releasing hormone.										

trimesters. In addition, the baby had been exposed to diagnostic X-rays and marijuana or cocaine at least once during the first 6 weeks of gestation [30]. Finally, another case report described a Pierre Robin sequence after 6

weeks of tamoxifen exposure. Noncraniofacial anomalies (acetabular and sacral dysplasia) were also reported [31]. The AstraZeneca Safety Database divided the cases according to the duration of the exposure [8]. In the group

Fig. 2. Duration of tamoxifen exposure during pregnancy. Pregnancies without malformations are depicted in green; pregnancies complicated by major malformations are depicted in red; pregnancies with minor malformations are depicted in orange. The letter by Clark [46] and the AstraZeneca Safety Database analysis [8] are not presented in this graph, as the exact duration of tamoxifen exposure was not specified.

of fetuses exposed during the 1st trimester only, 2 live births with congenital anomalies were included: 1 with a triple X karyotype and anomalies of the external genitalia (clitoris and labia) and 1 with idiopathic chylothorax. In the group of fetuses exposed after the 1st trimester, there was just 1 case of congenital malformation of one hand. This database also included 74 patients with unknown duration of tamoxifen exposure; of these, 6 women gave birth to 6 babies with congenital anomalies (Table 1).

Hence, among the 68 live births reported, there were 12 babies with congenital malformations (18%) [8, 29–31] and 56 without major anomalies (81%) [8, 32–44, 47, 48]; of these, 10 fetuses had been exposed to tamoxifen only from the 2nd trimester [8, 44], while 33 had been exposed from the 1st trimester. In 19 pregnancies, tamoxifen was stopped before the end of the 1st trimester [8, 34–37, 39, 47, 48], while in the other 16, administration continued till delivery [8, 32, 33, 38, 40, 42, 43]. Two of these 56 live births without major anomalies showed minor malformations; preauricular skin tags were reported in an otherwise normal infant following exposure to tamoxifen during the entire pregnancy [32]. Microph-

thalmos and severe hypermetropia were diagnosed in an infant at 1 year of age. This pregnancy had been exposed to tamoxifen both in the 1st and in the 2nd trimester, together with carmustine, dacarbazine, and cisplatin [33] (Fig. 2).

In summary, if we consider all the 248 pregnancies (249 fetuses) with in utero exposure to tamoxifen, 68 live births were reported, while there were 25 elective terminations, 12 spontaneous abortions, 1 ectopic pregnancy, 2 stillbirths, and 55 pregnancies whose outcome was unknown. As the letter of Clark [46] reporting on 85 healthy women who became pregnant while receiving chemoprophylactic tamoxifen did not describe the obstetric outcome of pregnancies or the exposure duration, we decided not to include these data in our systematic review.

Discussion and Conclusions

According to current guidelines [49, 50], use of tamoxifen is contraindicated during pregnancy and should be postponed after delivery. Nevertheless, as

shown in a recent survey investigating physicians' knowledge, practice, and attitudes regarding fertility and pregnancy issues in young breast cancer patients, 25 and 36% of the respondents disagreed or were neutral, respectively, on the statement that endocrine therapy should be avoided in pregnant patients [51]. In the present paper, we reported 3 more cases of patients accidentally exposed to tamoxifen during the 1st trimester of pregnancy and performed a systematic review of the literature to gather preclinical and clinical data about tamoxifen exposure during pregnancy and its effect on pregnancy and newborns.

Our 3 patients were exposed to tamoxifen during the 1st trimester only, without subsequent fetal malformations (Fig. 1). In our review, we documented 2 newborns with minor malformations [32, 33], whereas major malformations were observed in 12 infants of the 68 live births (17.6%) [8, 29–31]. As a point of reference, the prevalence of major malformations in the general population of the USA is 3% [52, 53].

The malformations in the reproductive tract were consistent with those reported in developmental toxicity studies in rats. Craniofacial malformations were observed in another 3 infants. Similar craniofacial malformations have been observed in cases of retinoic acid embryopathy, leading some researchers to hypothesize that tamoxifen may act on early organogenesis in a way similar to that of retinoic acid drugs [31, 53]. Thus, the apparent rate of malformations among all tamoxifen-exposed offspring, regardless of the severity of the malformations (both minor and major) or the gestational stage at exposure, was 20.5% (14/68). Both spontaneous abortions and stillbirths were documented in the literature, but no effective causal link with tamoxifen use can be established because of the paucity of cases. Notably, some patients received tamoxifen together with chemotherapy, illicit drugs, and X-rays, which may represent major confounding factors (Table 1).

Because of animal studies and case reports showing congenital abnormalities after tamoxifen exposure during pregnancy and because of the lack of long-term data on pediatric outcomes, no definitive conclusions on the teratogenic risk following tamoxifen use in pregnant women can be drawn. Although international guidelines contraindicate tamoxifen use during pregnancy, each case of accidental exposure must be approached individually. Our case series and systematic review of the literature supports the indication to stop tamoxifen during pregnancy and reinforces the need to provide adequate information on contraception to patients undergoing adjuvant endocrine therapy. The choice of contraceptive method among breast cancer survivors should consider the benefits of the method versus the risk of cancer recurrence. Thus, the ideal contraception meth-

od would be nonhormonal. In fact, the World Health Organization (WHO) contraindicates any hormonal contraception during breast cancer treatment. The first choice should be a copper intrauterine device [54]. A Cochrane study showed that in women with breast cancer treated with tamoxifen, the levonorgestrel-releasing intrauterine system led to a reduction in the incidence of endometrial polyps and hyperplasia [55]. Barrier methods (condoms and diaphragms) may also be chosen. Their limited efficacy is a major concern if pregnancy must be stringently controlled during breast cancer treatment. Female or male sterilization may also be discussed with couples who do not wish to have another pregnancy. Emergency contraception may be used if needed [56].

Patients receiving tamoxifen and interested in having a pregnancy should be informed about the need for a washout period of 3 months before conceiving. If an inadvertent pregnancy occurs, the potential risks for the fetus and newborn should be clearly discussed with patients, along with possible options, allowing women to make an informed decision. This also highlights the importance of framing an appropriate and patient-centered reproductive counselling in a dedicated setting.

Acknowledgements

For the conduction of this study, M. Lambertini acknowledges the support from the ESMO for a Translational Research Fellowship at the Institut Jules Bordet in Brussels (Belgium).

Statement of Ethics

The authors have no ethical conflicts to disclose. The subjects gave informed consent to published their clinical history anonymously.

Disclosure Statement

A. Brunello served as a consultant for Roche and Eli Lilly, and she is part of the advisory board for Eisai and for Eli Lilly; L. Del Mastro received honoraria from Takeda and personal fees from Ipsen and Takeda outside the submitted work; M. Lambertini served as a consultant for Teva, and he received honoraria from Theramex outside the submitted work; F.A. Peccatori received honoraria from Roche, AstraZeneca, Clovis, Takeda, and Ipsen outside the submitted work; the other authors do not have any financial disclosure to declare.

Funding Sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author Contributions

A. Brunello was the medical oncologist that provided care to the patient in case 1 and contributed to writing of the manuscript; L. Del Mastro and L. Miglietta were the medical oncologists that provided care to the patient in case 2 and contributed to write the manuscript; B. Buonomo and M. Lambertini developed the research strategy, searched the literature according to the PRISMA

methodology, and contributed to writing of the manuscript; B. Buonomo and S. Noli performed the data analysis, worked on the images and editorial adaptations, and contributed to writing of the manuscript; M. Lambertini and F.A. Peccatori coordinated the project, contributed to writing of the manuscript, and edited the last version of the paper; all authors contributed, revised parts of the manuscript, and agreed on the final version of the manuscript.

References

- Peccatori FA, Lambertini M, Scarfone G, Del Pup L, Codacci-Pisanelli G. Biology, staging, and treatment of breast cancer during pregnancy: reassessing the evidences. *Cancer Biol Med*. 2018 Feb;15(1):6–13.
- Rosenberg SM, Newman LA, Partridge AH. Breast cancer in young women: rare disease or public health problem? *JAMA Oncol*. 2015 Oct;1(7):877–8.
- Fidler MM, Gupta S, Soerjomataram I, Ferlay J, Steliarova-Foucher E, Bray F. Cancer incidence and mortality among young adults aged 20–39 years worldwide in 2012: a population-based study. *Lancet Oncol*. 2017 Dec;18(12):1579–89.
- Burstein HJ, Lacchetti C, Anderson H, Buchholz TA, Davidson NE, Gelmon KA, et al. Adjuvant Endocrine Therapy for Women with Hormone Receptor-Positive Breast Cancer: ASCO Clinical Practice Guideline Focused Update. *J Clin Oncol*. 2019 Feb 10;37(5):423–38.
- Curigliano G, Burstein HJ, Winer EP, Gnant M, Dubsy P, Loibl S, et al.; St. Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2017, André F, Baselga J, Bergh J, Bonnefoi H, Y Brucker S, Cardoso F, et al. De-escalating and escalating treatments for early-stage breast cancer: the St. Gallen International Expert Consensus Conference on the Primary Therapy of Early Breast Cancer 2017. *Ann Oncol*. 2017 Aug;28(8):1700–12.
- Paluch-Shimon S, Pagani O, Partridge AH, Abulkhair O, Cardoso MJ, Dent RA, et al. ESO-ESMO 3rd international consensus guidelines for breast cancer in young women (BCY3). *Breast*. 2017 Oct;35:203–17.
- Burstein HJ, Lacchetti C, Anderson H, Buchholz TA, Davidson NE, Gelmon KE, et al. Adjuvant Endocrine Therapy for Women with Hormone Receptor-Positive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline Update on Ovarian Suppression. *J Clin Oncol*. 2016 May;34(14):1689–701.
- Braems G, Denys H, De Wever O, Cocquyt V, Van den Broecke R. Use of tamoxifen before and during pregnancy. *Oncologist*. 2011;16(11):1547–51.
- Sonmezer M, Oktay K. Fertility preservation in young women undergoing breast cancer therapy. *Oncologist*. 2006 May;11(5):422–34.
- Castro-Sanchez A, Martinez-Cannon BA, Platas A, Mohar A, Fonseca A, Vega Y, et al. Sub-optimal Use of Effective Contraceptive Methods in Young Mexican Women with Breast Cancer. *J Glob Oncol*. 2018 Oct;4(4):1–7.
- Barthelmes L, Gateley CA. Tamoxifen and pregnancy. *Breast*. 2004 Dec;13(6):446–51.
- Nolvadex. In: *Physicians' Desk Reference Health*. 57th ed. Thomsen Healthcare; 2003.
- Mehta DK, editor. *Pregnancy. British National Formulary No. 45*. London: British Medical Association and Royal Pharmaceutical Society; 2004. p. 726. [Appendix 4].
- Briggs GG, Freeman RK, Yaffe SJ. *Drugs in pregnancy and lactation*. 6th ed. Philadelphia: Williams & Wilkins; 2002. pp. 1307–13.
- Pagani O, Ruggeri M, Manunta S, Saunders C, Peccatori F, Cardoso F, et al. Pregnancy after breast cancer: are young patients willing to participate in clinical studies? *Breast*. 2015 Jun;24(3):201–7.
- Zubor P, Kubatka P, Kapustova I, Miloseva L, Dankova Z, Gondova A, et al. Current approaches in the clinical management of pregnancy-associated breast cancer – pros and cons. *EPMA J*. 2018 Jun;9(3):257–70.
- Royal College of Obstetricians and Gynaecologists. *Pregnancy and Breast Cancer (Green-Top Guideline No. 12)* [Internet]. London: Royal College of Obstetricians and Gynaecologists; 2011. Available from: <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg12/>.
- Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009 Jul;339:b2535.
- Chamness GC, Bannayan GA, Landry LA Jr, Sheridan PJ, McGuire WL. Abnormal reproductive development in rats after neonatally administered antiestrogen (tamoxifen). *Biol Reprod*. 1979 Dec;21(5):1087–90.
- Iguchi T, Hirokawa M, Takasugi N. Occurrence of genital tract abnormalities and bladder hernia in female mice exposed neonatally to tamoxifen. *Toxicology*. 1986 Dec;42(1):1–11.
- Diwan BA, Anderson LM, Ward JM. Proliferative lesions of oviduct and uterus in CD-1 mice exposed prenatally to tamoxifen. *Carcinogenesis*. 1997 Oct;18(10):2009–14.
- Tucker MJ, Adam HK, Patterson JS. Tamoxifen. In: Laurence DR, McLean AE, Wetherall M, editors. *Safety testing of new drugs: laboratory predictions and clinical performance*. London: Academic Press; 1984. Chapter 6; pp. 125–61.
- Cunha GR, Taguchi O, Namikawa R, Nishizuka Y, Robboy SJ. Teratogenic effects clomiphene, tamoxifen, and diethylstilbestrol on the developing human female genital tract. *Hum Pathol*. 1987 Nov;18(11):1132–43.
- Hines M, Alsum P, Roy M, Gorski RA, Goy RW. Estrogenic contributions to sexual differentiation in the female guinea pig: influences of diethylstilbestrol and tamoxifen on neural, behavioral, and ovarian development. *Horm Behav*. 1987 Sep;21(3):402–17.
- Durrani S, Akbar S, Heena H. Breast Cancer during Pregnancy. *Cureus*. 2018 Jul;10(7):e2941.
- Halakivi-Clarke L, Cho E, Onojafé I, Liao DJ, Clarke R. Maternal exposure to tamoxifen during pregnancy increases carcinogen-induced mammary tumorigenesis among female rat offspring. *Clin Cancer Res*. 2000 Jan;6(1):305–8.
- Yamasaki K, Noda S, Muroi T, Mitoma H, Takakura S, Sakamoto S. Effects of in utero and lactational exposure to tamoxifen in SD rats. *Toxicol Lett*. 2005 Apr;156(2):289–96.
- Nobakht M, Najafzadeh N, Kordestani Shargh B. Effects of tamoxifen on morphological and ultrastructural aspects of developing hippocampus of rat. *Iran Biomed J*. 2009 Oct;13(4):237–43.
- Tewari K, Bonebrake RG, Asrat T, Shanberg AM. Ambiguous genitalia in infant exposed to tamoxifen in utero [letter]. *Lancet*. 1997 Jul;350(9072):183.
- Cullins SL, Pridjian G, Sutherland CM. Goldenhar's syndrome associated with tamoxifen given to the mother during gestation. *JAMA*. 1994 Jun;271(24):1905–6.
- Berger JC, Clericuzio CL. Pierre Robin sequence associated with first trimester fetal tamoxifen exposure. *Am J Med Genet A*. 2008 Aug;146A(16):2141–4.
- Isaacs RJ, Hunter W, Clark K. Tamoxifen as systemic treatment of advanced breast cancer during pregnancy – case report and literature review. *Gynecol Oncol*. 2001 Mar;80(3):405–8.
- Li RH, Tam WH, Ng PC, Mok TS, Tam B, Lau TK. Microphthalmos associated with Dartmouth combination chemotherapy in pregnancy: a case report. *J Reprod Med*. 2007 Jun;52(6):575–6.
- Koizumi K, Aono T. Pregnancy after combined treatment with bromocriptine and tamoxifen in two patients with pituitary prolactinomas. *Fertil Steril*. 1986 Aug;46(2):312–4.
- Öksüzoglu B, Güler N. An infertile patient with breast cancer who delivered a healthy child under adjuvant tamoxifen therapy. *Eur J Obstet Gynecol Reprod Biol*. 2002 Aug;104(1):79.
- Koca T, Akgun Z, Yucel SB, Dag NZ, Teomete M. Pregnancy a short time after multimodal therapy for bilateral breast cancer: a case report and review of literature. *J Oncol Pharm Pract*. 2011 Dec;17(4):440–3.

- 37 Koca E, Kuzan TY, Babacan T, Turkbeyler IH, Furkan S, Altundag K. Safety of tamoxifen during pregnancy: 3 case reports and review of the literature. *Breast Care (Basel)*. 2013 Dec;8(6):453–4.
- 38 Ishizuka S, Satou S. A case of delivery of healthy infant in breast cancer patient incidentally treated with goserelin acetate and tamoxifen during pregnancy. *Breast Cancer*. 2016 Jan;23(1):164–6.
- 39 Jyoti B, Bharat C, Ankita N, Munita B, Sudeep G. Pregnancy on tamoxifen: case-report and review of literature. *South Asian J Cancer*. 2016 Oct-Dec;5(4):209–10.
- 40 Mohamed KEH, Mirghani S. Breast cancer case using tamoxifen during pregnancy: a case report and literature review. *Sudan J Med Sci*. 2017;12(2).
- 41 Andreadis C, Charalampidou M, Diamantopoulos N, Chouchos N, Mouratidou D. Combined chemotherapy and radiotherapy during conception and first two trimesters of gestation in a woman with metastatic breast cancer. *Gynecol Oncol*. 2004 Oct;95(1):252–5.
- 42 Beale JM, Tuohy J, McDowell SJ. Herceptin (trastuzumab) therapy in a twin pregnancy with associated oligohydramnios. *Am J Obstet Gynecol*. 2009 Jul;201(1):e13–4.
- 43 Warraich Q, Smith N. Herceptin therapy in pregnancy: continuation of pregnancy in the presence of anhydramnios. *J Obstet Gynaecol*. 2009 Feb;29(2):147–8.
- 44 Dipaola RS, Goodin S, Ratzell M, Florczyk M, Karp G, Ravikumar TS. Chemotherapy for metastatic melanoma during pregnancy. *Gynecol Oncol*. 1997 Sep;66(3):526–30.
- 45 Grandvuillemin A, Rousseau T, Laurent N, Meyer F, Lebouvier M, Disson-Dautriche A. A case of sexual ambiguity under tamoxifen during pregnancy. *Fundam Clin Pharmacol*. 2009;23 Suppl 1:37.
- 46 Clark S. Prophylactic tamoxifen. *Lancet* 1993; 342:168.
- 47 Schuurman TN, Witteveen PO, van der Wall E, Passier JL, Huitema AD, Amant F, et al. Tamoxifen and pregnancy: an absolute contraindication? *Breast Cancer Res Treat*. 2019 May;175(1):17–25.
- 48 Lambertini M, Martel S, Campbell C, Guillaume S, Hilbers FS, Schuehly U, et al. Pregnancies during and after trastuzumab and/or lapatinib in patients with human epidermal growth factor receptor 2-positive early breast cancer: analysis from the NeoALTTO (BIG 1-06) and ALTTO (BIG 2-06) trials. *Cancer*. 2019 Jan 15;125(2):307–16.
- 49 Peccatori FA, Azim HA Jr, Orecchia R, Hoekstra HJ, Pavlidis N, Kesic V, et al.; ESMO Guidelines Working Group. Cancer, pregnancy and fertility: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2013 Oct;24 Suppl 6:vi160–70.
- 50 Loibl S, Schmidt A, Gentilini O, Kaufman B, Kuhl C, Denkert C, et al. Breast cancer diagnosed during pregnancy: adapting recent advances in breast cancer care for pregnant patients. *JAMA Oncol*. 2015 Nov;1(8):1145–53.
- 51 Lambertini M, Di Maio M, Pagani O, Curiigliano G, Poggio F, Del Mastro L, et al. The BCY3/BCC 2017 survey on physicians' knowledge, attitudes and practice towards fertility and pregnancy-related issues in young breast cancer patients. *Breast*. 2018 Dec;42:41–9.
- 52 Correa A, Cragan JD, Kucik JE, Alverson CJ, Gilboa SM, Balakrishnan R, et al. Reporting birth defects surveillance data 1968–2003. *Birth Defects Res A Clin Mol Teratol*. 2007 Feb;79(2):65–186.
- 53 National Toxicology Program. NTP Monograph: Developmental Effects and Pregnancy Outcomes Associated with Cancer Chemotherapy Use during Pregnancy. *NTP Monogr*. 2013 May;(2):i-214.
- 54 Gompel A, Ramirez I, Bitzer J; European Society of Contraception Expert Group on Hormonal Contraception. Contraception in cancer survivors – an expert review. Part I. Breast and gynaecological cancers. *Eur J Contracept Reprod Health Care*. 2019 Jun;24(3):167–74.
- 55 Dominick S, Hickey M, Chin J, Su HI. Levonorgestrel intrauterine system for endometrial protection in women with breast cancer on adjuvant tamoxifen. *Cochrane Database Syst Rev*. 2015 Dec;(12):CD007245.
- 56 Medica AC, Stark SS, Hadnott TN, Dietz AC, Romero SA, Natarajan L, et al. Use of emergency contraception among female young adult cancer survivors. *Fertil Steril*. 2018 Jun; 109(6):1114–1120.e1.