Rational Development of Peptide Vaccination in Clinical Oncology

Concerning Jäger E, et al.: Peptide vaccination in clinical oncology. *Onkologie* 2000;23:410–415.

Antigen-specific vaccination is a very promising new option for cancer patients. As reviewed by Jäger et al., results from early clinical vaccination trials are of interest, since induction of T-cell responses and even tumor regressions in a few patients were observed. Objective tumor remissions, however, were rarely observed in most studies performed so far (review in [1]) and were usually restricted to melanoma patients with limited disease. In the MAGE-3 peptide study cited by Jäger et al. with a 30% response rate, the 7 patients showing an objective tumor remission all had metastatic disease limited to soft tissue or lymph nodes and the response evaluation did not include the 14 patients with early progressive disease during vaccination [2]. Jäger et al. so far have published in 1996 only the results from 3 patients with objective tumor regression by peptide vaccination with GM-CSF [3], but since then not the results on further patients vaccinated with this protocol. In a recent two-center phase II trial, in which we had vaccinated 16 patients with metastatic melanoma with tyrosinase peptides and GM-CSF using similar dosages as Jäger et al., no objective tumor remissions were observed and T-cell responses were recorded in only 2 of 16 patients with a mixed response and a stable disease [4].

Future perspectives need to focus on more potent vaccination strategies. This can be either achieved using more potent molecular or cellular adjuvants or by vaccinating patients in an adjuvant situation or with minimal residual disease, as suggested by recent studies [5]. Systematic comparative development of vaccination strategies is urgently warranted, since so far its role in adjuvant and therapeutic situations remains undefined. This is best achieved in form of controlled clinical phase I–III trials as currently performed within the EORTC (European Organization for Research and Treatment), and the ADO (Arbeitsgemeinschaft Dermato-Onkologie). Also crucial for the development of antigen-specific vaccination are sensitive and standardized T-cell assays to monitor the induction of specific T-cell responses.

Within the EORTC melanoma group we have performed several expert workshops on T-cell assay methodology and have evaluated different assays in a multicenter laboratory study showing the high sensitivity and reproducibility of the ELISPOT assay (introduced into the field by Thomas Woelfel) in contrast to chromium release CTL assays [6]. As a basis for a randomized adjuvant phase III trial, we have subsequently performed a series of phase I trials to evaluate the immunogenicity of different adjuvants in peptide-based vaccination. In two consecutive phase I trials we could not detect a difference

in frequency, strength or duration of peptide-induced T-cell responses if peptides were applied alone or in combination with GM-CSF, using a similar schedule as reported by Jäger et al. [3], but we could show a strong adjuvant effect if immunogenic 'T helper proteins' were added as adjuvants [7].

Randomized phase III peptide vaccination trials will be activated shortly by the EORTC for patients with resected stage III and IV melanoma as well as for patients with high-risk ocular melanoma. In the USA, the ECOG (Eastern Cooperative Oncology Group) is following the same path and has recently activated a similar study evaluating the effect of peptide vaccination and the role of GM-CSF in a randomized phase III trial in stage IV patients after resection of metastases. In patients with metastatic melanoma the therapeutic efficacy of peptide vaccination with dendritic cells is currently being evaluated in a randomized trial against standard DTIC chemotherapy within the ADO.

Before results from these ongoing studies are available, patients should be treated whenever possible within controlled multicenter trials to obtain reliable data on the role of antigen-specific immunotherapy in adjuvant and therapeutic cancer treatment.

U. Keilholz, C. Scheibenbogen, E. Thiel, Berlin
D. Schadendorf, Mannheim

References

- Schadendorf D, Schuler G, Paschen A, Scheibenbogen C. Experimentelle Therapie des Melanom. *Onkologie* 2001;7:76–83.
- Marchand M, Weynants P, Rankin E, Arienti F, Belli F, Parmiani G, Cascinelli N, Bourlond A, Vanwijk R, Humblet Y: Tumor regression responses in melanoma patients treated with a peptide encoded by gene MAGE-3. *Int J Cancer* 1995;63:883–885.
- Jäger E, Ringhoffer M, Dienes HP, Arand M, Karbach J, Jäger D, Ilsemann C, Hagedorn M, Oesch F, Knuth A: GM-CSF enhances immune responses to melanoma-associated peptides in vivo. *Int J Cancer* 1996; 67:54–62.
- Scheibenbogen C, Schmitt A, Keilholz U, Allgauer T, Hofmann U, Max R, Thiel E, Schadendorf D: Phase II trial of vaccination with tyrosinase peptides and GM-CSF in melanoma. *J Immunother* 2000;23:275–281.
- Keilholz U, Scheibenbogen C, Hoffmann U, Servetopoulou F, Schmitt A, Siehl J, Foerster MH, Thiel E, Schadendorf D: Therapeutic vs. adjuvant peptide vaccination: Comparison of two phase I/II trials in melanoma patients. *Proc Am Soc Clin Oncol* 2000;2102.
- Scheibenbogen C, Romero P, Rivoltini L, Herr W, Schmitt A, Cerrotini JC, Woelfel T, Eggermont AMM, Keilholz U: Quantitation of antigen-reactive T cells in peripheral blood by IFNγ-ELISPOT assay and chromium release assay: A four-centre comparative study. *J Immunol Meth* 2000;244:81–89.
- Keilholz U, Scheibenbogen C, Foerster MH, Hofmann U, Nagorsen D, Servetopoulou F, Bechrakis N, Schmitt A, Philipp A, Siehl J, Thiel E, Schadendorf D: Phase I study in high-risk melanoma patients of vaccination with MHC class I binding tyrosinase peptides comparing various immunological adjuvants. 2001, submitted.

Concerning Bembenek A, Schlag PM: Sentinel lymphadenectomy – ready for clinical routine. *Onkologie* 2000;23:408–409.

The timely editorial by Bembenek and Schlag [1] falls somewhat short to the current controversy surrounding the issue of sentinel lymphadenectomy in breast cancer. A glance at a recent issue of a leading journal [2] shows that – with regard to breast cancer and the axilla – we are ‘not entirely out of the labyrinth’. From a surgical standpoint, the problem may be that sentinel lymphadenectomy has become ready for clinical routine. However, in view of individualization of (surgical) therapy and the current widespread use of hormonal and/or cytotoxic therapy even in patients with early breast cancer, the more important point appears to be that axillary lymphonodectomy might not be needed at all. Perhaps, the authors of the editorial wish to elaborate on that matter.

D. Fritze, Darmstadt

References

- 1 Bembenek A, Schlag PM: Sentinel lymphadenectomy – ready for clinical routine? *Onkologie* 2000;23:408–409.
- 2 Costa A: Editorial. Breast cancer and the axilla: Not entirely out of the labyrinth. *Ann Oncol* 2000;11:1079–1080.

Authors' Reply

We are pleased about the comments on our publication. Raised was the interesting question ‘Do we need axillary dissection at all?’ This statement is based on two assumptions:

1. Removal of axillary nodal metastases does not alter survival nor influence local control.
2. Systemic treatment can always be guided by the features of the primary tumor.

Nodal status is still the strongest predictor of survival. Not yet defined is the role of micrometastases. But recent studies gave rise to the assumption, that even immunohistochemically detected metastases play a prognostic role [1]. Studies in the adjuvant setting and the studies on the survival benefit of axillary lymph node dissection (ALND) were conducted without any information about micrometastases and, thus, are at risk to be confounded by these metastatic entities. The selective examination of the sentinel lymph nodes (SN) including serial sections and immunohistochemistry increases the percentage of patients with potential nodal disease in about 10–20% of the cases [2].

Recent retrospective meta-analyses showed a significant survival benefit [3] resp. a risk reduction [4] in early breast cancer patients after ALND with [3] or without [4] adjuvant therapy. This rather small but net difference may reflect the percentage of patients with an early stage of nodal metastasis confined to the axilla, not reflecting systemic disease, according to the spectrum model proposing the existence of all stages of disease, replacing the Fisher hypothesis [5].

All the available studies proving the efficacy of systemic adjuvant therapy had been conducted either after axillary dissection or without effective axillary staging, because non-invasive diagnostics have by far not reached the sensitivity and accuracy of the sentinel lymph node biopsy (SLNB). This means that all the studies were either conducted after surgical treatment of the axilla or did not consider the nodal status: They had to accept a compromise in one of the two directions. With SLNB we have got a completely new setting, that reveals the nodal status exactly, but leaves behind an otherwise intact axillary region. It is evident, that apart from nodal-negative patients, who are spared from ALND, this setting is attractive even for SN-positive patients with small tumors: SLNB removes in a high percentage of patients the only involved node or, at least, the center of nodal disease without significant morbidity [6] and leaves an otherwise untouched axilla in situ: It would be logic that the results of adjuvant therapy in this setting could be different to former therapies and its indications. Ongoing studies in the USA have recently been created to evaluate this setting for (sentinel) nodal positive patients.

Another potential field of application would be the neoadjuvant concept, where SLNB may select patients with small tumors for preoperative therapy.

Finally, the avoidance of lymphnode dissection would ignore the selectivity of SLNB with 15% of newly diagnosed micrometastases, the possibility to determine the strongest predictor of survival exactly with low morbidity and refuse the potential of surgical cure for some of the early breast cancer patients. We agree that individualization of the surgical therapy is one of the most important principles of future medicine. But – in our opinion – SLNB is an important step forward to individualized therapy because it overcomes ultimately the Halstedian age of standardized radical surgery and adjuvant therapy. What has to be renewed in the future are the current strategies for adjuvant therapy considering the new potentials of axillary staging.

A. Bembenek and P. M. Schlag, Berlin

References

- 1 Clare SE, Sener SF, Wilkens W, Goldschmidt R, Merkel D, Winchester DJ: Prognostic significance of occult lymph node metastases in node-negative breast cancer. *Ann Surg Oncol* 1997;4:447–451.
- 2 Schreiber RH, Pendas S, Ku NN, et al: Microstaging of breast cancer patients using cytokeratin staining of the sentinel lymph node. *Ann Surg Oncol* 1999;6:95–101.
- 3 Bland KI, Scott-Conner CE, Menck H, Winchester DP: Axillary dissection in breast-conserving surgery for stage I and II breast cancer: A national cancer data base study of patterns of omission and implications for survival. *J Am Coll Surg* 1999;188:586–596.
- 4 Orr RK: The impact of prophylactic axillary node dissection on breast cancer survival – a Bayesian metaanalysis. *Ann Surg Oncol* 1999;6:109–116.
- 5 Harris JR, Osteen RT: Patients with early breast cancer benefit from effective axillary treatment. *Breast Cancer Res Treat* 1985;5:17–25.
- 6 Schrenk P, Rieger R, Shamiyeh A, Wayand W: Morbidity following sentinel lymph node biopsy versus axillary lymph node dissection for patients with breast carcinoma. *Cancer* 2000;88:608–614.

Significance of Chromosomal Aberrations in CLL

Concerning Jarošová M, et al.: Contribution of comparative genomic hybridization and fluorescence in situ hybridization to the detection of chromosomal abnormalities in B-cell chronic lymphocytic leukemia. *Onkologie* 2001;24:60–65.

The introduction of purine analogs a few years ago has initiated a major interest in the therapeutic strategies in chronic lymphocytic (CLL). Remarkable remission rates including clinical complete remissions (CR) intensified the search of potentially curative strategies with autologous or allogeneic stem cell transplantation [1]. For the development of optimal strategies for a individual patient, risk-oriented management of the CLL may be an important task.

Besides some clinical parameters, the diagnosis of chromosomal aberrations in CLL is increasingly of interest, as risk groups may be defined by special aberrations (e.g. 17p-, 11q-, trisomy 12, 13q-). Whereas the median survival in patients with 17p- is < 3 years, patients with 13q- have a favorable life expectancy of about 10 years [2, 3]. Conventional cytogenetic analysis, however, is hampered by the low mitotic rate of CLL cells. Here, techniques as fluorescence in situ hybridization (FISH) and comparative genomic hybridization (CGH) have provided sensitive tools for defining specific chromosomal aberrations in CLL. Döhner et al. [2, personal communication] reported chromosomal aberrations using a broad panel of specific probes in up to 78% of cases, whereby 13q- deletion (44%), 11q- (25%), 17p- (10%) and trisomy 12 (13%) were the most common ones.

Jarošová et al. [4] investigated 88 patients with CLL for chromosomal aberrations with conventional banding technique analysis as well as with FISH and CGH. With conventional cytogenetic analysis, in only 19% of the patients an abnormal karyotype was identified. But performing interphase cytogenetic analysis with FISH or CGH, in 54% and 51% of CLL patients, respectively, an abnormal karyotype was found.

The data of Jarošová et al. [4] clearly indicate that using banding techniques in conventional cytogenetics is difficult in CLL due to low rate of mitosis. Interphase techniques as FISH and/or CGH are much more sensitive methods [3, 4]. While banding techniques identify only gross chromosomal aberrations, molecular cytogenetic approaches as FISH identify small aberration regions and translocation breakpoints. On the First and Second International Working Party on Chromosomes in CLL 1991, a comparison of banding techniques with FISH on a large series of patients showed that structural 13q aberrations were identified only in 10% of cases by the conventional banding techniques but in 53% of patients using FISH [5]. Additionally, the incidence of chromosomal aberrations in

CLL varies considerably in the literature, even when using FISH and/or CHG. These different incidences may be caused by the quantity and quality of specific probes used and by the target size. Jarošová et al. [4] identified a deletion of 13q14 in only 18% of their patients. This low incidence in comparison to others [3] may be due to used probes, as translocations are not only involve the retinoblastoma tumor suppressor gene (RBI) on chromosome 13 but more frequently a more distal locus on this chromosome [6].

In contrast to FISH, CGH allows screening for the presence of chromosomal imbalances in a tumor genome without knowing regions that might be altered. Especially for the detection of additional imbalances, CGH is an adequate method. Recently, however, Barth et al. [7] reported their data on CGH and showed that genomic imbalances might be identified with a high sensitivity but dependent on the used method also false-positive results may be obtained.

Taken together, the identification of chromosomal aberrations in CLL by FISH and/or CGH is a promising tool for the identification of risk groups. However, standardizations of the probes and quality controls between the laboratories are required, especially if treatment stratifications will be based on aberration-defined risk groups.

L. Bergmann, Frankfurt/M.

References

- 1 Bergmann L: Present status of purine analogs in the therapy of chronic lymphocytic leukemias. *Leukemia* 1997;11(suppl 2):29–34.
- 2 Döhner H, Stilgenbauer S, Fischer K, Bentz M, Lichter P: Cytogenetic and molecular cytogenetic analysis of B-cell chronic lymphocytic leukemia: Specific chromosome aberrations identify prognostic subgroups of patients and point to loci of candidate genes. *Leukemia* 1997; 11(suppl 2):4–13.
- 3 Döhner H, Stilgenbauer S, Döhner K, Bentz M, Lichter P: Chromosome aberration in B-cell chronic lymphocytic leukemia: Reassessment based on molecular cytogenetic analysis. *J Mol Med* 1999;77:266–281.
- 4 Jarošová M, Jedlicková K, Holzerová M, Urbanová R, Papajik T, Raida L, Pikalová Z, Lakomá I, Pepoková I, Kropáková J, Indrák K: Contribution of comparative genomic hybridization and fluorescence in situ hybridization to the detection of chromosomal abnormalities in B-cell chronic lymphocytic leukemia. *Onkologie* 2001;24:60–65.
- 5 Juliusson B, Oscier D, Gahrton G for the International Working Party on chromosomes in CLL (IWCCLL): Cytogenetic findings and survival in B-cell chronic lymphocytic leukemia. Second IWCCLL compilation of data on 662 patients. *Leuk Lymphoma* 1991;5:21–25.
- 6 Brown AG, Ross FM, Dunne EM, Steel CM, Weit-Thompson EM: Evidence for a new tumour suppressor locus in human B-cell neoplasia telomeric to the retinoblastoma gene. *Nat Genet* 1993;3:67–72.
- 7 Barth TF, Benner A, Bentz M, Döhner H, Moller P, Lichter P: Risk of false positive results in comparative genomic hybridization. *Genes Chromosomes Cancer* 2000;28:353–357.