Case Report

Rebiopsy with Thoracoscopy under Local Anesthesia for the Detection of EGFR T790M Mutation

Asako Matsuda^a Yasuko Fuchimoto^a Sae Wada^a Takaaki Tanaka^a
Tetsuya Takeguchi^a Sho Mitsumune^a Yuki Takigawa^a
Yosuke Miyamoto^a Shinji Ozaki^a Norichika Iga^b Hideyuki Nishi^b
Nobukazu Fujimoto^c

^aDepartment of Respiratory Medicine, Okayama Rosai Hospital, Okayama, Japan;

^bDepartment of Surgery, Okayama Rosai Hospital, Okayama, Japan; ^cDepartment of Medical Oncology, Okayama Rosai Hospital, Okayama, Japan

Keywords

EGFR-mutated non-small cell lung cancer · Osimertinib · Rebiopsy · T790M · Thoracoscopy

Abstract

A 70-year-old woman underwent right upper lobectomy for adenocarcinoma of the lung (pT1bN2M0 stage IIIA). Five years after the surgery, lymph node recurrence was detected. Gefitinib was administered because epidermal growth factor (EGFR) exon 19 deletion mutation was detected in the previously resected surgical specimen. After a treatment of first-generation EGFR tyrosine kinase inhibitors, an FDG-PET/CT scan demonstrated abnormal FDG uptake in the pleura indicating pleural dissemination. Pleurocentesis revealed tumor cells in the pleural fluid; however, EGFR mutation testing failed due to inadequate tumor cellularity. Thoracoscopy under local anesthesia revealed multiple nodules on the parietal pleura. A biopsy specimen confirmed the diagnosis of lung adenocarcinoma with pleural dissemination and revealed EGFR exon 20-T790M mutation.

© 2019 The Author(s)
Published by S. Karger AG, Basel

Dr. Nobukazu Fujimoto
Department of Medical Oncology, Okayama Rosai Hospital
1-10-25 Chikkomidorimachi,
Minamiku, Okayama 7028055 (Japan)
E-Mail nobufujimoto@gmail.com

Introduction

Epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors (TKIs) improved the survival of patients with EGFR-mutated non-small cell lung cancer [1–3]. However, most of the patients developed resistance to EGFR-TKIs during treatment. EGFR exon 20-T790M (T790M) mutation develops as one of the mechanisms of acquired resistance in about 50% of patients treated with first-generation EGFR-TKIs [4, 5].

Case Report

A 70-year-old woman underwent right upper lobectomy for adenocarcinoma of the lung (pT1bN2M0 stage IIIA) followed by adjuvant chemotherapy with paclitaxel and carboplatin. Five years after the surgery, lymph node recurrence was detected in the mediastinum (#4R) by fluorodeoxyglucose-positron emission tomography (FDG-PET)/computed tomography (CT). Gefitinib was administered because EGFR exon 19 deletion mutation was detected in the previously resected surgical specimen. Although the lymph node shrank, the patient was switched from gefitinib to afatinib 7 months later because of hepatotoxicity. After 9 months of administration, afatinib was discontinued due to anorexia and skin rash. After another 8 months, multiple brain metastases developed and afatinib was reinstituted. After 8 months of afatinib administration, an FDG-PET/CT scan demonstrated abnormal FDG uptake in the pleura indicating pleural dissemination (Fig. 1a). Pleurocentesis revealed tumor cells in the pleural fluid; however, EGFR mutation testing failed due to inadequate tumor cellularity. Therefore, the patient underwent thoracoscopy under local anesthesia (TULA), which revealed multiple nodules on the parietal pleura (Fig. 1b). A drainage tube was placed into the pleural space via a port site and the tube was removed after pleurodesis with talc. Post-operative bleeding or pneumothorax was not observed.

A biopsy specimen confirmed the diagnosis of lung adenocarcinoma with pleural dissemination and revealed T790M mutation. Osimertinib treatment was started 2 weeks after TULA. Right pleural effusion was decreased, and serum carcinoembryonic antigen levels significantly declined (2,773.5 to 622.6 ng/mL) after 2 months of treatment.

Discussion

The third-generation EGFR-TKI, osimertinib, is recommended for patients who develop T790M mutation [6]. EGFR testing of the resistant tumor to detect T790M mutation is essential for the administration of osimertinib. Kawamura et al. [7] reported the procedures of obtaining tumor tissues for EGFR testing. They reported that among the 75 cases, procedures included 30 thoracocenteses, 24 transbronchial biopsies, 13 CT-guided needle biopsies, and 8 other procedures. None of the patients underwent TULA. Ichihara et al. [8] also reported 55 rebiopsy cases and TULA was also not performed for any patient. To the best of our knowledge, this is the first report of T790M mutation detection by TULA.

It is often difficult to obtain enough tumor tissues for T790M mutation testing because of the peripheral location or small size of the tumor. In the current case, pleurocentesis was first performed to obtain tumor samples in the pleural fluid. However, T790M mutation testing failed because of inadequate tumor cellularity. There was no target region with a large enough size to perform transbronchial or percutaneous needle biopsy, so we selected TULA as the procedure for rebiopsy. TULA has been established as a safe and effective diagnostic procedure for pleural diseases or pleural effusion of unknown etiology [9, 10]. The advantage of this

Fig. 1. PET/CT image demonstrating the abnormal uptake of FDG on the right side of the pleura (a), and thoroscopic exploration revealing multiple elevated lesions on the parietal pleura (b).

method is the ability to directly visualize the biopsy target. In addition, chest tube placement via a port site can be used for both drainage of the effusion and for pleurodesis. In the current case, pleurodesis was performed with talc through the chest tube after rebiopsy and drainage of the effusion. All of these procedures were completed without complication.

Liquid biopsy attracts a lot of attention as a minimally invasive method for T790M mutation testing, and many findings have been obtained in this field. Currently, liquid biopsy is only recommended by Japanese Lung Cancer Society guidelines in cases for which rebiopsy has failed or is unavailable. In fact, there is a 30% false-negative rate for T790M detection by liquid biopsy [11]. Tumor biopsy and repeated T790M testing should be considered to confirm the presence or absence of T790M mutation using transbronchial biopsy, percutaneous needle biopsy, or TULA, particularly in cases with pleural effusion and/or pleural dissemination.

Further development of genetic testing for the personalization of chemotherapy will increase the importance of obtaining enough tumor samples. For this purpose, it is extremely important to select the appropriate biopsy target and procedure. TULA may be a useful option as one of the procedures.

Statement of Ethics

The patient gave written informed consent to publish the case.

Disclosure Statement

The authors have no conflicting interests to declare.

Funding Sources

This work is supported by grants-in-aid from the Ministry of Health, Labor and Welfare, Japan.

Author Contributions

A. Matsuda designed the study and drafted the article. Y. Fuchimoto, S. Wada, N. Iga, and H. Nishi provided clinical samples and provided scientific advice. Y. Takigawa, S. Mitsumune, T. Tanaka, T. Takeguchi, Y. Miyamoto, and S. Ozaki provided scientific advice. N. Fujimoto supervised the entire project and completed the article.

References

- 1 Maemondo M, Inoue A, Kobayashi K, Sugawara S, Oizumi S, Isobe H. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med*. 2010 Jun 24;362(25):2380–8.
- 2 Mitsudomi T, Morita S, Yatabe Y, Negoro S, Okamoto I, Tsurutani J. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol*. 2010 Feb;11(2):121–8.
- 3 Sequist LV, Bell DW, Lynch TJ, Haber DA. Molecular predictors of response to epidermal growth factor receptor antagonists in non-small-cell lung cancer. *J Clin Oncol*. 2007 Feb 10;25(5):587–95.
- 4 Yu HA, Arcila ME, Rekhtman N, Sima CS, Zakowski MF, Pao W. Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers. *Clin Cancer Res*. 2013 Apr 15;19(8):2240–7.
- 5 Nguyen KS, Kobayashi S, Costa DB. Acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in non-small-cell lung cancers dependent on the epidermal growth factor receptor pathway. *Clin Lung Cancer*. 2009 Jul;10(4):281–9.
- 6 Mok TS, Wu YL, Ahn MJ, Garassino MC, Kim HR, Ramalingam SS. Osimertinib or platinum-pemetrexed in EGFR T790M-positive lung cancer. *N Engl J Med*. 2017 Feb 16;376(7):629–40.
- 7 Kawamura T, Kenmotsu H, Taira T, Omori S, Nakashima K, Wakuda K. Rebiopsy for patients with non-small-cell lung cancer after epidermal growth factor receptor-tyrosine kinase inhibitor failure. *Cancer Sci*. 2016 Jul;107(7):1001–5.
- 8 Ichihara E, Hotta K, Kubo T, Higashionna T, Ninomiya K, Ohashi K. Clinical significance of repeat rebiopsy in detecting the EGFR T790M secondary mutation in patients with non-small cell lung cancer. *Oncotarget*. 2018 Jun 29;9(50):29525–31.
- 9 Gao BA, Zhou G, Guan L, Zhang LY, Xiang GM. Effectiveness and safety of diagnostic flexi-rigid thoracoscopy in differentiating exudative pleural effusion of unknown etiology: a retrospective study of 215 patients. *J Thorac Dis*. 2014 May;6(5):438–43.
- 10 Murthy V, Bessich JL. Medical thoracoscopy and its evolving role in the diagnosis and treatment of pleural disease. *J Thorac Dis*. 2017 Sep;9(Suppl 10):S1011–S21.
- 11 Oxnard GR, Thress KS, Alden RS, Lawrance R, Paweletz CP, Cantarini M. Association between plasma genotyping and outcomes of treatment with osimertinib (AZD9291) in advanced non-small-cell lung cancer. *J Clin Oncol*. 2016 Oct 1;34(28):3375–82.