The NASA Twins Study: The Effect of One Year in Space on Long-Chain Fatty Acid Desaturases and Elongases

Michael A. Schmidt^{a, b} Cem Meydan^{d, e} Caleb M. Schmidt^{b, c}
Ebrahim Afshinnekoo^{d–f} Christopher E. Mason^{d–g}

^aAdvanced Pattern Analysis & Countermeasures Group, Boulder, CO, USA; ^bSovaris Aerospace, Boulder, CO, USA; ^cDepartment of Systems Engineering, Colorado State University, Fort Collins, CO, USA; ^dDepartment of Physiology and Biophysics, Weill Cornell Medicine, New York, NY, USA; ^eThe HRH Prince Alwaleed Bin Talal Bin Abdulaziz Alsaud Institute for Computational Biomedicine, Weill Cornell Medicine, New York, NY, USA; ^fThe WorldQuant Initiative for Quantitative Prediction, Weill Cornell Medicine, New York, NY, USA; ^gThe Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, New York, NY, USA

Keywords

NASA · Spaceflight · Fatty acids · Desaturase · Elongase · Eicosapentaenoic acid · Docosahexaenoic acid · Arachidonic acid · International Space Station · NASA Twins Study · Multiscale omics

Abstract

Background: At present, there is no clear understanding of the effect of long-duration spaceflight on the major enzymes that govern the metabolism of omega-6 and omega-3 fatty acids. To address this gap in knowledge, we used data from the NASA Twins Study, which includes a multiscale omics investigation of the changes that occurred during a year-long (340 days) human spaceflight. Embedded within the NASA Twins data are specific analytes associated with fatty acid metabolism. **Objectives:** To examine the long-chain fatty acid desaturases and elongases in a single human during 1 year in space. **Method:** One male twin was on board the International Space Station (ISS) for 1 year, while his monozygotic twin served as a genetically matched ground control.

Longitudinal assessments included the genome, epigenome, transcriptome, proteome, metabolome, microbiome, and immunome during the mission, as well as 6 months before and after. The gene-specific fatty acid desaturase and elongase transcriptome data (FADS1, FADS2, ELOVL2, and ELOVL5) were extracted from untargeted RNA-seq measurements derived from white blood cell fractions. **Results:** Most data from the elongases and desaturases exhibited relatively similar expression profiles ($R^2 > 0.6$) over time for the CD8, CD19, and lymphocyte-depleted (LD) cell fractions, indicating overall conservation of function within and between the subjects. Both cell-type and temporal specificity was observed in some cases, and some differences were also apparent between the polyadenylated (polyA) fraction of processed RNAs versus the ribodepleted (ribo-) fraction. The flight subject showed a stronger enrichment of the fatty acid metabolic process pathway across almost all cell types (col-

Michael A. Schmidt
Sovaris Aerospace
1942 Broadway Avenue, Suite 406
Boulder, CO 80302 (USA)
mschmidtphd@patternanalysis.org

Christopher E. Mason
Weill Cornell Medicine
1305 York Avenue, Y13-05
New York, NY 10021 (USA)
chm2042@med.cornell.edu

umns, CD4, CD8, CPT, and LD), most especially in the ribodepleted fraction of RNA, but also with the polyA⁺ fraction of RNA. Gene set enrichment analysis (GSEA) measures across three related fatty acid metabolism pathways showed a differential between the ground and the flight subject. **Conclusions:** There appears to be no persistent alteration of desaturase and elongase gene expression associated with 1 year in space. However, these data provide evidence that cellular lipid metabolism can be responsive and dynamic to spaceflight, even though it appears cell-type and context specific, most notably in terms of the fraction of RNA measured and the collection protocols. These results also provide new evidence of mid-flight spikes in expression of selected genes, which may indicate transient responses to specific insults during spaceflight.

© 2020 The Author(s)
Published by S. Karger AG, Basel

Introduction

Future human exploration missions to Mars will include transit to and from the planet, with eventual exploration of the planet surface. These will be preceded or paralleled by lunar orbital or lunar colony missions of extended duration. The safe execution of such missions, including some lasting up to 500 days, will require a more detailed understanding of human physiology and aerospace medicine than has been available to date. Presently, only 4 humans have participated in spaceflight missions of 1 year or more. While our understanding of the biological consequences of 4- to 6-month missions on the International Space Station (ISS) has increased substantially [1], there is almost no experience with exploration-class spaceflight. In addition, comprehensive molecular analytics, such as the multiscale omics studies now common on Earth, have not previously been applied to humans on long-duration missions in space.

Longitudinal measures of the genome, epigenome, transcriptome, proteome, metabolome, and microbiome in space represent a considerable advancement in the ability to detect patterns of variance in molecular networks that are related to spaceflight. These new multiscale omics platforms and metrics are also expected to aid in the identification of actionable biomarkers and targets from which to develop spaceflight countermeasures. Such knowledge would further allow space programs to develop truly personalized interventions, which would represent an evolution in spaceflight medicine [2].

There are presently no studies reporting on the biological response of long-chain polyunsaturated fatty acid

(PUFA) desaturase and elongase enzymes, and their associated metabolites, in astronauts who enter the space environment. Long-chain PUFA represent an important class of molecules to study in spaceflight, because they shape the basic architecture of the lipid bilayer in all cell membranes. This includes shaping the membrane architecture of neurons, synapses, the retina, and other highly electrically active elements of the brain [3]. It is noteworthy that the dry weight of the human brain is roughly 60% fat with an important dependence upon 20- and 22-carbon long-chain PUFA, especially docosahexaenoic acid (DHA) [4]. Given that the brain is increasingly being considered as one of the most vulnerable organs in the environment of space, understanding how the brain maintains and remodels the lipid fractions important to its architecture is likely among the more pressing items in spaceflight medicine [5].

Beyond structure, these long-chain fatty acids act as crucial signaling molecules, regulators of inflammation, and pro-resolving mediators in the recovery process following injury [6]. Such mediators have a strong regulatory influence on systems relevant to space, such as the cardiovascular system [7]. Muscle catabolism and bone loss are two primary phenotypic changes that occur on exposure to microgravity in space. The role of omega-3 fatty acids, in particular eicosapentaenoic acid (EPA), has been extensively studied on Earth, with limited but substantive examination in space. Zwart et al. [1] have investigated the relationship between the intake of omega-3 fatty acids and bone resorption markers in three models: (1) a cell culture system that models weightlessness, (2) *N*-telopeptide levels in healthy volunteers during 60 days of bed rest [1], and (3) the relationship between fish (and EPA) intake and *N*-telopeptide levels in astronauts on missions of 4–6 months on the ISS. The cell, bed rest, and flight data all support the hypothesis that EPA intake inhibited osteoclastogenesis. These models showed that the effect of EPA may be mediated, in part, via the NF- κ B pathway [6]. Disuse atrophy of muscle is also associated with NF- κ B activation, which similarly appears responsive to the modulating effect of omega-3 fatty acids [8].

Since selected fatty acids are essential and cannot be synthesized by the human body, understanding the dynamics of fatty acid metabolism in space is fundamental to forming proper exploration mission dietary regimens and clinical therapeutics [9]. Characterizing genetic variants of fatty acid-metabolizing enzymes in an astronaut would further inform the application of individualized nutritional approaches in order to prevent specific fatty acid deficits, while optimizing for mission performance

Fig. 1. Role of elongases and desaturases in the synthesis of long chain polyunsaturated fatty acids in humans. ELOVL, elongation of very long chain fatty acids protein; FADS, fatty acid desaturase; PG, prostaglandins; LT, leukotrienes; TBX, thromboxanes.

and health [10]. The fatty acids linoleic acid ($18:2n-6$) and α -linolenic acid ($18:3n-3$) are the two essential PUFA that, by definition, cannot be synthesized by humans and must be obtained from the diet. Their conversion into longer-chain fatty acids (Fig. 1) is governed by a series of steps that extend the fatty acid chain length by 2-carbon increments (elongation) and increase unsaturation by inserting double bonds (desaturation). Impairment in the elongation or desaturation enzymes can limit precursor fatty acid conversion, thereby causing accumulation of precursor fatty acids or causing longer-chain fatty acids to become *conditionally essential*, thus requiring direct dietary supplementation or other interventions [11].

Fatty acid desaturases (FADS) are enzymes key to human metabolism that remove two hydrogen atoms from a fatty acid, creating a carbon-carbon double bond. FADS1 ($\Delta 5$ -desaturase) facilitates the direct conversion of omega-6 $20:3n-6$ (DGLA) into $20:4n-6$ (AA). FADS1 also facilitates the direct conversion of omega-3 $20:4n-3$ (ETA) into $20:5n-3$ (EPA). FADS2 ($\Delta 6$ -desaturase) facilitates the direct conversion of omega-6 $18:2n-6$ (LA) into $18:3n-6$ (GLA). FADS2 also facilitates the direct conversion of omega-3 $18:3n-3$ (ALA) into $18:4n-3$ (SDA). Additionally, FADS2 converts $24:4n-6$ into $24:5n-6$ and $24:5n-4$ into $24:6n-3$ prior to the final beta-oxidation step to produce $22:5n-6$ and $22:6n-3$, respec-

Fig. 2. Gene expression profile for FADS1. Shown are the line plot expression data for fatty acid desaturase 1 (FADS1), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue). L, launch; FD, flight duration; R, return; FPKM, fragments per kilobase million.

tively. Collectively, these are the primary desaturation steps in converting PUFA precursors into their long-chain end products [12].

In addition, human elongases (ELOVL) are enzymes that extend the chain length of fatty acids by adding 2-carbon units during specific steps of fatty acid synthesis. There are two primary elongases in long-chain PUFA metabolism. These are elongase 2 (ELOVL2) and elongase 5 (ELOVL5). ELOVL5 facilitates the direct elongation of omega-6 18:3 n -6 (GLA) to 20:3 n -6 (DGLA). ELOVL5 also facilitates the direct elongation of omega-3 18:4 n -3 (SDA) to 20:4 n -3 (ETA). ELOVL2 facilitates the direct elongation of omega-6 20:4 n -6 (AA) to 22:4 n -6 (adrenic acid, DTA). ELOVL2 also facilitates the direct elongation of omega-3 20:5 n -3 (EPA) to 22:5 n -3 (DPA). It is important to note that ELOVL5 can also participate

in this conversion into omega-6 22:4 n -6 (adrenic acid, DTA) and omega-3 22:5 n -3 (DPA). Similarly, ELOVL2 acts in a second step of the pathway to elongate 22:4 n -6 to 24:4 n -6 and 22:5 n -3 to 24:5 n -3 [13, 14].

The NASA Twins Study examined a wide range of molecular dynamics in space for a pair of identical twins (one on the ISS and one on Earth). Data were generated from specimens collected over a 25-month period, including before (preflight), during (inflight), and after (postflight) spaceflight. The resulting data represent NASA's first multiscale omics profile of astronauts [15]. From this high-dimensional data set, we report for the first time on the targeted transcriptome of the genes that govern long-chain PUFA elongation and desaturation, and consider the implications for long-duration spaceflight.

Fig. 3. Gene expression profile for FADS1. Shown are the box plot expression data for fatty acid desaturase 1 (FADS1), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue).

Materials and Methods

Sample Collection

A pair of monozygotic twins were studied for 25 months, during which one subject (flight subject, TW) spent 340 days aboard the ISS, while his identical twin (ground subject, HR) remained on Earth. The subjects were male and aged 50 years at the beginning of the study. Both subjects had experienced different amounts of spaceflight exposure prior to the study (flight subject: 180 days total; ground subject: 54 days total). Multiple blood samples were collected ($n_{\text{flight}} = 19$; $n_{\text{ground}} = 12$) from both subjects, beginning approximately 6 months prior to the launch date, during the 340 days aboard the ISS, and 6 months after return.

Transcriptome samples were either mixed by inversion and immediately frozen at -80°C (for both Earth and ISS collections), or were freshly collected, sorted, and processed. Fresh samples collected aboard the ISS were returned to Earth in the Soyuz capsule

and kept at 4°C , until processing (approx. 35 h from collection to processing). Fresh samples were sorted into CD4, CD8, and CD19 cell selection and the remaining lymphocyte-depleted (LD) fraction, while frozen samples were processed to isolate peripheral blood mononuclear cells (PBMC) from the CPT vacutainers. RNA was isolated from the samples and sequenced with RNA-seq using two separate library preparation methods (polyadenylated [polyA] fraction selection and ribodepleted RNA). Blood sample collection, sorting, RNA extraction, quality control, library preparation, and sequencing were performed as previously described [15].

All of the samples freshly collected aboard the ISS from the flight subject were returned on the Soyuz capsule. Therefore, samples derived from ISS habitation were confounded by ambient return to Earth. To correct for the effects of ambient return on gene expression, blood samples were collected from an unrelated individual, which were then exposed to conditions that simulated the fresh and ambient return collections. The fresh and ambient sam-

Fig. 4. Gene expression profile for FADS2. Shown are the line plot expression data for fatty acid desaturase 2 (FADS2), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue). L, launch; FD, flight duration; R, return; FPKM, fragments per kilobase million.

ples from the control individual were processed and sequenced with the same protocol used for the study subjects. Data generated from the control individual were used to model and correct the effects of ambient return, using ComBat [16, 17] or multivariate analyses in differential expression, as previously described [15].

Data Processing

Adapter removal was performed using Cutadapt and Trim Galore! (v0.4.1) [18, 19]. Reads were aligned with a STAR transcriptome aligner (v2.5.1b), using two-pass alignment to hg19 genome and GENCODE v19 gene reference [20, 21]. Read quantification was performed, using Subread featureCounts and kallisto on ENSEMBL transcripts [22–24]. Transcript expression levels were summarized into their respective genes.

Differential gene expression analysis was performed using DESeq2 on gene counts [25]. Comparisons were done either within the flight subject only, e.g., comparing samples collected preflight to those collected in flight, or in a model that included samples from the ground subject to model the normal baseline variance

during the study duration. Ambient and fresh samples from the control individual were also included as a term in the multivariate model for any comparison that included ambient return samples. Each cell type was treated separately. Two different library preparations for polyA+ and rRNA depletion were combined together with an added covariate in the multivariate model. All differential gene expression analyses were performed as previously described [13]. Gene set enrichment analysis (GSEA) was performed using fgsea by ranking the genes on the resulting Wald statistic from DESeq2 and calculating enrichment on the Molecular Signatures Database (MSigDB) entries [26–28].

For normalized expression, gene counts were normalized using the trimmed mean of M [29]. ComBat was applied to gene expression values including the fresh and ambient samples from the control subject to correct for the potential effect of ambient return on gene expression. The resulting normalized and corrected gene expression values were used for visualization and gene set variation analysis (GSVA) [30]. GSVA was performed for each cell type and library preparation method separately using the corrected gene expression values for MSigDB signatures.

Fig. 5. Gene expression profile for FADS2. Shown are the box plot expression data for fatty acid desaturase 2 (FADS2), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue).

Results

We examined the gene expression profiles of four key elongases and desaturases, specifically FADS1, FADS2, ELOVL2, and ELOVL5, as well as their overall biological pathways, represented from GSEA and GSVA. These are represented by linear gene plots from before, during, and after the mission for the flight subject (TW), and compared to the ground control (HR) box plots summarizing the variance of the flight intervals and enrichment heat maps for the pathways.

Across these data, we observed several trends. First, most data from the elongases and desaturases exhibited relatively similar expression profiles ($R^2 > 0.6$) over time for the CD8, CD19, and LD cell fractions, indicating over-

all conservation of function within and between the subjects. Yet, both cell-type and temporal specificity was observed in some cases, and some differences were also apparent between the polyA fraction of processed RNAs and the ribodepleted (ribo-) fraction, which represent more nascent transcripts. For example, the bulk blood collection of PBMC from the CPT tubes showed a mid-mission, flight-specific increase in FADS1 (Fig. 2, 3) in both RNA fractions. Yet, other genes showed more “spikes” in their expression, indicating a transient response associated with spaceflight or sensitivity to other biological influences. This includes FADS2 (Fig. 4, 5), most notably in the polyA fraction of RNA, an increase in ELOVL2 during flight (Fig. 6, 7) in the polyA fraction, and a decrease in ELOVL5 during flight (Fig. 8, 9) in the polyA fraction.

Fig. 6. Gene expression profile for ELOVL2. Shown are the line plot expression data for elongation of very long chain fatty acids protein 2 (ELOVL2), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue). L, launch; FD, flight duration; R, return; FPKM, fragments per kilobase million.

To discern if related genes in these biochemical processes were also responsive to spaceflight, we examined the GSEA and GSVA enrichments for fatty acid metabolic pathways. The flight subject showed a stronger enrichment across almost all cell types (columns, CD4, CD8, CPT, and LD) for the fatty acid metabolic process pathway, most especially in the ribodepleted fraction of RNA, but also with the polyA+ fraction of RNA (Fig. 10). Also, the enrichment was nearly always above that of the ground control. Also, we used the GSEA enrichment measures across the related Fatty Acid Metabolism pathways, including Gene Ontology sets for “fatty acid derivative biosynthetic processes,” “phosphate-containing compound metabolic processes,” and “regulation of cell differentiation.” All three of these Gene Ontology sets showed a differential between the ground control and the

flight subject, with a notable shift in the third flight sample (Fig. 11). These represent both key process genes and related networks of genes that are dynamic in spaceflight.

Discussion and Conclusions

The NASA Twins Study represents the first integrated, multiscale omics, molecular, physiological, and cognitive portrait of an astronaut engaged in a year-long space mission, which revealed myriad phenotypic responses of the human body. Our results are the first to demonstrate transient gene expression profiles of desaturases and elongases in space, through examination of the transcriptome of FADS1, FADS2, ELOVL2, and ELOVL5 derived from untargeted RNA-seq analysis. Based on the tran-

Fig. 7. Gene expression profile for ELOVL2. Shown are the box plot expression data for elongation of very long chain fatty acids protein 2 (ELOVL2), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue).

scriptome analysis presented herein, there appears to be no persistent alteration of these fatty acid-metabolizing enzymes associated with 1 year in space, even though the second half of the mission (months 6–12) showed more differentially expressed genes (15). However, we observed that the enrichment of fatty acid metabolic pathway processes was already “primed” before spaceflight, and there were variable expression patterns across cell types. These patterns were isolated from CD4, CD8, and CD19 cell types, the remaining LD fraction, and PBMC.

These data provide evidence that cellular lipid metabolism can be responsive and dynamic during spaceflight, and that it can also be quite cell-type and context specific, most notably in terms of the fraction of RNA measured and the collection protocols. This is consistent with other

studies that have examined expression patterns by cell type [31–33]. These results also provide new evidence of mid-flight spikes in expression of these key genes, analogous to those observed for DNA repair and immune activation, which may indicate transient responses to specific radiation, vascular, or physiologic insults during spaceflight. Moreover, the enrichment and differential pathways utilized as part of fatty acid metabolism also suggest potential new mechanisms of adaptation by the body during long-duration spaceflight.

Several limitations of this study warrant consideration. First, the small number of subjects and the large number of variables (analytes) render such high-dimensional data prone to overfitting. One method to address this limitation is through serial measures of the baseline,

Fig. 8. Gene expression profile for ELOVL5. Shown are the line plot expression data for elongation of very long chain fatty acids protein 5 (ELOVL5), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue). L, launch; FD, flight duration; R, return; FPKM, fragments per kilobase million.

inflight, and postflight conditions, which was the approach used in this study. Second, the cohort of males is limiting on the generalizability of these results to females. Nonetheless, the matched cohort favors homogeneity, which is valuable in studies involving high-dimensional data. Third, the data reported herein do not contain SNP, indel, CNV, and structural genomic elements for reasons of confidentiality. In particular, SNP variants of FADS1, FADS2, ELOVL2, and ELOVL5 would have been additionally informative, as their impact on the conversion of precursor fatty acids into downstream metabolites has been characterized [15]. Thus, this analysis provides some insight into changes in molecular functional elements in the space environment, but it does not include an examination of structural genomic elements that might alter human risk in the space environment.

Fourth, we were unable to describe the precise relationship between FADS, elongases, and their fatty acid metabolites in space. In the original *Science* paper describing the first results of the Twins Study, it was reported that the omega-6 20:4 fatty acid (described as arachidonic acid [AA]) increased, while lysophospholipids containing the omega-3 20:5 fatty acid (EPA) decreased in space [34]. While this represents a potentially important finding, our subsequent analysis revealed that AA was not individually isolated, as 20:4 consists of more than one isomer. More specifically, the 20:4 (eicosatetraenoic acid [ETA]) measurement that was reported in the original methods did not differentiate between the omega-6 and omega-3 isomers of ETA (20:4), which includes the omega-6 ETA (*all-cis*-5,8,11,14-eicosatetraenoic acid) and the omega-3 ETA (*all-cis*-8,11,14,17-eicosatetraenoic acid).

Fig. 9. Gene expression profile for ELOVL5. Shown are the box plot expression data for elongation of very long chain fatty acids protein 5 (ELOVL5), including the polyA+ and ribodepleted specimens for various timepoints and cell types (CD4, CD8, CD19, lymphocyte-depleted [LD] fraction, and CPT [peripheral blood mononuclear cells]), representing ground (green) and flight (blue).

Thus, while it is likely that AA may have increased (as AA commonly predominates over ETA), the percent composition of AA in relation to ETA could not be established, and thus warrants higher-resolution methods in future studies.

Likewise, in reporting the 22:5 as docosapentaenoic acid (DPA), the analysis did not differentiate between the two isomers of DPA, which includes the omega-6 DPA (*all-cis*-4,7,10,13,16-docosapentaenoic acid) and the omega-3 DPA (*all-cis*-7,10,13,16,19-docosapentaenoic acid). Similarly, in reporting the 18:3 as linolenic acid, the analysis did not differentiate between the two isomers of linolenic acid, which includes the omega-6 γ -linolenic acid (*all-cis*-6,9,12-octadecatrienoic acid) and the omega-3 α -linolenic acid (*all-cis*-9,12,15-octadecatrienoic acid).

While a general estimate of the omega-6 trajectory in relation to omega-3 fatty acids was reported in the original *Science* paper, we were unable to complete a specific analysis of individual fatty acid metabolites in relation to elongase and desaturase enzymes because of insufficient data surrounding the identity of individual fatty acid isomers.

Our ongoing work, using the extant data set, is examining the elongase and desaturase proteins, SNPs in elongase and desaturase genes, and epigenomic changes in genes that govern fatty acid metabolism, derived from the multiscale omics analysis of 1 year in space. This work also explores fatty acid molecular dynamics in relation to various clinical phenotypes. Included is an examination of phenotypes that were observed to have changed during the 1-year mission, such as cognitive, mitochondrial, im-

Fig. 10. Gene set variation analysis (GSVA) pathway enrichment for fatty acid metabolic processes. The data for the flight subject (blue) showed a stronger enrichment across almost all cell types (columns, CD4, CD8, CPT, and lymphocyte-depleted [LD] fraction) for the fatty acid metabolic process pathway, most especially in the ribodepleted fraction of RNA (bottom row), but also with the polyA+ fraction of RNA (top row). L, launch; FD, flight duration; R, return.

munological, cardiovascular, ocular, and other phenotypes. Potential correlations between fatty acid trajectories and cognitive associations will be of particular interest, since (1) cognitive speed decreased for all tests (except for the visual object learning test) after 1 year in space, (2) accuracy decreased for 80% of the domains tested post-flight, and (3) fatty acids have previously been associated with changes in cognition [15].

In the future, new methods for direct sequencing and molecular assays that can be performed directly on the ISS can help reduce some of the confounding variables [35–37], as well as provide longer sequence reads and improved resolution of isoform changes [38]. The use of red blood cell fatty acid analysis is expected to provide a more stable measure of fatty acid status that is more reflective of tissue stores and shows less day-to-day variation associated with fatty acid intake. If serum or whole blood is used,

strict controls related to dietary intake and fasting blood samples should be employed. Isolation of individual fatty acid isomers within each group (e.g., 18:3 n -6/ n -3, 20:4 n -6/ n -3, 22:5 n -6/ n -3, etc.), using methods such as GC-FID (gas chromatography-flame ionization detection) or GC-MS (mass spectrometry), will add further precision. Incorporating the analysis of eicosanoids, docosanoids, and pro-resolving mediators (downstream metabolites of long-chain PUFA) will further elucidate an important class of signaling and regulatory derivatives of fatty acids. New dietary monitoring methods, tracking of dietary supplement intake, and preflight quantification of the ISS foods can further aid in the identification of potential drivers of variations in fatty acid molecular networks. Since double-bonded fatty acids are uniquely sensitive to oxidation, such as that initiated by ionizing radiation, improved ISS radiation monitoring would be of benefit.

Fig. 11. Fatty acid metabolism Gene Ontology (GO) differences. Plotted is the enrichment (positive = orange, negative = purple) for specific fatty acid pathways (rows) for various timepoints in the study (top row labels), including preflight (dark purple), flight (blue), postflight (green), and longitudinal ground control (brown). Ambient return samples are labelled in black. L, launch; FD, flight duration; R, return; GD, ground duration.

The degree to which the present study of 1 year in low Earth orbit accurately reflects the biological dynamics that would emerge from a 3-year mission to Mars or prolonged lunar habitation is currently unknown. Therefore, the future support of exploration-class missions will benefit from additional multiscale omics studies of longer duration and varying conditions.

Acknowledgements

We would like to thank the Epigenomics Core Facility at Weill Cornell Medicine, the Scientific Computing Unit (SCU), XSEDE Supercomputing Resources, and funding from the Irma T. Hirsch and Monique Weill-Caulier Charitable Trusts, The Bert L & N Kuggie Vallee Foundation, the WorldQuant Foundation, The Pershing Square Sohn Cancer Research Alliance, NASA (NNX14AH50G and NNX17AB26G), the National Institutes of Health (R01AI125416, R01MH117406, R01CA249054, R01AI151059, P01HD067244, and P01CA214274), TRISH (NNX16AO69A:0107 and NNX16AO69A:0061), the Bill & Melinda Gates Foundation (OPP1151054), and the Leukemia and Lymphoma Society (LLS) grants (LLS 9238-16, Mak, LLS-MCL-982, Chen-Kiang). We would like to thank the NASA JSC Nutritional Biochemistry Lab personnel, especially YaVonne Bourbeau, who led efforts on this project for project coordination, sample collection, processing, and analysis; the NASA Hu-

man Research Program's ISS Medical Projects element, Daniel Mollicone and Christopher Mott from Pulsar Informatics Inc., Emanuel Hermosillo, and Sarah McGuire for support on the cognition measures; Dr. Yael-Rosenberg Hasson from the Stanford Human Immune Monitoring Center for advice and support on cytokine profiling data; Keith Bettinger from Stanford University for support with the Stanford Twins data repository; Pawel R. Kiela, DVM, PhD, Daniel Laubitz, PhD (University of Arizona), and Eunjo Song (Northwestern University) for support on the microbiome project sample collection; and Jiwan "John" Kim for laboratory support in targeted metabolomics. We thank Jason X.-Y. Yuan, MD, PhD, at the University of Arizona, Tucson, for providing his support and laboratory for the Tucson sample collections. Our thanks to Michael Snyder and Tejas Mishra of Stanford University for untargeted metabolomic and fatty acid analysis. We thank Julian C. Schmidt, M.S. (Sovaris Aerospace), for manuscript support. Finally, we thank Robert Hubbard, EdD, MA (Sovaris Aerospace), for manuscript review and illustrations.

Statement of Ethics

The study participants have given their written informed consent. The study protocol has been approved by the research institutes' (both NASA and WCM) committee on human research, specifically IRB# 1309014347.

Disclosure Statement

C.E.M. is a cofounder and board member for Biotia, Inc., and Onegevity Health, Inc.

Funding Sources

The study was supported by NASA: NNX14AH51G (All Twins PIs); TRISH: NNX16AO69A:0107 and NNX16AO69A:0061 (C.E.M.); and The Bert L & N Kuggie Vallee Foundation, the WorldQuant Foundation, The Pershing Square Sohn Cancer Research Alliance, and the Bill & Melinda Gates Foundation (OPP1151054) for funding (C.E.M.).

Author Contributions

M.A.S. and C.E.M. conceived the overall study objectives; M.A.S. and C.E.M. led the manuscript writing, organization, and editing efforts; C.E.M. and C.M. processed the data, and C.M. performed the statistical analysis and multiomics data integration; M.A.S. led the molecular target selection and pathway elucidation; all authors performed the data interpretation, translation into biological meaning, summary of results, discussion, conclusion, and future directions.

References

- 1 Zwart SR, Pierson D, Mehta S, Gonda S, Smith SM. Capacity of omega-3 fatty acids or eicosapentaenoic acid to counteract weightlessness-induced bone loss by inhibiting NF- κ B activation: from cells to bed rest to astronauts. *J Bone Miner Res*. 2010 May;25(5):1049–57.
- 2 Schmidt MA, Goodwin TJ. Personalized medicine in human space flight: using Omics based analyses to develop individualized countermeasures that enhance astronaut safety and performance. *Metabolomics*. 2013; 9(6):1134–56.
- 3 Cholewski M, Tomczykowa M, Tomczyk M. A Comprehensive Review of Chemistry, Sources and Bioavailability of Omega-3 Fatty Acids. *Nutrients*. 2018 Nov;10(11):1662.
- 4 Chang CY, Ke DS, Chen JY. Essential fatty acids and human brain. *Acta Neurol Taiwan*. 2009 Dec;18(4):231–41.
- 5 Jandial R, Hoshida R, Waters JD, Limoli CL. Space-brain: the negative effects of space exposure on the central nervous system. *Surg Neurol Int*. 2018 Jan;9(1):9.
- 6 Bazinet RP, Layé S. Polyunsaturated fatty acids and their metabolites in brain function and disease. *Nat Rev Neurosci*. 2014 Dec; 15(12):771–85.
- 7 Calder PC. Omega-3 fatty acids and inflammatory processes: from molecules to man. *Biochem Soc Trans*. 2017 Oct;45(5):1105–15.
- 8 Khal J, Tisdale MJ. Downregulation of muscle protein degradation in sepsis by eicosapentaenoic acid (EPA). *Biochem Biophys Res Commun*. 2008 Oct;375(2):238–40.
- 9 Tocher DR, Betancor MB, Sprague M, Olsen RE, Napier JA. Omega-3 Long-Chain Polyunsaturated Fatty Acids, EPA and DHA: Bridging the Gap between Supply and Demand. *Nutrients*. 2019 Jan;11(1):89.
- 10 Mathias RA, Pani V, Chilton FH. Genetic Variants in the FADS Gene: Implications for Dietary Recommendations for Fatty Acid Intake. *Curr Nutr Rep*. 2014 Jun;3(2):139–48.
- 11 Burdge GC, Calder PC. Conversion of α -linolenic acid to longer-chain polyunsaturated fatty acids in human adults. *Reprod Nutr Dev*. 2005 Sep-Oct;45(5):581–97.
- 12 Zhang JY, Kothapalli KS, Brenna JT. Desaturase and elongase-limiting endogenous long-chain polyunsaturated fatty acid biosynthesis. *Curr Opin Clin Nutr Metab Care*. 2016 Mar; 19(2):103–10.
- 13 Jakobsson A, Westerberg R, Jacobsson A. Fatty acid elongases in mammals: their regulation and roles in metabolism. *Prog Lipid Res*. 2006 May;45(3):237–49.
- 14 Robichaud PP, Munganyiki JE, Boilard E, Surette ME. Polyunsaturated fatty acid elongation and desaturation in activated human T-cells: ELOVL5 is the key elongase. *J Lipid Res*. 2018 Dec;59(12):2383–96.
- 15 Garrett-Bakelman FE, Darshi M, Green SJ, Gur RC, Lin L, Macias BR, et al. The NASA Twins Study: a multidimensional analysis of a year-long human spaceflight. *Science*. 2019 Apr;364(6436):eaau8650.
- 16 Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics*. 2007 Jan;8(1):118–27.
- 17 Leek JT, Johnson WE, Parker HS, Fertig EJ, Jaffe AE, Storey JD, et al. sva: Surrogate Variable Analysis. R package version 3.30.1. 2019.
- 18 Martin M. Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads. *EMBnet.journal*. 2011;17(1):10–12.
- 19 Krueger F. *Trim Galore!* [accessed 2019 Oct 2]. 2016. Available from: https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/.
- 20 Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013 Jan;29(1):15–21.
- 21 Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, et al. GENCODE: the reference human genome annotation for The ENCODE Project. *Genome Res*. 2012 Sep;22(9):1760–74.
- 22 Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*. 2014 Apr;30(7):923–30.

Appendix: Glossary of Terms

Omega-6

LA: linoleic acid (18:2n–6)
GLA: γ -linolenic acid (18:3n–6)
DGLA: dihomo- γ -linolenic acid (20:3n–6)
AA: arachidonic acid (eicosatetraenoic acid; 20:4n–6)
DTA: docosatetraenoic acid (22:4n–6)
DPAn–6: docosapentaenoic acid (22:5n–6)
TTA: tetracosatetraenoic acid (24:4n–6)
TPAn–6: tetracosapentaenoic acid (24:5n–6)

Omega-3

ALA: α -linolenic acid (18:3n–3)
SDA: stearidonic acid (18:4n–3)
ETA: eicosatetraenoic acid (20:4n–3)
EPA: eicosapentaenoic acid (20:5n–3)
DPAn–3: docosapentaenoic acid (22:5n–3)
DHA: docosahexaenoic acid (22:6n–3)
TPAn–3: tetracosapentaenoic acid (24:5n–3)
THA: tetracosahexaenoic acid (24:6n–3)

Enzymes

FADS1: fatty acid desaturase 1
FADS2: fatty acid desaturase 2
ELOVL2: elongation of very long chain fatty acids protein 2
ELOVL5: elongation of very long chain fatty acids protein 5

- 23 Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol*. 2016 May;34(5):525–7.
- 24 Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, et al. Ensembl 2018. *Nucleic Acids Res*. 2018 Jan;46 D1:D754–61.
- 25 Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550.
- 26 Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA*. 2005 Oct;102(43):15545–50.
- 27 Sergushichev A. An algorithm for fast pre-ranked gene set enrichment analysis using cumulative statistic calculation. *bioRxiv*. 2016 Jun:060012.
- 28 Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP. Molecular Signatures Database (MSigDB) 3.0. *Bioinformatics*. 2011 Jun;27(12):1739–40.
- 29 Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol*. 2010;11(3):R25.
- 30 Hänzelmann S, Castelo R, Guinney J. GSEA: gene set variation analysis for microarray and RNA-seq data. *BMC Bioinformatics*. 2013 Jan;14(1):7.
- 31 Li S, Tighe SW, Nicolet CM, Grove D, Levy S, Farmerie W, et al. Multi-platform assessment of transcriptome profiling using RNA-seq in the ABRF next-generation sequencing study. *Nat Biotechnol*. 2014 Sep;32(9):915–25.
- 32 Li S, Labaj PP, Zumbo P, Sykacek P, Shi W, Shi L, et al. Detecting and correcting systematic variation in large-scale RNA sequencing data. *Nat Biotechnol*. 2014 Sep;32(9):888–95.
- 33 Reynolds LM, Howard TD, Ruczinski I, Kanchan K, Seeds MC, Mathias RA, et al. Tissue-specific impact of FADS cluster variants on FADS1 and FADS2 gene expression. *PLoS One*. 2018 Mar;13(3):e0194610.
- 34 Koletzko B, Reischl E, Tanjung C, Gonzalez-Casanova I, Ramakrishnan U, Meldrum S, et al. FADS1 and FADS2 Polymorphisms Modulate Fatty Acid Metabolism and Dietary Impact on Health. *Annu Rev Nutr*. 2019 Aug;39(1):21–44.
- 35 McIntyre AB, Rizzardi L, Yu AM, Alexander N, Rosen GL, Botkin DJ, et al. Nanopore sequencing in microgravity. *NPJ Microgravity*. 2016 Oct;2(1):16035.
- 36 Castro-Wallace SL, Chiu CY, John KK, Stahl SE, Rubins KH, McIntyre AB, et al. Nanopore DNA sequencing and genome assembly on the International Space Station. *Sci Rep*. 2017 Dec;7(1):18022.
- 37 McIntyre AB, Alexander N, Grigorev K, Bezdan D, Sichtig H, Chiu CY, et al. Single-molecule sequencing detection of N6-methyladenine in microbial reference materials. *Nat Commun*. 2019 Feb;10(1):579.
- 38 Chhangawala S, Rudy G, Mason CE, Rosenfeld JA. The impact of read length on quantification of differentially expressed genes and splice junction detection. *Genome Biol*. 2015 Jun;16(1):131.