Case and Review

Asymptomatic Colonic Anisakiasis: Is It So Rare?

Tetsuro Takasaki^a Toru Yamada^b Junji Kinoshita^a Yasuaki Motomura^a

^aDepartment of Gastroenterology, Tokyo Bay Urayasu Ichikawa Medical Center, Urayasu, Japan; ^bDepartment of Family Medicine, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Tokyo, Japan

Keywords

Adult · *Anisakis* · Colon · Colonoscopy · Case report

Abstract

Most cases of anisakiasis occur in the stomach. Colonic anisakiasis, especially when asymptomatic, is extremely rare. We report a case of asymptomatic colonic anisakiasis and present a literature review of cases of colonic anisakiasis. A 52-year-old man underwent colonoscopy for colorectal cancer screening. Although no colorectal neoplasm was found, an *Anisakis* larva was detected in the transverse colon and the larva was removed by grasping forceps. Our review of PubMed revealed that 40% of cases of colonic anisakiasis are asymptomatic. We conclude that asymptomatic colonic anisakiasis may not be as rare as conventionally assumed.

© 2020 The Author(s)
Published by S. Karger AG, Basel

Introduction

Anisakiasis is a zoonotic infection most frequently caused by *Anisakis simplex* and *Pseudoterranova decipiens*. Marine mammals are the natural host, and humans can become

incidental hosts by eating raw or undercooked seafood harboring *Anisakidae* larvae [1]. Anisakiasis is common in East Asia, especially in Japan, because of the dietary culture of eating raw fish. The stomach is the most common site of larval infection; the incidence of colonic anisakiasis is only 0.25% [2]. Without treatment, abdominal pain lasts from several weeks to 1 month, and more than half of colonic anisakiasis lesions are present on the right side [3]. Because of the nonspecific symptoms, such as abdominal pain, the differential diagnoses of symptomatic colonic anisakiasis are broad and include appendicitis, ileitis, diverticulitis, eosinophilic gastroenteritis, cholecystitis, colonic tumor, and inflammatory bowel disease [1]. Asymptomatic colonic anisakiasis is considered extremely rare [4, 5]. We report a case of asymptomatic anisakiasis and summarize previously reported cases of colonic anisakiasis regarding the presenting symptoms.

Case Presentation

A 52-year-old man underwent an immunochemical fecal occult blood test for a routine medical checkup at his local hospital. He had no symptoms and did not report eating any raw seafood. He was taking no medications. He had stopped smoking 1 year previously, but his alcohol consumption was in excess of 60 g/day. On physical examination, his temperature was 36.5°C (97.7°F), his blood pressure was 129/75 mm Hg, and his pulse rate was 75 beats/min. Abdominal examination revealed no masses or pain. Laboratory examinations were not performed because the patient was being screened specifically for colon cancer. The result of a fecal occult blood test was positive; consequently, he underwent colonoscopy, which identified an *Anisakis* nematode penetrating the wall of the transverse colon; the surrounding mucosa was slightly swollen (Fig. 1). The nematode was removed completely using grasping forceps, and no symptoms occurred following the colonoscopy.

Discussion and Conclusion

Asymptomatic colonic anisakiasis is considered rare because the number of case reports is low [5]. We reviewed previously reported cases of colonic anisakiasis in PubMed up to March 2019 using the keywords “*Anisakis*” and “colon” (Table 1) [2–15]. Papers that were not written in English and papers that contained no description of the patient’s symptoms were excluded. Fourteen patients were identified; 6/15 patients (40.0%, including our patient) were asymptomatic [3]. Among the 9 symptomatic patients, the most common symptom was abdominal pain ($n = 7$, 77.8%). The second most common symptoms were hematochezia ($n = 3$, 33.3%) and nausea ($n = 3$, 33.3%). Our review showed that asymptomatic colonic anisakiasis may account for a large percentage of all cases of colonic anisakiasis beyond the conventional assumption. Notably, the main limitation of our study is the small number of cases, which occurred because most cases of anisakiasis are reported in Japanese.

The symptoms of anisakiasis are caused by the immune response to the *Anisakis* larvae. *Anisakis* larvae secrete potent proteolytic enzymes, called excretory-secretory products, to invade the gastrointestinal mucosa. The larval antigens, including the excretory-secretory products, induce allergic reactions. Anisakiasis is classified into three types according to the

immune response: indolent-type anisakiasis, which is caused by an innate immune response; fulminant-type anisakiasis, which is caused by hypersensitivity and induces severe symptoms; and chronic anisakiasis. Initial infection can cause an indolent-type allergic reaction with no or only mild symptoms [5], as in the present case. Anisakiasis should not be left untreated in asymptomatic patients. Chronic anisakiasis caused by degenerating larvae can lead to the development of an abscess and/or granuloma [1]. In some cases, a granuloma caused by *Anisakis* larvae needed differentiation from a tumor [6, 7]. Complete removal of the larvae is strongly recommended when *Anisakis* is identified. Colonoscopy is not usually performed if a patient with colonic anisakiasis has no symptoms; therefore, asymptomatic colonic anisakiasis can be missed.

Anisakiasis rarely involves sites other than the stomach, and the incidence of asymptomatic colonic anisakiasis has been considered extremely low. However, our review of the worldwide literature showed that 40% of cases of colonic anisakiasis are asymptomatic. The consumption of raw fish has recently extended beyond Asia to become more globally widespread. As a result, we predict that the number of reports of asymptomatic colonic anisakiasis will increase.

Acknowledgment

We wish to acknowledge the assistance of Aki Miyagaki, Eriko Yamaguchi, Junji Kinoshita, Kento Fujioka, Risa Okamoto, and Tsubasa Yoshioka. We thank Jane Charbonneau, DVM, from Edanz Group (www.edanzediting.com/ac) for editing a draft of this paper.

Statement of Ethics

Written informed consent was obtained from the patient prior to the publication of this case report.

Conflict of Interest Statement

The authors have no competing interests to declare.

Funding Sources

The authors have no financial disclosures or funding to declare.

Author Contributions

T. Takasaki performed the literature review and wrote the manuscript. T. Yamada critically revised the manuscript and the literature review. J. Kinoshita performed the colonos-

copy. Y. Motomura reviewed and assisted in writing the manuscript. All authors approved the final version of the manuscript.

References

- 1 Hochberg NS, Hamer DH. Anisakidosis: perils of the deep. *Clin Infect Dis*. 2010 Oct;51(7):806–12.
- 2 Mineta S, Shimanuki K, Sugiura A, Tsuchiya Y, Kaneko M, Sugiyama Y, et al. Chronic anisakiasis of the ascending colon associated with carcinoma. *J Nippon Med Sch*. 2006 Jun;73(3):169–74.
- 3 Tsukui M, Morimoto N, Kurata H, Sunada F. Asymptomatic anisakiasis of the colon incidentally diagnosed and treated during colonoscopy by retroflexion in the ascending colon. *J Rural Med*. 2016;11(2):73–5.
- 4 Zullo A, Balsamo G, Baldini D, Francesco V, Manta R. Asymptomatic Anisakis and erosive lesions in the colon. *Ann Gastroenterol*. 2018 Mar–Apr;31(2):246.
- 5 Taniguchi G, Nagahara A, Matsumoto K, Ritsuno H, Igusa Y, Sasaki H, et al. Asymptomatic anisakiasis of the colon incidentally found by colonoscopy. *Clin J Gastroenterol*. 2011 Dec;4(6):371–3.
- 6 Yorimitsu N, Hiraoka A, Utsunomiya H, Imai Y, Tatsukawa H, Tazuya N, et al. Colonic intussusception caused by anisakiasis: a case report and review of the literature. *Intern Med*. 2013;52(2):223–6.
- 7 Hernandez-Prera JC, Polydorides AD. Anisakidosis of the sigmoid colon disguising as metastatic carcinoma: a case report and review of the literature. *Pathol Res Pract*. 2012 Jul;208(7):433–5.
- 8 Mladineo I, Popović M, Drmić-Hofman I, Poljak V. A case report of *Anisakis pegreffii* (Nematoda, Anisakidae) identified from archival paraffin sections of a Croatian patient. *BMC Infect Dis*. 2016 Feb;16(1):42.
- 9 Cavallero S, Scribano D, D'Amelio S. First case report of invasive pseudoterranoviasis in Italy. *Parasitol Int*. 2016 Oct;65(5 Pt A):488–90.
- 10 Amir A, Ngui R, Ismail WH, Wong KT, Ong JS, Lim YA, et al. Anisakiasis causing acute dysentery in Malaysia. *Am J Trop Med Hyg*. 2016 Aug;95(2):410–2.
- 11 Kim SH, Park CW, Kim SK, Won S, Park WK, Kim HR, et al. A case of anisakiasis invading the stomach and the colon at the same time after eating anchovies. *Clin Endosc*. 2013 May;46(3):293–6.
- 12 Herranz-Bachiller MT, Atienza-Sánchez R, Barrio-Andrés J, Alcaide-Suárez N, Ruiz-Zorrilla R, Sancho-Del-Val L, et al. Colonic polyp secondary to *Anisakis simplex*. *Rev Esp Enferm Dig*. 2012 Oct–Nov;104(10):554–5.
- 13 Do KR, Cho YS, Kim HK, Hwang BH, Shin EJ, Jeong HB, et al. Intestinal helminthic infections diagnosed by colonoscopy in a regional hospital during 2001–2008. *Korean J Parasitol*. 2010 Mar;48(1):75–8.
- 14 Higashi M, Tanaka K, Kitada T, Nakatake K, Tsuji M. Anisakiasis confirmed by radiography of the large intestine. *Gastrointest Radiol*. 1988;13(1):85–6.
- 15 Matsui T, Iida M, Murakami M, Kimura Y, Fujishima M, Yao Y, et al. Intestinal anisakiasis: clinical and radiologic features. *Radiology*. 1985 Nov;157(2):299–302.

Fig. 1. A 52-year-old man underwent colonoscopy during a routine medical checkup. An *Anisakis* larva was observed in the transverse colon.

Table 1. Published cases of colonic anisakiasis

Case No.	Year	Age, years/sex	Country	Detection site	Symptoms	Ref.
1	2018	50/M	Italy	sigmoid colon	none	4
2	2016	74/M	Japan	ascending colon	none	3
3	2016	65/F	Italy	ascending colon	nonspecific symptoms	8
4	2016	64/M	Malaysia	ascending colon	abdominal discomfort, hematochezia	9
5	2016	70/F	Croatia	sigmoid colon	abdominal pain, nausea	10
6	2013	47/M	Korea	transverse colon	abdominal pain, nausea	11
7	2013	67/F	Japan	ascending colon	abdominal pain	7
8	2012	47/F	Spain	rectum	hematochezia	12
9	2012	45/F	USA	sigmoid colon	hematochezia, anemia	6
10	2011	42/M	Japan	ascending colon	none	5
11	2010	49/M	Korea	ascending colon	none	13
12	2006	69/M	Japan	ascending colon	abdominal pain, diarrhea, urticaria	2
13	1988	37/M	Japan	ascending colon	abdominal pain, nausea, vomiting	14
14	1985	62/F	Japan	ascending colon	abdominal pain	15

M, male; F, female.