Can Diet Influence the COVID-19 Mortality Rate?

Amanda Avery

Division of Food, Nutrition & Dietetics, University of Nottingham, Sutton Bonington Campus, Leicestershire, UK

Abstract from Bousquet J, Anto JM, Iaccarino G, et al.: Is diet partly responsible for differences in COVID-19 death rates between and within countries? Clin Transl Allergy. 2020;10:16.

Keywords

Angiotensin-converting enzyme · Antioxidant · Coronavirus · Diet · Food

Abstract

Reported COVID-19 deaths in Germany are relatively low as compared to many European countries. Among the several explanations proposed, an early and large testing of the population was put forward. Most current debates on COVID-19 focus on the differences among countries, but little attention has been given to regional differences and diet. The low-death rate European coun-

tries (e.g. Austria, Baltic States, Czech Republic, Finland, Norway, Poland, Slovakia) have used different quarantine and/or confinement times and methods and none have performed as many early tests as Germany. Among other factors that may be significant are the dietary habits. It seems that some foods largely used in these countries may reduce angiotensin-converting enzyme activity or are anti-oxidants. Among the many possible areas of research, it might be important to understand diet and angiotensin-converting enzyme-2 (ACE2) levels in populations with different COVID-19 death rates since dietary interventions may be of great benefit.

© 2020 The Author(s)

Knowledge Transfer

Background

This commentary paper [1] considers why some European countries have much lower death rates from COVID-19 than other countries. The countries with notable low death rates include Austria, the Baltic States, Czech Republic, Finland, Norway, Poland, Slovakia and Germany. Of course, there could be a number of plausible reasons including the early and wide-scale testing adopted by Germany or some of the different approaches to quarantine adopted by the other countries in this list. Large differences in death rates exist within a country as well and, for example, whilst Bavaria started the earliest testing it remains the most affected region within Germany. However, the fact that these countries, and regions within, may have different dietary habits to countries and regions with higher death rates has been largely overlooked. It is increasingly being recognised that COVID symptoms are more prevalent and worse in people with sub-clinical inflammatory conditions such as obesity and type 2 diabetes [2], and there is some indication that vitamin D insufficiency may also be associated with an increased risk [3]. However, it could be that these countries with lower death rates from the virus may have specific dietary habits which may be protective. The authors of the commentary paper suggest that this may indeed be the case. This review will outline some of their theories in the context of our wider knowledge and the supporting evidence base.

Obesity as a Starting Point

The fact that obesity is a recognised risk factor for mortality does suggest that nutritional status may be an important factor in COVID-19 risk severity. Both obesity and type 2 diabetes have insulin resistance and impaired lipid metabolism as common features which lead to increased oxidative stress and inflammation [4]. Many foods have some level of anti-oxidant activity, particularly fruit and vegetables. Some dietary components have both anti-oxidative and anti-inflammatory properties and an example is resveratrol, a natural polyphenol found in red wine, rhubarb, fruits such as blueberries, cranberries, red grapes, and also in peanuts. From in vitro work it has been shown that resveratrol can significantly prevent Middle East Respiratory Syndrome coronavirus (MERS-CoV) and also prolong cellular survival after infection [5].

Dietary fatty acids, and particularly the omega-6 to omega-3 ratio, may influence the immune response in different situations with omega-3 polyunsaturated fatty acids (PUFAs) generally exhibiting a protective effect. For example, the airway epithelium is the first line of defence in the respiratory system and docosahexaenoic acid (DHA) supplementation has been shown to decrease pro-inflammatory cytokine production from the airway epithelium in the presence of an underlying inflammatory condition [6]. Supplements of omega-3 PUFAs may also help to reduce barrier permeability, mucus production and oxidative stress whilst improving healing in inflammatory conditions [7]. Oily fish and some vegetable oils are the best dietary sources of omega-3 PUFAs.

Angiotensin-Converting Enzyme 2 (ACE2)

An imbalance in ACE2 activation pathways may lead to an inflammatory response in people with diabetes [8]. ACE2 has a number of physiological roles including the receptor for SARS-CoV and SARS-CoV-2 and thus represents the main entry point for the coronavirus into cells [9]. Some genetic differences in ACE have been reported as being associated with susceptibility to vascular disease in the Asian population [10].

Of interest is that dietary patterns can influence ACE levels with circulating levels in the blood being very sensitive to food intake. ACE inhibitors can be found in food proteins of various origins. Peptides possessing ACE inhibitory activity have been found in milk, eggs, fish, meat, and plants with milk and dairy proteins having significant inhibitory activity [11]. Fermentation, using different live microorganism cultures, increases ACE inhibitory properties in a number of foods [12]. In contrast, a high saturated fat diet may increase ACE levels in some individuals [13].

So Could Diet Be Partly Responsible for Differences in COVID-19 Death Rates between and within Countries?

Certainly people in the southern region of Germany, where the mortality rates are higher, traditionally have a higher fat diet compared to people in the northern states. Fermented food products with potent anti-ACE activity are consumed in a number of European countries with reduced COVID severity. Fermented milk is frequently consumed in Bulgaria and fermented cabbage in Romania. People in Turkey consume both fermented milk and cabbage products. All three countries have relatively low mortality rates from COVID [1].

The easy availability to a wide range of processed foods of poor nutritional quality in a number of the countries and regions with a higher level of mortality may be a contributory factor for a number of reasons. Not only does this level of availability increase the risk of obesity but also highly processed, fibre-depleted foods have a negative effect on the gut microbiome. In contrast, probiotics can have a beneficial effect on maintaining a healthy intestinal or lung microbiota. A number of studies have used probiotic supplementation to prevent or treat respiratory tract infections, and the potential role of both probiotics and prebiotics for the prevention and treatment of COVID-19 is being investigated [14]. Of course, fermented food products are often sources of a range of these potentially beneficial live microorganisms.

In conclusion, dietitians are well placed to consider whether dietary differences both between and within countries may contribute to the severity of the COVID-19 virus, and from a public health perspective, key nutritional messages may be of importance going forward both in the treatment and the management of the disease.

Disclosure Statement

I hereby declare that there are no conflicts of interest with regard to this commentary.

References

- 1 Bousquet J, Anto JM, Laccarino G, et al.: Is diet partly responsible for differences in COVID-19 death rates between and within countries? *Clin Transl Allergy*. 2020;10:16.
- 2 Simonnet A, Chetboun M, Poissy J, et al.: High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity*. 2020;28:1195–1199.
- 3 Lewis PJ: Rapid response to Sixty seconds on vitamin D. *BMJ*. 2020;371:bmj.m3872.
- 4 Gianchandani R, Esfaniari NH, Ang L, et al.: Managing hyperglycemia in the COVID-19 inflammatory storm. *Diabetes*. 2020;69 (10):2048–2053.
- 5 Lin SC, Ho CT, Chuo WH, et al.: Effective inhibition of MERS-CoV infection by resveratrol. *BMC Infect Dis*. 2017;17(1):144.
- 6 Saedisomeolia A, Wood LG, Garg ML, et al.: Anti-inflammatory effects of long-chain n-3 PUFA in rhinovirus-infected cultured airway epithelial cells. *Br. J. Nutr.* 2009;101:533–540.
- 7 Radzikowska U, Rinaldi AO, Celebi Sozener Z, et al.: The influence of dietary fatty acids on immune responses. *Nutrients*. 2019;11(12):2990.
- 8 Cuschieri S, Grech S: COVID-19 and diabetes: The why, the what and the how. *J Diabetes Complications*. 2020;34(9):107637.
- 9 Gheblawi M, Wang K, Viveiros A, et al.: Angiotensin converting enzyme 2:SARS-CoV-2 receptor and regulator of the renin-angiotensin system. *Circ Res*. 2020;126:1456–1474.
- 10 Han C, Han XK, Liu FC, et al.: Ethnic differences in the association between angiotensin-converting enzyme gene insertion/deletion polymorphism and peripheral vascular disease: a meta-analysis. *Chronic Dis Transl Med*. 2017;3(4):230–241.
- 11 Iwaniak A, Minkiewicz P, Darewicz M: Food-originating ACE inhibitors, including antihypertensive peptides, as preventive food components in blood pressure reduction. *Comprehens Rev Food Sc Food safety*. 2014;13(2):111–134.
- 12 Nahariah N, Legowo AM, Abustam E, et al.: Angiotensin I-converting enzyme inhibitor activity on egg albumen fermentation. *Asian-Australas J Anim Sci*. 2015;28(6):855–861.
- 13 Schuler R, Osterhoff MA, Frahnw T, et al.: High-saturated-fat diet increases circulating angiotensin-converting enzyme, which is enhanced by the rs4343 polymorphism defining persons at risk of nutrient-dependent increases of blood pressure. *J Am Heart Assoc*. 2017;6(1):e004465.
- 14 Olaimat AN, Aolymat I, Al-Holy M, et al.: The potential application of probiotics and prebiotics for the prevention and treatment of COVID-19. *npj Sci Food*. 2020;4:17.

Correspondence: Dr Amanda Avery, PhD RD, Division of Food, Nutrition & Dietetics, University of Nottingham, Sutton Bonington Campus, Leicestershire, LE12 5RD, UK, amanda.avery@nottingham.ac.uk