Introduction

K. Satake S. Matsuno T. Takeuchi

The 100th anniversary of the founding of the Japanese Society of Gastroenterology was celebrated this October, meaning that pancreatology in Japan also has a history of 100 years.

In this landmark year, we were pleased to be able to host the 8th Meeting of the International Association of Pancreatology. An international group of nearly 500 research-oriented pancreatologists assembled in Tokyo to update the current status of topics related to basic and clinical pancreatology and to discuss future prospects. Four experts presented special lectures. The keynote lecture, entitled '1998, Perspectives in Pancreatic Cancer Research', was presented by Sugimura (National Cancer Center, Japan), and provided us with very important information on the future perspectives of pancreatic cancer. There were two state-of-the-art lectures. The first, entitled 'Ulcerogenic pancreatic tumor syndrome – 1998', was presented by Chey (University of Rochester), and provided us with information on a new pancreatic tumor syndrome. Beger gave the second: 'International documentation system for pancreatic cancer: A chance to overcome international barriers?' This lecture also provided us with a very important viewpoint for establishing an 'International documentation system for pancreatic cancer' to understand the current status of pancreatic cancer in each country from the same standpoint, especially the surgical standpoint. Pour (UNMC/Eppler Cancer Center) presented 'The link between exocrine pancreatic cancer and the endocrine pancreas' as an IAP Founder's Lecture. The symposia were presented by 47 experts in 8 different sessions on acute and chronic pancreatitis, pancreatic cancer, diagnostic procedures, treatment of these dis-

eases, and neurohormonal regulation of pancreatic secretion.

77 papers were presented in 16 oral sessions and 222 papers were presented in two lunchtime poster sessions, which allowed young investigators to present their findings. Fifty of 150 young investigators received the young investigator's award for their excellent presentations.

The assessments and treatment of severe acute pancreatitis are still unsolved matters in clinical situations. Yokoi at Mie University School of Medicine presented an own scoring system for assessing the severity of acute pancreatitis by the Japanese Criteria of Acute Pancreatitis and APACHE II, and established the treatment schedule for severe acute pancreatitis according to the score. Yokoi concluded that gallstone pancreatitis should first be treated by biliary drainage, and that non-gallstone necrotizing pancreatitis should be initially treated by peritoneal lavage, intra-arterial infusion of a protease inhibitor (nafamostat mesilate), and antibiotics. Bassi (University of Verona) described his experience with antiproteolytic therapy for acute pancreatitis, and concluded that antiproteases can still play a role when given prophylactically or when used in the very early phases of the disease. He also stated that a 'multiple-drugs approach' (including for example, appropriate antibiotics) seems to represent the most up-to-date and rational treatment of acute pancreatitis. Takeda (Tohoku University School of Medicine) presented a paper entitled 'The role of early continuous regional arterial infusion of protease inhibitor and antibiotics in nonsurgical management of acute necrotizing pancreatitis.' He demonstrated that, when continuous regional arterial infusion of a protease inhibitor (nafamos-

tat mesilate) and antibiotic (imipenem) were employed in patients with acute necrotizing pancreatitis within 72 h after the onset, most of the patients responded to this treatment with reduced mortality rates and incidence of infected pancreatic necrosis, and he concluded that early intervention by continuous regional arterial infusion of a protease inhibitor and antibiotic may become a very useful method for nonsurgical management for acute necrotizing pancreatitis. Bradley (State University of New York at Buffalo) reviewed 'Operative vs. non-operative therapy in necrotizing pancreatitis' from the standpoint of three important clinical corollaries: (1) the majority of acute pancreatitis patients can be treated nonoperatively; (2) early identification of patients likely to suffer complications is valuable in terms of prognosis and therapy, and (3) surgical approaches are reserved for patients developing complications. This session included a variety of important information for assessing and treating severe acute pancreatitis. Schoenberg (Rotkreuz-Krankenhaus, Munich) presented a paper entitled 'New approaches in surgical management of severe acute pancreatitis (SAP)'. He stated that, to date, the only available treatment for SAP is intensive care medicine and all its consequences, antibiotic prophylaxis, and treatment to avoid or treat infected necrosis. In patients who do not benefit from intensive care medicine or even deteriorate in their organ functions, surgical debridement of necrosis and further supportive therapy should be considered. In sterile necrosis, a delay of surgical intervention seems advisable. Finally, he concluded that the experienced surgeon will always treat the severely ill patient to his/her individual needs.

Concepts on the pathophysiology of acute pancreatitis have changed as a result of the discovery of cytokines and other mediators of inflammation. Saluja (Harvard Medical School) reviewed some of his findings regarding the steps involved in the initiation of pancreatitis and factors which determine its severity. He identified several factors that play an important pro- and anti-inflammatory role in pancreatitis. Future studies will identify many other such mediators, and drug or other interventions designed to favor anti-inflammatory or counteract pro-inflammatory events will prove useful in treating clinical pancreatitis. Yotsuya (Ishihara Sangyo Kaisha) developed a new N-(2-sulfonylamino-5-trifluoromethyl-3-pyridyl) carboxamide derivative IS 741. This drug has been found to have an anti-inflammatory effect mediated by actions that inhibit cell adhesion *in vitro* and neutrophil infiltration *in vivo*. He used this drug to treat experimental acute pancreatitis in various models and found that it improved the survival

rate of severe acute pancreatitis in various species. He also suggested that the drug inhibits pancreatic tissue destruction by neutrophils through its inhibitory action on neutrophil infiltrations of the pancreas, and that it may have the same effect on other important organs during severe acute pancreatitis. Yamauchi (Tohoku University School of Medicine) investigated the mechanism underlying leukocyte accumulation in the pancreas and attempted to elucidate the therapeutic efficacy of IS 741 on leukocyte-endothelial cell interaction in a rat necrotizing pancreatitis model. His group demonstrated that modulation of leukocyte kinetics in the early phase of the disease could be an attractive strategy for preventing aggravation of acute pancreatitis, and IS 741 may be a candidate drug for reducing expression of CD 11b/18, and subsequently inhibit leukocyte endothelial cell interaction. Isaji (Mie University School of Medicine) presented their paper entitled 'Effect of IS 741 on acute necrotizing pancreatitis in dogs: Significance of its inhibitory effect on cytosolic phospholipase A₂.' They found that the survival rate of IS 741-treated animals was significantly higher than in the control, and that cytosolic phospholipase A₂ activity in the pancreas, lung, and renal cortex was much lower in the IS 741-treated animals than in the control. They concluded that IS 741 exerts a therapeutic effect on acute pancreatitis by preventing histological damage in the pancreas, lung, and kidney, and that the inhibitory effect of this drug on cytosolic phospholipase A₂ activity may play an important role attenuating acute necrotizing pancreatitis. Yamaguchi (Kumamoto University Medical School) demonstrated a reduced production of cytokine-induced neutrophil chemoattractant by bronchoalveolar macrophages and effective prevention of pancreatitis-associated lung injury by intravenous infusion of IS 741 following the septic challenge in rats. This new drug may open some new approaches to the treatment of acute pancreatitis, although more careful investigations of this drug are needed in regard to its clinical aspects. Norman (University of South Florida) reviewed 'New approaches to acute pancreatitis: The role of inflammatory mediators' in regard to four different aspects: (1) cytokinin activation during clinical pancreatitis; (2) inflammatory mediators in experimental pancreatitis; (3) antagonism of inflammatory mediators during acute pancreatitis, and (4) the clinical possibility of anticytokine therapy for acute pancreatitis. He concluded that the majority of the morbidity and mortality of acute pancreatitis is due to systemic manifestations secondary to inflammatory mediator overproduction. As we learn more about the production and activities of these powerful mediators, a scenario is

likely to develop that allows specific therapies for each phase of the disease to prevent progression of gland destruction and the development of systemic complications.

It is well known that CCK may have an important role in the development and progression of acute pancreatitis. In this session, Beglinger (University Hospital, Basel) reviewed 'Potential role of CCK in the development of acute pancreatitis' including: (1) the role of CCK in the development of acute pancreatitis, and (2) the effect of CCK-A receptor antagonists in models of acute pancreatitis. He also touched on the importance of cytokines during the progression of acute pancreatitis, and concluded that (1) endogenous CCK plays a modulatory, aggravating role in the development and progression of acute pancreatitis; (2) the effects of CCK are mediated by CCK-A receptor, and (3) CCK-A receptor antagonists may have therapeutic potential for treatment of acute pancreatitis. Finally, he stated that clinical trials of CCK-A receptor antagonists are necessary to document their efficacy in human disease. Satake (Osaka City University Medical School) presented the work of his groups: 'Therapeutic effects of CCK-A receptor antagonist (Loxiglumide) on experimental acute pancreatitis in various models', and reported findings that the CCK-A receptor antagonist displayed therapeutic and/or prophylactic effects on acute pancreatitis in various experimental models. Niederau (St. Josef-Hospital) reviewed the role of CCK as a contributory factor to the development and progression of acute pancreatitis and prospects of CCK receptor antagonists as treatment for acute pancreatitis based on his tremendous experience in this field. He concluded that CCK acts as a permissive or contributory factor in the development and progression of acute pancreatitis, and that CCK antagonists have a potential therapeutic benefit. He also stated that clinical studies will evaluate their therapeutic potential for acute pancreatitis patients. Funakoshi (National Kyushu National Cancer Center) presented a paper entitled 'Mechanism of CCK-A receptor antagonist on human exocrine secretion: localization of CCK-A receptor in the human duodenum'. He reported that CCK-A receptor mRNA is not expressed in the pancreas, but is expressed in the gallbladder and duodenum. This is a very important finding in regard to whether CCK-A receptor antagonists can be used to treat acute pancreatitis in humans as experimental animals. They speculated that the mechanism of CCK-A receptor antagonist in human pancreatic exocrine secretion is that specific CCK-A receptor antagonists inhibit pancreatic exocrine secretion acting via the receptor sites of terminal afferent fibers,

because CCK and duodenal stimuli may act on distinct duodenal afferent fibers to potentiate pancreatic exocrine secretion. Their findings suggested that CCK-A receptor antagonist may become a useful tool for the treatment of acute pancreatitis. The first preliminary clinical trials of a CCK-A receptor antagonist (loxiglumide) for acute pancreatitis were presented by the loxiglumide study group in Japan. They found that CCK-A receptors had efficacy on human acute pancreatitis, although we need more detailed clinical analysis in the future to determine whether loxiglumide possesses efficacy on human acute pancreatitis.

In chronic pancreatitis, Kataoka (Kyoto Prefectural University of Medicine) evaluated the diagnostic value of two noninvasive tests, BT-PABA and fecal chymotrypsin (FTC), to assess pancreatic exocrine function in chronic pancreatitis. They concluded that, although noninvasive tests are less sensitive and specific for diagnosing pancreatic exocrine dysfunction than the secretion test, a greater reliability for diagnosing chronic pancreatitis can be obtained by using BT-PABA and FCT simultaneously. Nakamura (Hirosaki University School of Medicine) presented his group's works on pancreatic exocrine insufficiency (chronic pancreatitis) and steatorrhea in Japanese patients. They found that the incidence of steatorrhea in Japanese is lower than in Europeans and Americans, and that the grade of steatorrhea is milder in Japanese because the fat intake is lower than that of Europeans and Americans, making it difficult to compare the findings regarding pancreatic exocrine insufficiency between Japan and other countries. However, pancreatic enzyme preparations are very important in controlling steatorrhea in Japanese chronic pancreatitis patients. Lankish (Municipal Clinic Lüneburg) gave a very important lecture entitled 'What to do when a patient with exocrine pancreatic insufficiency does not respond to pancreatic enzyme substitution'. He recommended: increasing enzyme dosage, checking the patient's compliance, reevaluating the diagnosis, and, as a last resort, decreasing fat intake or adding H₂-blocker or proton pump inhibitors. Neoptolemos (University of Liverpool) made a presentation on pancreatic exocrine insufficiency following pancreatic resection. How to control this problem after pancreatic resection is a very important question among surgeons. He stated that about 70% of patients with steatorrhea can be controlled with the effective doses of pancreatic enzyme supplements, but that the remaining 30% still have evidence of significant levels of steatorrhea. He concluded that complete eradication of steatorrhea is not feasible at present, even using the best currently available agents, underscoring the need for more

effective therapies. Imrie (Glasgow Royal Infirmary) lectured on 'The diagnosis of chronic pancreatitis and new aspects of pain control'. He concluded that minimal invasive thoroscopic splachnicectomy is an adequate method for controlling the pain of chronic pancreatitis.

The Japanese experience with surgery for pancreatic cancer was presented in the form of a video symposium. All speakers described their operative methods and results of their operations in this supplement. Takada (Teikyo University School of Medicine) reviewed historical developments in surgery for pancreatic cancer in Japan and discussed its future prospects based on our vast experience. Kwarada (Mie University School of Medicine) presented his results of modified standard pancreatoduodenectomy (P-D) for the treatment of pancreatic head cancer. He has performed three different operative procedures during the past 22 years: (1) standard; (2) extended, and (3) modified standard P-D. He obtained better post-operative QOL and survival rates with modified standard P-D than with the other procedures. Imamura (Kyoto University) reported his group's results with extended resection surgery and multimodality treatment. He concluded that it may be possible to control the local recurrence of pancreatic cancer by extended resection and

intraoperative irradiation, and that the early development of hepatic metastasis might be controlled by regional chemotherapy. Takahashi (Keio University School of Medicine) presented his group's treatment strategy for pancreatic head cancer and their results with pylorus-preserving pancreatoduodenectomy (PPPD), intraoperative radiotherapy (IORT), and portal catheterization with fluorouracil infusion (PC). He concluded that (1) PPPD allows almost the same extent of D2 dissection as conventional P-D, and achieves radical treatment without any problems. (2) The local recurrence of pancreatic cancer after surgery can not be controlled by IORT. (3) Liver metastasis in such patients can be suppressed by PC. Ishikawa (Osaka Medical Center for Cancer and Cardiovascular Disease) presented his group's results with extended P-D and liver perfusion chemotherapy. He concluded that extended P-D and liver perfusion chemotherapy (without radiation) improves the 5-year survival of these patients. The perspective of the investigators in this session is different, but their presentation contained very important information on the treatment of pancreatic cancer in the future, although a standard method of staging pancreatic cancer worldwide is needed.