Surveillance, Diagnosis, and Treatment Outcomes of Hepatocellular Carcinoma in Japan: 2021 Update

Masatoshi Kudo

Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, Osaka, Japan

Prof. M. Kudo

Editor *Liver Cancer*

Keywords

Hepatocellular carcinoma · Surveillance · Diagnosis · Treatment outcomes

Introduction

The Liver Cancer Study Group of Japan has recently published the results of its 21st Nationwide Follow-up Survey of Primary Liver Cancer in Japan [1]. The group's report covers a 2-year period from January 1, 2010, to December 31, 2011, and provides basic statistics on 22,134 prospectively enrolled patients from 546 institutions, and survival data on 41,956 previously enrolled patients, who were followed through this reporting period. The Liver Cancer Study Group of Japan was founded in 1965, and

this year's follow-up survey report is the 21st in a series that started with the first such survey in 1969. The participating institutions in this survey register newly enrolled patients with primary liver cancer, and enter follow-up data on previously enrolled patients in the National Clinical Database once every 2 years, based on the General Rules for the Clinical and Pathological Study of Primary Liver Cancer [2–4]. This database is used for a range of analyses performed by the Liver Cancer Study Group of Japan, and the analysis results are published in Japanese in booklet form every 2 years. When each of the previous 20 survey reports was released, a concise version was published in Japanese in *Kanzo*, the journal of the Japan Society of Hepatology and in English in an international journal at the same time [5–49]. This editorial highlights the main results in the concise English version of the report and the longevity survey and analysis results for the patients registered in the 2-year period 2010–2011 and the previously registered patients (follow-up cases) [1]. In addition, surveillance, diagnosis, and treatment outcomes for hepatocellular carcinoma (HCC) in Japan are discussed.

Patient Characteristics at the Time of Initial Detection

The Japan Society of Hepatology's clinical practice guideline recommends that HCC surveillance cover the points outlined below. Patients at super-high risk for

Fig. 1. Numbers of nodules at the time of initial detection in patients with HCC ($n = 19,536$) registered in the nationwide follow-up survey of the LCSGJ between January 1, 2010, to December 31, 2011, in 546 institutions throughout Japan. HCC, hepatocellular carcinoma; LCSGJ, Liver Cancer Study Group of Japan. Modified from Ref. [1] and [50] with permission.

Fig. 2. Maximum tumor size at the time of initial detection in patients with HCC ($n = 19,537$) registered in the nationwide follow-up survey of the LCSGJ between January 1, 2010 to December 31, 2011 in 546 institutions throughout Japan. HCC, hepatocellular carcinoma; LCSGJ, Liver Cancer Study Group of Japan. Modified from Ref. [1] and [50] with permission.

HCC (those with hepatitis B or C cirrhosis or nonviral cirrhosis) are recommended to undergo ultrasonography and measurements of 3 tumor markers – namely, α -fetoprotein (AFP), protein induced by vitamin K absence or antagonist II (PIVKA-II), and *Leish culinaris* agglutinin-reactive AFP fraction (AFP-L3) – once every 3–4 months, with optional dynamic CT or Gd-EOB-DTPA-enhanced MRI once every 6–12 months.

High-risk patients (defined as those chronic hepatitis B or C are recommended to undergo surveillance by ultrasonography and measurements of the 3 tumor markers (AFP, PIVKA-II, and AFP-L3) once every 6 months. All imaging and tumor marker examinations are covered by national health insurance for super-high-risk and

high-risk patients, and thus can be performed by almost all relevant institutions and private practitioners in Japan, thus contributing to the early detection of liver cancer.

The 21st follow-up survey presents a number of findings for patients newly diagnosed with HCC in the 2010–2011 period [50]. Among these patients ($n = 19,536$), 63.5% had a solitary nodule when they were diagnosed with HCC (Fig. 1). Measurements of maximum tumor size (irrespective of the number of nodules) revealed that small HCCs accounted for a large proportion of the newly diagnosed cases, with maximum tumor size measuring ≤3 cm in 56.6% of patients, and ≤2 cm in 34% of patients (Fig. 2). Thus, many of the HCCs in the Japanese population are detected as small and/or single tumors. Extrahe-

Fig. 3. Presence or absence of extrahepatic spread at the time of initial detection in patient with HCC ($n = 19,887$) registered in the nationwide follow-up survey of the LC-SGJ between January 1, 2010, to December 31, 2011, in 546 institutions throughout Japan. HCC, hepatocellular carcinoma; LC-SGJ, Liver Cancer Study Group of Japan. Modified from Ref. [1] and [50] with permission.

Fig. 4. Presence or absence of portal vein invasion at the time of initial detection in patients with HCC ($n = 19,167$) registered in the nationwide follow-up survey of the LC-SGJ between January 1, 2010, to December 31, 2011, in 546 institutions throughout Japan. Portal vein invasion was observed in 13.2% of the patients. HCC, hepatocellular carcinoma; LC-SGJ, Liver Cancer Study Group of Japan; Vp1, invasion (or tumor thrombus) distal to the second-order branches of the portal vein, but no invasion of the second-order branches; Vp2, invasion (or tumor thrombus) of the second-order branches of the portal vein; VP3, invasion (or tumor thrombus) of the first-order branches of the portal vein; Vp4, invasion (or tumor thrombus) of the main trunk of the portal vein and/or the portal vein branch contralateral to the primarily involved lobe. Modified from Ref. [1] and [50] with permission.

patic spread was found in 802 of 19,887 patients (4.0%) examined at the time of initial detection, showing that extrahepatic spread is considerably less common in Japan than in other countries (Fig. 3). Portal vein invasion was also less common in Japan than in other countries, but its incidence was higher than expected, occurring in 2,523 of 19,167 patients (13.2%) examined at the time of initial detection. This higher-than-expected incidence could be explained by recent advances in CT and MRI [51, 52] and abdominal contrast-enhanced ultrasound [53], which have made minor vascular invasion graded as Vp1 or Vp2 clinically discernable from the time of initial detection for small HCCs.

At initial detection, 22.8% of HCC patients had tumors measuring ≥ 5 cm, and 12.8% had multiple tumors with ≥ 4 nodules. Considering these findings, we can say that some cases had advanced HCC at the time of initial detection, and major vascular invasion graded as Vp3 or Vp4 occurred to some extent (Fig. 4). Hepatic vein invasion was found in 1,179 of 18,700 patients (6.2%) examined at the time of initial detection (Fig. 5). This result also demonstrates that, to a certain extent, HCC is detected in patients who do not regularly undergo the periodical surveillance. Up to now, the surveillance system has been designed to provide full coverage to patients with hepatitis B virus (HBV) or hepatitis C virus (HCV)-related

Fig. 5. Presence or absence of hepatic vein invasion at the time of initial detection in patients with HCC ($n = 18,770$) registered in the nationwide follow-up survey of the LCSGJ between January 1, 2010, to December 31, 2011, in 546 institutions throughout Japan. HCC, hepatocellular carcinoma; LCSGJ, Liver Cancer Study Group of Japan; Vv1, invasion (or tumor thrombus) of the peripheral branches of the hepatic vein; Vv2, invasion (or tumor thrombus) of the right, middle, or left hepatic vein, the inferior right hepatic vein, or the short hepatic vein; Vv3, invasion (or tumor thrombus) of the main hepatic vein or the inferior vena cava. Modified from Ref. [1] and [50] with permission.

Fig. 6. Incidence rate of HCC according to etiology. HCC, hepatocellular carcinoma; HCV, hepatitis C virus. Modified from Ref. [1] and [50] with permission.

Fig. 7. AFP level (ng/mL) at the time of initial detection in patients with HCC ($n = 19,466$) registered in the nationwide follow-up survey of the LCSGJ between January 1, 2010, to December 31, 2011, in 546 institutions throughout Japan. HCC, hepatocellular carcinoma; LCSGJ, Liver Cancer Study Group of Japan; AFP, serum α -fetoprotein. Modified from Ref. [1] and [50] with permission.

Fig. 8. AFP-L3 (*Lens culinaris* agglutinin-reactive AFP fraction; %) at the time of initial detection in patients with HCC ($n = 9,504$) registered in the nationwide follow-up survey of the LCSGJ between January 1, 2010 to December 31, 2011 in 546 institutions throughout Japan. HCC, hepatocellular carcinoma; LCSGJ, Liver Cancer Study Group of Japan; AFP, serum α -fetoprotein. Modified from Ref. [1] and [50] with permission.

Fig. 9. PIVKA-II level at the time of initial detection in patients with HCC ($n = 18,824$) registered in the nationwide follow-up survey of the LCSGJ between January 1, 2010, to December 31, 2011, in 546 institutions throughout Japan. HCC, hepatocellular carcinoma; LCSGJ, Liver Cancer Study Group of Japan; PIVKA-II, protein induced by vitamin K absence or antagonist II. Modified from Ref. [1] and [50] with permission.

chronic liver disease; however, non-HBV/non-HCV-related HCC, including HCC related to nonalcoholic steatohepatitis, has become relatively common in Japan [54]. This phenomenon may explain why cases of advanced disease have recently continued to be detected at the same rates as previously in Japan. Indeed, non-HBV/non-HCV-related HCCs account for a high proportion of patients (31.5%) in the report of the 21st Nationwide Follow-up Survey of Primary Liver Cancer in Japan (Fig. 6). We consider it necessary to further promote awareness-raising among the general public and private practitioners in Japan.

Serum AFP level was below the normal range (≤ 15 ng/mL) in 46.1% of patients and ≤ 200 ng/mL in 77.6% of patients; overall, AFP levels were low in many cases at diagnosis (Fig. 7). Abnormal levels of the AFP-L3 fraction ($\geq 10\%$) were noted in 3,273 of 9,504 patients (34.4%) (Fig. 8), and screening with this tumor marker had high specificity but not sensitivity [55–59]. The 3 tumor markers AFP, PIVKA-II, and AFP-L3 are known not to be correlated; therefore, measuring all 3 markers at the same time enables complementarily greater detection of HCCs [58]. The proportion of patients with normal levels of PIVKA-II, which is also called DCP (des-gamma-carboxy prothrombin), was 38.0% (Fig. 9).

Fig. 10. Initial treatment modality selected after diagnosis of HCC over time. Curative treatments (resection 41.3%, ablation 24.5%) were performed in 65.8% of HCC patients registered in the nationwide follow-up survey of the LCSGJ between January 1, 2010, to December 31, 2011, in 546 institutions throughout Japan. The rate of resection is gradually increasing over time. Systemic therapy

was used in only 2.1% of HCC patients. HCC, hepatocellular carcinoma; LCSGJ, Liver Cancer Study Group of Japan; HAIC, hepatic arterial infusion chemotherapy; TACE, transcatheter arterial chemoembolization. Modified from Ref. [1] and [50] with permission.

The above findings show that, with the establishment of the Japanese nationwide surveillance program [1, 60], detected HCCs tended to be solitary, small tumors and the levels of tumor markers tended to be low [61].

Treatment options selected by patients initially diagnosed with HCC in 2010–2011 were surgical resection in 41.3% of patients, ablation therapy in 24.5%, and transcatheter arterial chemoembolization (TACE) in 27.2%. Only 2.1% of patients opted for systemic therapy as initial treatment (Fig. 10). The proportion of patients opting for surgical resection has gradually increased in recent years,

and probably this may be related to the increase in non-HBV/non-HCV-related HCCs.

Overall Survival by Treatment Modality and Other Factors

Overall survival (OS) is presented by Child-Pugh grade for 27,903 patients who underwent liver resection between 2002 and 2013 in Figure 11. Of these patients, 25,492 had preserved liver function (Child-Pugh grade

Fig. 11. OS in HCC patients (registered in the follow-up survey from 2002 to 2013) who were treated with resection according to the Child-Pugh grade ($n = 27,903$). Median OS for Child-Pugh A patients treated with resection was 95.0 months, and 5- and 10-year survival rates were 64.9 and 41.3%, respectively. HCC, hepatocellular carcinoma; LSCGJ, Liver Cancer Study Group of Japan; OS, overall survival. Modified from Ref. [1] and [50] with permission.

A), Their median OS was 95.0 months (approximately 7.9 years), and their 5- and 10-years survival rates were 65 and 41%, respectively (Fig. 11). Portal vein invasion was associated with survival, and patients with portal vein invasion graded as Vp4 who underwent surgical resection had a median OS of 11.1 months (Fig. 12). Serum AFP level was correlated with OS in patients with resected HCC (Fig. 13). This is highly consistent with the fact that AFP is a strong prognostic factor [62].

OS is presented by Child-Pugh grade for patients who underwent local ablation therapy in Figure 14. Median OS for patients with Child-Pugh grade A liver function was 79.9 months (approximately 6.7 years), and their 5- and 10-years survival rates were 64.1 and 28.3%, respectively (Fig. 14). Numerically, local ablation therapy showed OS inferior to that of surgical resection: 6.7 versus 7.9 years. However, survival rates did not differ between radiofrequency ablation and surgery in a multicenter, randomized control study (the SURF Trial, recently conducted in Japan) of 300 HCC patients with ≤ 3 tumors measuring ≤ 3 cm [63]. Accordingly, the difference in survival between surgical resection and ablation was attributable to patients with a more favorable condition, among those with Child-Pugh grade A liver function, opting for surgical resection.

Fig. 12. OS in HCC patients (registered in the follow-up survey from 2002 to 2013) who treated with resection according to extent of portal vein invasion. HCC, hepatocellular carcinoma; LSCGJ, Liver Cancer Study Group of Japan; OS, overall survival; Vp0, absence of portal vein invasion; Vp1, invasion (or tumor thrombus) distal to the second-order branches of the portal vein, but no invasion of the second-order branches; Vp2, invasion (or tumor thrombus) of second-order branches of the portal vein; Vp3, invasion (or tumor thrombus) of the first-order branches of the portal vein; Vp4, invasion (or tumor thrombus) of the main trunk of the portal vein and/or the portal vein branch contralateral to the primarily involved lobe. Modified from Ref. [1] and [50] with permission.

OS is presented by Child-Pugh grade for patients who underwent TACE in Figure 15. Median OS for patients with Child-Pugh grade A liver function was 45.3 months, and the 5- and 10-years survival rates for these patients were 38.3 and 14.7%, respectively (Fig. 15). This 45.3-month OS represents a markedly prolonged survival period. Given that the longest OS shown in 6 randomized control trials of TACE monotherapy (post TACE [64], TACE-2 [65], SPACE [66], ORIENTAL [67], BRISQ-TA [68], and TACTICS [69]) was 30 months in the TACTICS study [70], we consider that many of the patients in Japan may have corresponded to Stage A in the Barcelona Clinic Liver Cancer system (a single tumor of any size or ≤ 3 tumors measuring ≤ 3 cm).

OS is presented by the Child-Pugh grade for patients who underwent reservoir-based continuous hepatic arterial infusion chemotherapy (HAIC) in Figure 16. Median OS for patients with Child-Pugh grade A liver function was 12.9 months, and the 5- and 10-years survival

Fig. 13. OS in HCC patients (registered in the follow-up survey from 2002 to 2013) who were treated with resection according to serum AFP level (ng/mL). HCC, hepatocellular carcinoma; OS, overall survival; AFP, serum α -fetoprotein. Modified from Ref. [1] and [50] with permission.

Fig. 15. OS in HCC patients (registered in the follow-up survey from 2002 to 2013) who were treated with TACE according to Child-Pugh grade (n = 18,750). Median OS for Child-Pugh A patients treated with TACE was 45.3 months, and 5- and 10-year survival rates were 38.3 and 14.7%, respectively. HCC, hepatocellular carcinoma; OS, overall survival; TACE, transcatheter arterial chemoembolization. Modified from Ref. [1] and [50] with permission.

Fig. 14. OS in HCC patients (registered in the follow-up survey from 2002 to 2013) treated with local ablation therapy according to Child-Pugh grade (n = 22,776). Median OS for Child-Pugh A patients (registered in the follow-up survey from 2002 to 2013) who were treated with local ablation therapy was 79.9 months, and 5- and 10-year survival rates were 64.1 and 28.3%, respectively. HCC, hepatocellular carcinoma; OS, overall survival. Modified from Ref. [1] and [50] with permission.

Fig. 16. OS in HCC patients (registered in the follow-up survey from 2002 to 2013) treated with continuous HAIC using a reservoir (implanted port system) according to Child-Pugh grade (n = 1,429). Median OS for Child-Pugh A patients treated with HAIC was 12.9 months, and 5- and 10-year survival rates were 14.0 and 5.0%, respectively. HCC, hepatocellular carcinoma; OS, overall survival; HAIC, hepatic arterial infusion chemotherapy. Modified from Ref. [1] and [50] with permission.

Fig. 17. Improvement of 5-year survival rates in all patients with BCLC Stage 0, A, B, C, and D registered in the follow-up survey during the 7 periods listed below. The 5-year survival rates for HCC patients registered in the nationwide follow-up survey in the periods 1978–1980, 1981–1985, 1986–1990, 1991–1995, 1996–

2000, 2001–2005, and 2006–2009 were 5, 14, 25, 32, 39, 43, and 50%, respectively. HCC, hepatocellular carcinoma; HAIC, hepatic arterial infusion chemotherapy; PIVKA-II, protein induced by vitamin K absence or antagonist II; BCLC, Barcelona Clinic Liver Cancer. Modified from Ref. [1] and [50] with permission.

rates for these patients were 14.0 and 5.0%, respectively (Fig. 16). Reservoir-based arterial infusion achieved similar outcomes to molecularly targeted therapy with sorafenib, which is restricted to patients with Child-Pugh grade A liver function. The SILIUS study [71] compared OS between sorafenib plus arterial infusion chemotherapy and sorafenib alone and demonstrated that addition of HAIC to sorafenib was not effective; however, sorafenib plus arterial infusion chemotherapy demonstrated clear superiority among patients with tumors showing Vp4 portal vein tumor thrombus, with a median OS of 11.4 versus 6.5 months for sorafenib alone

(hazard ratio: 0.493; 95% confidence interval: 0.240–1.014) [71]. According to this 21st nationwide follow-up survey in Japan, OS for patients with Vp4 portal vein invasion receiving HAIC monotherapy was 5.7 months [1]. A large proportion of these patients were classified as Child-Pugh grade B rather than Child-Pugh grade A, and this raises the possibility that HAIC is at least equivalent or slightly superior to sorafenib. Nationwide data on arterial infusion chemotherapy versus sorafenib were evaluated retrospectively by propensity score matching, and the results showed that HAIC was significantly superior to sorafenib in patients with Vp3/Vp4 vascular

Fig. 18. Improvement of OS in all patients with BCLC Stage 0, A, B, C, and D registered in the follow-up survey during the 7 periods listed below. Median OS for patients with HCC registered in the nationwide follow-up survey in the periods 1978–1980, 1981–1985, 1986–1990, 1991–1995, 1996–2000, 2001–2005, and 2006–

2009 was 4, 16, 26, 36, 44, 50, and 60 months, respectively. HCC, hepatocellular carcinoma; OS, overall survival; BCLC, Barcelona Clinic Liver Cancer. Modified from Ref. [1] and [50] with permission.

invasion, with a median OS of 10.6 months versus 9.1 for sorafenib [72].

HAIC with a FOLFOX regimen plus sorafenib was reported to be significantly superior to sorafenib alone in cases of advanced HCC with portal vein invasion in a study in China [73]. Accordingly, among the treatment options, HAIC will probably be used as the first-line therapy for patients with major vascular invasion (Vp3 or Vp4) in Asia [74, 75]. Meanwhile, a groundbreaking therapy, atezolizumab plus bevacizumab combination immunotherapy, was approved in 2020 [76]. As a result, with the accumulation of cases given this therapy, consideration is now being given to atezolizumab plus bevacizum-

ab combination immunotherapy becoming the treatment of choice for patients with major vascular invasion (Vp3 or Vp4). This is because of the high efficacy shown in a subset of patients in IMbrave150 trial – the so-called REFLECT out patients set (101 cases, comprising 19% of the original study population) – who were excluded from the REFLECT study in patients with Vp4 vascular invasion, tumors occupying >50% and invasion to the bile duct [77]. “REFLECT out” patents with Vp4 vascular invasion (48 cases, 14%) had OS of 7.6 months, progression-free survival of 5.4 months, and an objective response rate of 25% [78].

Improvement of Treatment Outcomes in Patients with HCC

Between 1978 and 1980, the 5-year survival rate for the 2,323 HCC patients registered was 5%; however, this rate has gradually improved for subsequently registered patients. The 5-years survival rate for the 39,423 patients registered between 2006 and 2009 was 50%. This includes survival for all patients at all Barcelona Clinic Liver Cancer stages, from the early stage of 0 to stage D (Fig. 17). Median OS for patients registered between 1978 and 1980 was 4 months but has gradually improved since then, with median OS of 60 months for patients registered between 2006 and 2009 (Fig. 18). The major reasons for this improvement were as follows: AFP tests and ultrasound were first introduced into the Japanese nationwide surveillance program in the 1980s, surgical resection was established and TACE started being adopted as a treatment option across Japan around 1985, and percutaneous ethanol injection therapy was developed in Japan in the 1990s. All these factors are considered to have contributed to improved outcomes for HCC. Other contributing factors have been proposed: the growing use of helical CT and abdominal MRI across Japan since the 1990s, and the approval of interferon therapy for HCV and PIVKA-II testing approved by insurance in 1989. These developments have further enabled early detection of HCC.

Furthermore, HAIC has been implemented in many hospitals in Japan since around 1995, improving survival for HCC patients with vascular invasion. Insurance coverage was extended to AFP-L3 testing in 1996, and methods were developed for analysis of the 3 tumor markers AFP, AFP-L3, and DCP (PIVKA-II). These developments have further advanced the potential for early HCC detection.

The years just after 2000 saw rapidly increasing use of radiofrequency ablation as well as an increase in the use of multi-detector row CT scanners, which were also factors in improving early HCC detection. Since that time, sorafenib was approved in Japan in 2009 [79]. Accordingly, some of the patients registered in the 2001–2005 or 2006–2009 period received sorafenib or other molecular targeted agents after undergoing surgical resection or locoregional therapy, and this is also considered to be a factor in the improved survival outcomes. With the approval of regorafenib in 2017 [80], lenvatinib in 2018 [77], ramucirumab in 2019 [81], and atezolizumab plus bevacizumab combination immunotherapy [76] and cabozantinib 2020 [82], further improvements in survival for HCC patients are expected [83, 84].

The survival of intermediate-stage HCC patients is also being markedly improved through the evolution of therapeutic strategies, with a gradual change to a strategy of selective TACE after initial introduction of systemic therapy [85–87]. Clinical trials now in progress are evaluating combination immunotherapies (anti-PD-1/PD-L1 antibody plus anti-CTLA4 antibody [88] and anti-PD-1/anti-PD-L1 antibody plus anti-VEGF antibody [89]) applied after surgical resection or ablation as adjuvant therapy, and in combination with TACE [90]. With the application of these therapies in actual clinical settings, we can anticipate further improvements in survival for Japanese patients with HCC who previously would be out of treatment options.

Statement of Ethics

Not applicable.

Conflict of Interest Statement

Lecture: Eisai, Bayer, MSD, BMS, EA Pharma, Eli Lilly, and Chugai; grants: Eisai, Takeda, Otsuka, Taiho, EA Pharma, Gilead Sciences, Abbvie, Sumitomo Dainippon Pharma, Chugai, and Ono Pharma; advisory consulting: Eisai, Ono, MSD, BMS, and Roche. Masatoshi Kudo is the Editor-in-Chief of Liver Cancer.

Funding Sources

The authors did not receive any funding.

Author Contributions

M. Kudo conceived, wrote, and approved the final manuscript.

References

- 1 Kudo M, Izumi N, Kokudo N, Sakamoto M, Shiina S, Takayama T, et al. Report of the 21st nationwide follow-up survey of primary liver cancer in Japan (2010–2011). *Hepatol Res.* 2021;51:335–405.
- 2 Liver Cancer Study Group of Japan. *The general rules for the clinical and pathological study of primary liver cancer*. 1st ed. Tokyo: Kanehara & Co., Ltd.; 1997.
- 3 Liver Cancer Study Group of Japan. *General rules for the clinical and pathological study of primary liver cancer*. 2nd ed. Tokyo: Kanehara & Co., Ltd.; 2003.

- 4 Liver Cancer Study Group of Japan. [General rules for the clinical and pathological study of primary liver cancer](#). 3rd ed. Tokyo: Kanehara & Co., Ltd.; 2010.
- 5 Okuda K. Primary liver cancers in Japan. [Cancer](#). 1980;45:2663–9.
- 6 Liver Cancer Study Group of Japan. Primary liver cancer in Japan. The Liver Cancer Study Group of Japan. [Cancer](#). 1984;54:1747–55.
- 7 Liver Cancer Study Group of Japan. Primary liver cancer in Japan. Sixth report. [Cancer](#). 1987;60:1400–11.
- 8 Liver Cancer Study Group of Japan. Primary liver cancer in Japan. Clinicopathologic features and results of surgical treatment. [Ann Surg](#). 1990;211:277–87.
- 9 Tobe T, Kameda H, Okudaira M, Ohto M, Endo Y, Mito M, et al., editors. [Primary liver cancer in Japan](#). Berlin, Heidelberg, New York, London, Paris, Hong Kong Barcelona: Springer-Verlag Tokyo; 1992.
- 10 Liver Cancer Study Group of Japan. Predictive factors for long term prognosis after partial hepatectomy for patients with hepatocellular carcinoma in Japan. The Liver Cancer Study Group of Japan. [Cancer](#). 1994;74:2772–80.
- 11 Arii S, Yamaoka Y, Futagawa S, Inoue K, Kobayashi K, Kojiro M, et al. Results of surgical and nonsurgical treatment for small-sized hepatocellular carcinomas: a retrospective and nationwide survey in Japan. Liver Cancer Study Group of Japan. [Hepatology](#). 2000;32:1224–9.
- 12 Ikai I, Itai Y, Okita K, Omata M, Kojiro M, Kobayashi K, et al. Report of the 15th follow-up survey of primary liver cancer. [Hepatol Res](#). 2004;28:21–9.
- 13 Ikai I, Arii S, Kojiro M, Ichida T, Makuuchi M, Matsuyama Y, et al. Reevaluation of prognostic factors for survival after liver resection in patients with hepatocellular carcinoma in a Japanese nationwide survey. [Cancer](#). 2004;101:796–802.
- 14 Ikai I, Arii S, Ichida T, Okita K, Omata M, Kojiro M, et al. Report of the 16th follow-up survey of primary liver cancer. [Hepatol Res](#). 2005;32:163–72.
- 15 Ikai I, Arii S, Okazaki M, Okita K, Omata M, Kojiro M, et al. Report of the 17th nationwide follow-up survey of primary liver cancer in Japan. [Hepatol Res](#). 2007;37:676–91.
- 16 Takayasu K, Arii S, Ikai I, Omata M, Okita K, Ichida T, et al. Prospective cohort study of transarterial chemoembolization for unresectable hepatocellular carcinoma in 8,510 patients. [Gastroenterology](#). 2006;131:461–9.
- 17 Ikai I, Takayasu K, Omata M, Okita K, Nakanuma Y, Matsuyama Y, et al. A modified Japan integrated stage score for prognostic assessment in patients with hepatocellular carcinoma. [J Gastroenterol](#). 2006;41:884–92.
- 18 Minagawa M, Ikai I, Matsuyama Y, Yamaoka Y, Makuuchi M. Staging of hepatocellular carcinoma: assessment of the Japanese TNM and AJCC/UICC TNM systems in a cohort of 13,772 patients in Japan. [Ann Surg](#). 2007;245:909–22.
- 19 Eguchi S, Kanematsu T, Aii S, Okazaki M, Okita K, Omata M, et al. Comparison of the outcomes between an anatomical subsegmentectomy and a non-anatomical minor hepatectomy for single hepatocellular carcinomas based on a Japanese nationwide survey. [Surgery](#). 2008;143:469–75.
- 20 Hasegawa K, Makuuchi M, Takayama T, Kokudo N, Aii S, Okazaki M, et al. Surgical resection vs. percutaneous ablation for hepatocellular carcinoma: a preliminary report of the Japanese nationwide survey. [J Hepatol](#). 2008;49:589–94.
- 21 Takayasu K, Aii S, Ikai I, Kudo M, Matsuyama Y, Kojiro M, et al. Overall survival after transarterial lipiodol infusion chemotherapy with or without embolization for unresectable hepatocellular carcinoma: propensity score analysis. [AJR Am J Roentgenol](#). 2010;194:830–7.
- 22 Ikai I, Kudo M, Aii S, Omata M, Kojiro M, Sakamoto M, et al. Report of the 18th follow-up survey of primary liver cancer in Japan. [Hepatol Res](#). 2010;40(11):1043–59.
- 23 Eguchi S, Kanematsu T, Aii S, Omata M, Kudo M, Sakamoto M, et al. Recurrence-free survival more than 10 years after liver resection for hepatocellular carcinoma. [Br J Surg](#). 2011;98:552–7.
- 24 Higashi T, Hasegawa K, Kokudo N, Makuuchi M, Izumi N, Ichida T, et al. Demonstration of quality of care measurement using the Japanese liver cancer registry. [Hepatol Res](#). 2011;41:1208–15.
- 25 Takayasu K, Aii S, Kudo M, Ichida T, Matsui O, Izumi N, et al. Superselective transarterial chemoembolization for hepatocellular carcinoma. Validation of treatment algorithm proposed by Japanese guidelines. [J Hepatol](#). 2012;56:886–92.
- 26 Takayasu K, Aii S, Sakamoto M, Matsuyama Y, Kudo M, Ichida T, et al. Clinical implication of hypovascular hepatocellular carcinoma studied in 4,474 patients with solitary tumour equal or less than 3 cm. [Liver Int](#). 2013;33:762–70.
- 27 Hasegawa K, Kokudo N, Makuuchi M, Izumi N, Ichida T, Kudo M, et al. Comparison of resection and ablation for hepatocellular carcinoma: a cohort study based on a Japanese nationwide survey. [J Hepatol](#). 2013;58:724–9.
- 28 Nouse K, Miyahara K, Uchida D, Kuwaki K, Izumi N, Omata M, et al. Effect of hepatic arterial infusion chemotherapy of 5-fluorouracil and cisplatin for advanced hepatocellular carcinoma in the nationwide survey of primary liver cancer in Japan. [Br J Cancer](#). 2013;109:1904–7.
- 29 Aoki T, Kokudo N, Matsuyama Y, Izumi N, Ichida T, Kudo M, et al. Prognostic impact of spontaneous tumor rupture in patients with hepatocellular carcinoma: an analysis of 1160 cases from a nationwide survey. [Ann Surg](#). 2014;259:532–42.
- 30 Utsunomiya T, Shimada M, Kudo M, Ichida T, Matsui O, Izumi N, et al. Nationwide study of 4741 patients with non-B non-C hepatocellular carcinoma with special reference to the therapeutic impact. [Ann Surg](#). 2014;259:336–45.
- 31 Hasegawa K, Makuuchi M, Kokudo N, Izumi N, Ichida T, Kudo M, et al. Impact of histologically confirmed lymph node metastases on patient survival after surgical resection for hepatocellular carcinoma: report of a Japanese nationwide survey. [Ann Surg](#). 2014;259:166–70.
- 32 Kitai S, Kudo M, Izumi N, Kaneko S, Ku Y, Kokudo N, et al. Validation of three staging systems for hepatocellular carcinoma (JIS score, biomarker-combined JIS score and BCLC system) in 4,649 cases from a Japanese nationwide survey. [Dig Dis](#). 2014;32:717–24.
- 33 Utsunomiya T, Shimada M, Kudo M, Ichida T, Matsui O, Izumi N, et al. A comparison of the surgical outcomes among patients with HBV-positive, HCV-positive, and non-B non-C hepatocellular carcinoma: a nationwide study of 11,950 patients. [Ann Surg](#). 2015;261:513–20.
- 34 Sakamoto Y, Kokudo N, Matsuyama Y, Sakamoto M, Izumi N, Kadoya M, et al. Proposal of a new staging system for intrahepatic cholangiocarcinoma: analysis of surgical patients from a nationwide survey of the Liver Cancer Study Group of Japan. [Cancer](#). 2016;122:61–70.
- 35 Kitai S, Kudo M, Nishida N, Izumi N, Sakamoto M, Matsuyama Y, et al. Survival benefit of locoregional treatment for hepatocellular carcinoma with advanced liver cirrhosis. [Liver Cancer](#). 2016;5:175–89.
- 36 Kudo M, Izumi N, Sakamoto M, Matsuyama Y, Ichida T, Nakashima O, et al. Survival analysis over 28 years of 173,378 patients with hepatocellular carcinoma in Japan. [Liver Cancer](#). 2016;5:190–7.
- 37 Kudo M, Izumi N, Ichida T, Ku Y, Kokudo N, Sakamoto M, et al. Report of the 19th follow-up survey of primary liver cancer in Japan. [Hepatol Res](#). 2016;46:372–90.
- 38 Kokudo T, Hasegawa K, Matsuyama Y, Takayama T, Izumi N, Kadoya M, et al. Survival benefit of liver resection for hepatocellular carcinoma associated with portal vein invasion. [J Hepatol](#). 2016;65:938–43.
- 39 Toyoda H, Tada T, Johnson PJ, Izumi N, Kadoya M, Kaneko S, et al. Validation of serological models for staging and prognostication of HCC in patients from a Japanese nationwide survey. [J Gastroenterol](#). 2017;52:1112–21.

- 40 Kokudo T, Hasegawa K, Matsuyama Y, Takayama T, Izumi N, Kadoya M, et al. Liver resection for hepatocellular carcinoma associated with hepatic vein invasion: a Japanese nationwide survey. *Hepatology*. 2017;66:510–7.
- 41 Hiraoka A, Michitaka K, Kumada T, Izumi N, Kadoya M, Kokudo N, et al. Validation and potential of albumin-bilirubin grade and prognostication in a nationwide survey of 46,681 hepatocellular carcinoma patients in Japan: the need for a more detailed evaluation of hepatic function. *Liver Cancer*. 2017;6:325–36.
- 42 Takayasu K, Arai S, Sakamoto M, Matsuyama Y, Kudo M, Kaneko S, et al. Impact of resection and ablation for single hypovascular hepatocellular carcinoma ≤ 2 cm analysed with propensity score weighting. *Liver Int*. 2018;38:484–93.
- 43 Kaibori M, Yoshii K, Hasegawa K, Ogawa A, Kubo S, Tateishi R, et al. Treatment optimization for hepatocellular carcinoma in elderly patients in a Japanese nationwide cohort. *Ann Surg*. 2019;270:121–30.
- 44 Kaibori M, Yoshii K, Yokota I, Hasegawa K, Nagashima F, Kubo S, et al. Impact of advanced age on survival in patients undergoing resection of hepatocellular carcinoma: report of a Japanese nationwide survey. *Ann Surg*. 2019;269:692–9.
- 45 Hiraoka A, Michitaka K, Kumada T, Izumi N, Kadoya M, Kokudo N, et al. Prediction of prognosis of intermediate-stage HCC patients: validation of the tumor marker score in a nationwide database in Japan. *Liver Cancer*. 2019;8:403–11.
- 46 Aoki T, Kubota K, Hasegawa K, Kubo S, Izumi N, Kokudo N, et al. Author response to: comment on: significance of the surgical hepatic resection margin in patients with a single hepatocellular carcinoma. *Br J Surg*. 2020;107:470–20.
- 47 Watanabe Y, Matsuyama Y, Izumi N, Kubo S, Kokudo N, Sakamoto M, et al. Effect of surgical margin width after R0 resection for intrahepatic cholangiocarcinoma: a nationwide survey of the Liver Cancer Study Group of Japan. *Surgery*. 2020;167:793–802.
- 48 Kudo M, Izumi N, Kubo S, Kokudo N, Sakamoto M, Shiina S, et al. Report of the 20th nationwide follow-up survey of primary liver cancer in Japan. *Hepatol Res*. 2020;50:15–46.
- 49 Fukami Y, Kaneoka Y, Maeda A, Kumada T, Tanaka J, Akita T, et al. Liver resection for multiple hepatocellular carcinomas: a Japanese nationwide survey. *Ann Surg*. 2020;272:145–54.
- 50 Liver Cancer Study Group of Japan. *Report of the 21st nationwide follow-up survey of primary liver cancer in Japan (2010–2011)*. Osaka: Liver Cancer Study Group of Japan; 2020. (in Japanese).
- 51 Wei Y, Ye Z, Yuan Y, Huang Z, Wei X, Zhang T, et al. A new diagnostic criterion with gadoxetic acid-enhanced MRI may improve the diagnostic performance for hepatocellular carcinoma. *Liver Cancer*. 2020;9:414–25.
- 52 Bae JS, Lee JM, Yoon JH, Jang S, Chung JW, Lee KB, et al. How to best detect portal vein tumor thrombosis in patients with hepatocellular carcinoma meeting the milan criteria: gadoxetic acid-enhanced MRI versus contrast-enhanced CT. *Liver Cancer*. 2020;9:293–307.
- 53 Kudo M, Ueshima K, Osaki Y, Hirooka M, Imai Y, Aso K, et al. B-mode ultrasonography versus contrast-enhanced ultrasonography for surveillance of hepatocellular carcinoma: a prospective multicenter randomized controlled trial. *Liver Cancer*. 2019;8:271–80.
- 54 Tateishi R, Uchino K, Fujiwara N, Takehara T, Okanoue T, Seike M, et al. A nationwide survey on non-B, non-C hepatocellular carcinoma in Japan: 2011–2015 update. *J Gastroenterol*. 2019;54:367–76.
- 55 Tamura Y, Igarashi M, Kawai H, Suda T, Satomura S, Aoyagi Y. Clinical advantage of highly sensitive on-chip immunoassay for fucosylated fraction of alpha-fetoprotein in patients with hepatocellular carcinoma. *Dig Dis Sci*. 2010;55:3576–83.
- 56 Tamura Y, Igarashi M, Suda T, Wakai T, Shirai Y, Umemura T, et al. Fucosylated fraction of alpha-fetoprotein as a predictor of prognosis in patients with hepatocellular carcinoma after curative treatment. *Dig Dis Sci*. 2010;55:2095–101.
- 57 Toyoda H, Kumada T, Tada T, Kaneoka Y, Maeda A, Kanke F, et al. Clinical utility of highly sensitive lens culinaris agglutinin-reactive alpha-fetoprotein in hepatocellular carcinoma patients with alpha-fetoprotein. *Cancer Sci*. 2011;102:1025–31.
- 58 Toyoda H, Kumada T, Osaki Y, Oka H, Urano F, Kudo M, et al. Staging hepatocellular carcinoma by a novel scoring system (BALAD score) based on serum markers. *Clin Gastroenterol Hepatol*. 2006;4:1528–36.
- 59 Kumada T, Toyoda H, Tada T, Kiriya S, Tanikawa M, Hisanaga Y, et al. High-sensitivity lens culinaris agglutinin-reactive alpha-fetoprotein assay predicts early detection of hepatocellular carcinoma. *J Gastroenterol*. 2014;49:555–63.
- 60 Kudo M. Management of hepatocellular carcinoma in Japan: current trends. *Liver Cancer*. 2020;9:1–5.
- 61 Yoon JH, Lee JM, Lee DH, Joo I, Jeon JH, Ahn SJ, et al. A comparison of biannual two-phase low-dose liver CT and US for HCC surveillance in a group at high risk of HCC development. *Liver Cancer*. 2020;9:503–17.
- 62 Galle PR, Foerster F, Kudo M, Chan SL, Llovet JM, Qin S, et al. Biology and significance of alpha-fetoprotein in hepatocellular carcinoma. *Liver Int*. 2019;39:2214–29.
- 63 Kudo M, Hasegawa K, Kawaguchi Y, Takayama T, Izumi N, Yamanaka N, et al. A multicenter randomized controlled trial to evaluate the efficacy of surgery vs. radiofrequency ablation for small hepatocellular carcinoma (SURF trial): analysis of overall survival. *ASCO Annual Meeting*. 2021 Jun 4–8.
- 64 Kudo M, Imanaka K, Chida N, Nakachi K, Tak WY, Takayama T, et al. Phase III study of sorafenib after transarterial chemoembolisation in Japanese and Korean patients with unresectable hepatocellular carcinoma. *Eur J Cancer*. 2011;47:2117–27.
- 65 Meyer T, Fox R, Ma YT, Ross PJ, James MW, Sturges R, et al. Sorafenib in combination with transarterial chemoembolisation in patients with unresectable hepatocellular carcinoma (TACE 2): a randomised placebo-controlled, double-blind, phase 3 trial. *Lancet Gastroenterol Hepatol*. 2017;2:565–75.
- 66 Lencioni R, Llovet JM, Han G, Tak WY, Yang J, Guglielmi A, et al. Sorafenib or placebo plus TACE with doxorubicin-eluting beads for intermediate stage HCC: the SPACE trial. *J Hepatol*. 2016;64:1090–8.
- 67 Kudo M, Cheng AL, Park JW, Park JH, Liang PC, Hidaka H, et al. Orantinib versus placebo combined with transcatheter arterial chemoembolisation in patients with unresectable hepatocellular carcinoma (ORIENTAL): a randomised, double-blind, placebo-controlled, multicentre, phase 3 study. *Lancet Gastroenterol Hepatol*. 2018;3:37–46.
- 68 Kudo M, Han G, Finn RS, Poon RT, Blanc JF, Yan L, et al. Brivanib as adjuvant therapy to transarterial chemoembolization in patients with hepatocellular carcinoma: a randomized phase III trial. *Hepatology*. 2014;60:1697–707.
- 69 Kudo M, Ueshima K, Ikeda M, Torimura T, Tanabe N, Aikata H, et al. Randomised, multicentre prospective trial of transarterial chemoembolisation (TACE) plus sorafenib as compared with TACE alone in patients with hepatocellular carcinoma: TACTICS trial. *Gut*. 2020;69:1492–501.
- 70 Kudo M, Ueshima K, Ikeda M, Torimura T, Tanabe M, Aikata H, et al. TACTICS: final overall survival data from a randomized, open label, multicenter, phase II trial of transcatheter arterial chemoembolization (TACE) therapy in combination with sorafenib as compared with TACE alone in patients with hepatocellular carcinoma. *ASCO-GI*. 2021.
- 71 Kudo M, Ueshima K, Yokosuka O, Ogasawara S, Obi S, Izumi N, et al. Sorafenib plus low-dose cisplatin and fluorouracil hepatic arterial infusion chemotherapy versus sorafenib alone in patients with advanced hepatocellular carcinoma (SILIUS): a randomised, open label, phase 3 trial. *Lancet Gastroenterol Hepatol*. 2018;3:424–32.
- 72 Ueshima K, Ogasawara S, Ikeda M, Yasui Y, Terashima T, Yamashita T, et al. Hepatic arterial infusion chemotherapy versus sorafenib in patients with advanced hepatocellular carcinoma. *Liver Cancer*. 2020;9:583–95.
- 73 He M, Li Q, Zou R, Shen J, Fang W, Tan G, et al. Sorafenib plus hepatic arterial infusion of oxaliplatin, fluorouracil, and leucovorin vs sorafenib alone for hepatocellular carcinoma with portal vein invasion: a randomized clinical trial. *JAMA Oncol*. 2019;5:953–60.

- 74 Cheng S, Chen M, Cai J, Sun J, Guo R, Bi X, et al. Chinese expert consensus on multidisciplinary diagnosis and treatment of hepatocellular carcinoma with portal vein tumor thrombus (2018 edition). *Liver Cancer*. 2020; 9:28–40.
- 75 Zhou J, Sun H, Wang Z, Cong W, Wang J, Zeng M, et al. Guidelines for the diagnosis and treatment of hepatocellular carcinoma (2019 edition). *Liver Cancer*. 2020;9(6):682–720.
- 76 Finn RS, Qin S, Ikeda M, Galle PR, Ducreux M, Kim TY, et al. Atezolizumab plus bevacizumab in unresectable hepatocellular carcinoma. *N Engl J Med*. 2020;382:1894–905.
- 77 Kudo M, Finn RS, Qin S, Han KH, Ikeda K, Piscaglia F, et al. Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. *Lancet*. 2018;391:1163–73.
- 78 Finn R, Qin S, Ikeda M, Galle P, ducreux M, Kim TY, et al. IMbrave150: updated overall survival (OS) data from a global, randomized, open-label phase III study of atezolizumab (atezo) + bevacizumab (bev) vs sorafenib (sor) in patients (pts) with unresectable hepatocellular carcinoma (HCC). *EASL Digital Liver Cancer Summit*. 2021 Feb 5–6.
- 79 Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, et al. Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med*. 2008;359:378–90.
- 80 Bruix J, Qin S, Merle P, Granito A, Huang YH, Bodoky G, et al. Regorafenib for patients with hepatocellular carcinoma who progressed on sorafenib treatment (RESORCE): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2017;389:56–66.
- 81 Zhu AX, Kang YK, Yen CJ, Finn RS, Galle PR, Llovet JM, et al. Ramucirumab after sorafenib in patients with advanced hepatocellular carcinoma and increased alpha-fetoprotein concentrations (REACH-2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol*. 2019;20:282–96.
- 82 Abou-Alfa GK, Meyer T, Cheng AL, El-Khoueiry AB, Rimassa L, Ryoo BY, et al. Cabozantinib in patients with advanced and progressing hepatocellular carcinoma. *N Engl J Med*. 2018;379:54–63.
- 83 Kudo M. A new era in systemic therapy for hepatocellular carcinoma: atezolizumab plus bevacizumab combination therapy. *Liver Cancer*. 2020;9:119–37.
- 84 Kudo M. Recent advances in systemic therapy for hepatocellular carcinoma in an aging society: 2020 update. *Liver Cancer*. 2020;9(6): 640–62.
- 85 Kudo M, Han KH, Ye SL, Zhou J, Huang YH, Lin SM, et al. A changing paradigm for the treatment of intermediate-stage hepatocellular carcinoma: Asia-Pacific primary liver cancer expert consensus statements. *Liver Cancer*. 2020;9:245–60.
- 86 Kudo M, Kawamura Y, Hasegawa K, Tateishi R, Kariyama K, Shiina S, et al. Management of hepatocellular carcinoma in Japan: JSH consensus statements and recommendations 2020 update. *Liver Cancer*. Forthcoming 2021.
- 87 Llovet JM, Villanueva A, Marrero JA, Schwartz M, Meyer T, Galle PR, et al. Trial design and endpoints in hepatocellular carcinoma: AASLD consensus conference. *Hepatology*. 2020;158–91.
- 88 Kudo M. Scientific rationale for combination immunotherapy of hepatocellular carcinoma with anti-PD-1/PD-L1 and anti-CTLA-4 antibodies. *Liver Cancer*. 2019;8(6):413–26.
- 89 Kudo M. Combination cancer immunotherapy with molecular targeted agents/anti-CTLA-4 antibody for hepatocellular carcinoma. *Liver Cancer*. 2019;8:1–11.
- 90 Kudo M. Immuno-oncology therapy for hepatocellular carcinoma: current status and ongoing trials. *Liver Cancer*. 2019;8:221–38.