Risk Factors of COVID-19 Symptoms in Old Subjects

Tomoyuki Kawada

Department of Hygiene and Public Health, Nippon Medical School, Tokyo, Japan

Dear Editor,

I read with great interest the study by Del Ser et al. [1] that explored the risk factors for severe COVID-19 in older people. Regarding the APOE $\epsilon 4$ allele, some queries were previously presented [2]. I want to discuss the risk of smoking and disease severity with special reference to statistical validity.

The authors described that more intense smoking was significantly associated with severe COVID-19 symptoms by applying linear regression analysis with the Poisson distribution. The age- and sex-adjusted odds ratio (95% confidence interval) of smoking intensity for severity of COVID-19 symptoms was 0.95 (0.91–0.99). I accept that the mean value of the smoking severity index was about 2-fold higher in the moderate/severe group than in the mild group; however, the multivariate statistical results presented an inverse relationship between smoking intensity and severity of COVID-19 symptoms.

Regarding smoking, COVID-19 infection, and disease progression, Simons et al. [3] conducted a meta-analysis by overviewing past epidemiological studies. Compared with never smokers, current smokers presented a reduced risk of COVID-19 infection, presenting a relative risk (95% confidence interval) of 0.74 (0.58–0.93). In contrast, the risk of hospitalization, disease severity, and mortality of COVID-19 in current smokers did not reach levels of significance.

A previous meta-analysis found that the risks of hospitalization, severe disease, and mortality of COVID-19

in former smokers might be significantly high [3]. The risk of disease progression in current smokers should be specified by further studies.

Acknowledgments

I greatly appreciate technical support by members of the Department of Hygiene and Public Health, Nippon Medical School.

Conflict of Interest Statement

The author has no conflicts of interest to declare.

Funding Sources

There was no funding for this letter.

References

- 1 Del Ser T, Fernandez-Blazquez MA, Valenti M, Zea-Sevilla MA, Frades B, Alfayate E, et al. Residence, clinical features, and genetic risk factors associated with symptoms of COVID-19 in a cohort of older people in Madrid. *Gerontology*. 2021;67(3):281–9.
- 2 Hubacek JA, Dlouha L, Dusek L, Majek O, Adamkova V. Apolipoprotein E4 allele in subjects with COVID-19. *Gerontology*. 2021; 67(3):320–2.
- 3 Simons D, Shahab L, Brown J, Perski O. The association of smoking status with SARS-CoV-2 infection, hospitalization and mortality from COVID-19: a living rapid evidence review with Bayesian meta-analyses (version 7). *Addiction*. 2021;116(6):1319–68.