Introduction

L. Boccon-Gibod

Department of Urology, Hôpital Bichat-Claude Bernard, Paris, France

There is no doubt that medical therapy, and more specifically alpha-blocking agents, have a dramatic impact on lower urinary tract symptoms (LUTS) associated with benign prostatic enlargement (BPE) or hypertrophy (BPH), to use the current urological dialect. Thus, the therapeutic approach towards BPH favoured by physicians, particularly urologists, has spectacularly changed since the introduction of medical treatments with evidence-based efficacy in the last 10–15 years.

The impact of medical treatment is all the more spectacular because of its effect on the most bothersome symptoms to the patient, i.e. storage symptoms, such as nocturia and urgency with or without incontinence. To a lesser degree, it also reduces symptoms of more concern to the urologist, i.e. emptying symptoms such as low flow. These symptoms directly reflect the presence of bladder outlet obstruction (BOO), with all its consequences on urinary function: acute or chronic urinary retention, obstructive renal insufficiency, detrusor deterioration leading to hyper- or acontractility, bladder stones and diverticula. The interest manifested by urologists toward emptying symptoms, however, is related to the fact that urologists are particularly good at relieving obstruction and have been for decades!

Regardless of how spectacular the symptomatic relief experienced by the patient, the urologists remain faced with a number of important questions:

- To what extent does the relief of LUTS relate to the real impact of medical treatment, specifically alpha-blocking agents, on the natural history of BPH/BOO? It is well accepted that BPH/BOO is not a life-threatening disease and that surgery does not improve life expectancy. Therefore, the impact of medical treatment on the natural history of the disease can only be measured by a reduction in the indications for surgery.

- If indeed alpha-blocking agents have a major impact on the natural history of the disease, may their use be extended to the treatment of obstructive complications, such as the most feared of all obstructive complications – acute urinary retention?
- If alpha blockers do impact on the natural history of the disease, how do they compare with the increasingly popular instrumental ‘minimally invasive therapies’, such as transurethral microwave thermotherapy and various lasers? These usually require no anaesthesia, can be used on an outpatient basis, and propose to treat the patient in a single session compared with continuous medical therapy.

The comparison between alpha blockers and minimally invasive therapies has to be made on three different levels:

- The safety of long-term medical treatment – a major issue when one considers the increasing life expectancy of the population.
- Patient satisfaction and acceptance of long-term medical therapy vs. a single treatment session.
- Medico-economic costs considerations: long-term medical treatment vs. a single session with minimally invasive therapies. Also to be incorporated into the costs of both groups is the ultimate risk of surgical therapy, which will probably increase with the increasing life expectancy of the population.

These questions have been addressed specifically during the symposium held in Brussels at the Congress of the European Association of Urology thanks to an unrestricted grant of Sanofi-Synthelabo Recherche. I hope that the readers will take the same interest and pleasure in reading these very interesting papers as the chairman took in listening to them.