Introduction

Bettina Borisch

The 2nd International Conference on Laser Microdissection took place in Chavannes-de-Bogis near Geneva. Since the first meeting held in Les Diablerets in April 1998, much progress has been made both in the laser microdissection microscope technology as well as in the downstream methods of analyzing the microdissected material.

In addition to a strong European participation, about 80 participants from 14 countries including Asia, Australia and America attended the meeting. The opportunities offered by the workshops were highly appreciated and allowed every participant to have hands-on experience with four different laser microdissection microscopes.

The present volume contains original articles that cover the contents of talks given at the meeting. The reader will no doubt appreciate the broad variety of topics discussing the powerful tool of microdissection. It will be readily understood that the development of micromethods in analytic and diagnostic biomedicine (such as chips/arrays, MALDI) requires morphologically verified and analyzed 'micromaterial'. Almost all questions addressed to whole tissue preparations some years ago are no longer useful and have to be redirected to well-defined parts of tissues or structures.

The first session on solid tumours readily underlined this aspect among other things. The opening presentation by Lemoine [1] gave an overview of the opportunities now available for the study of solid tumours and illustrated the (inherited) pancreatic carcinoma and the hepatocellular carcinoma. Studying precursor lesions such as regenerative nodules of the liver and intraductal pancreatic mucinous tumours using chip technology, he coined the term

of 'expressome' for this way of addressing the tumour biology of particular situations as a model for investigating solid tumours in general. The following presentations in the session were concerned with the problems of clonality and microheterogeneity of solid tumours. Knüchel et al. (p. 165) discussed the importance of the clonality during the establishment of urothelial cancerous lesions. An important part of the meeting was dedicated to breast carcinoma and its precursor lesions. Glöckner et al. (p. 173) described the combined use of microdissection and real-time PCR to study intraductal lesions and their adjacent invasive components. Diallo [2] showed that the HUMARA assay for clonality in breast lesions might be of limited value as terminal ductal lobular units are monoclonal to start with. These findings indeed support the hypothesis of distinct stem cell-derived monoclonal patches as a model for the organization of the normal mammary gland. By analyzing microsatellite instability (MSI) and loss of heterozygosity (LOH), Hartmann et al. (p. 180) showed a high degree of heterogeneity in breast tumours.

The haematopathology session was introduced by Hansmann [3, 4] who was a pioneer in microdissection and single cell analysis, starting out with manual microdissection of Hodgkin cells. Besides the old data on HD/RS cells and their origin, new results on composite lymphomas and lymphocyte-predominant Hodgkin's disease as well as the T cell aspect of Hodgkin's disease were presented. The second presentation (Fey et al.) summarized the results obtained on LOH of HD and RS cells. Microsatellites at 1q42, 4q26, 9p23 and 11q23 were analyzed. Clonal LOH was seen in all but one case, and the locus

most frequently altered was 4q26 [5]. Finally Leoncini et al. [6] presented the case of a patient with simultaneously occurring HD and large B cell lymphoma. Their analysis may suggest that RS and large B cell lymphoma cells originate from a common precursor in which a secondary V_H replacement took place during germinal centre reaction.

The following session showed that microanalysis is not only useful in tumour cell biology but also in the analysis of non-neoplastic tissues. After an overview talk by Köhlble [7], Bohle et al. (p. 191) presented an animal model which allows not only a cell type-specific mRNA quantification but also a cell type-specific characterization by combining this analysis with immunostaining. The Hannover group (Kleeberger et al., p. 196) showed that using a polymorphic short tandem repeat marker (STR) which is widely used in forensic medicine (SE33), graft chimerism after solid organ transplantation can be evaluated.

The keynote lecture by Liotta [8] covered the developments to come: functional proteomics, proteomics applied to molecular targets and protein pathway profiling. Combining these tools should allow us to describe more closely the functional state of a given protein pathway. He foresees that these questions will be answered using microdissection and combined downstream tools while concentrating especially on the proteomic approach. Lehmann and Kreipe (p. 202) pointed out that while doing a quantitative molecular analysis of laser-microdissected human tissues epigenetic modifications such as methyl-

ation can be evaluated. Fend (p. 209) gave a thorough review of the combination of immunohistochemistry and laser capture microdissection with all its pros and cons. More on the technical aspects, Ren et al. (p. 215) showed how they optimized the PCR conditions for microdissected formalin-fixed and paraffin-embedded materials.

In some instances, these new techniques are already used for diagnostics. Lahr et al. (p. 218) showed the usefulness of laser microdissection in the diagnosis of papillary thyroid carcinoma using an RT-PCR approach in archival material, and Rubbia et al. [9] the microanalysis of hepatic lesions. Dietmaier et al. (p. 227) discussed the application of LOH and MSI on microdissected colorectal tumours. Crnogorac-Jurcevic [1] extended the data of the Lemoine group by presenting cDNA arrays for generating a comprehensive gene expression profile of pancreatic adenocarcinoma. An important contribution (Clement-Sengewald et al., p. 232) focused on another application of laser microdissection, namely the use of the laser to extract polar bodies for a prefertilization diagnosis.

The microaspects being extensively covered, the macroaspects of life were highlighted by the sunny September weather that showed magnificent views of the Mont-Blanc and offered excellent opportunities to do tours to nearby vineyards and restaurants. The articles published herein give a complete overview of the present state of the art in laser microdissection and its applications and developments.

References

- 1 Sirivatanauskorn Y, Drury R, Crnogorac-Jurcevic T, Sirivatanauskorn V, Lemoine NR: Laser assisted-microdissection: Applications in molecular pathology. *J Pathol* 1999;1892:150–154.
- 2 Diallo R, Schaefer K-L, Poremba C, Shivazi N, Willmann V, Buerger H, Dockhorn-Dworniczak B, Boecker W: Monoclonality in the normal epithelium and in hyperplastic and neoplastic lesions of the breast. *J Pathol* 2001;193:27–32.
- 3 Küppers R, Rajewsky K, Zhao M, Simons G, Laumann R, Fischer R, Hansmann M-L: Hodgkin disease: Hodgkin- and Reed-Sternberg cells picked from histological sections show clonal immunoglobulin rearrangements and appear to be derived from B cells at various stages of development. *Proc Natl Acad Sci USA* 1994;91:10962–10966.
- 4 Küppers R, Klein U, Hansmann M-L, Rajewsky K: Cellular origin of human B-cell lymphomas. *N Engl J Med* 1999;341:1520–1529.
- 5 Hasse U, Tinguely M, Leibundgut EO, Cajot JF, Garvin AM, Tobler A, Borisch B, Fey MF: Clonal loss of heterozygosity in microdissected Hodgkin and Reed-Sternberg cells. *J Natl Cancer Inst* 1999;91:1581–1583.
- 6 Bellan C, Lazzi S, Zazzi M, Lalinga AV, Palumbo N, Galieni P, Marafioti T, Tonini T, Cinti C, Leoncini L, Pileri SA, Tosi P: Immunoglobulin gene rearrangement analysis in composite Hodgkin's disease and large B-cell lymphoma. Evidence for receptor revision of immunoglobulin heavy chain variable region genes in Hodgkin-Reed-Sternberg cells? *Diagn Mol Pathol*, in press.
- 7 Köhlble K: The LEICA microdissection system: Design and applications. *J Mol Med* 2000;787:B24–B25.
- 8 Emmert-Buck MR, Strausberg RL, Krizman DB, Bonaldo MF, Bonner RF, Bostwick DG, Brown MR, Buetow KH, Chuaqui RF, Cole KA, Duray PH, Englert CR, Gillespie JW, Greenhut S, Grouse L, Hillier LW, Katz KS, Klausner RD, Kuznetsov V, Lash AE, Lennon G, Linehan WM, Liotta LA, Marra MA, Munson PJ, Ornstein DK, Prabhu VV, Prange C, Schuler GD, Soares MB, Tolstoshev CM, Vocke CD, Waterston RH: Molecular profiling of clinical tissue specimens: Feasibility and applications. *Am J Pathol* 2000;156:1109–1115.
- 9 Paradis V, Dargere D, Bonvoust F, Rubbia-Brandt L, Ba N, Bioulac-Sage P, Bedossa P: Clonal analysis of micronodules in virus C-induced liver cirrhosis using laser capture microdissection (LCM) and HUMARA assay. *Lab Invest* 2000;80:1553–1559.