Summary

Ethan Rubinstein, Symposium Chairperson

Department of Internal Medicine and Unit of Infectious Diseases,
Tel Aviv University School of Medicine, Tel Aviv, Israel

Providing a base upon which the other presentations could build, I reviewed the historical development of fluoroquinolones, and then analysed the structure of individual agents in terms of their efficacy and tolerability, allowing the agents to be categorised based upon the likelihood of their causing adverse events due to their structure. This was followed by Professor Claude Carbon, who reported on differences between individual fluoroquinolones that have been described following use in patients. Such data is imperative in clearly defining which agents are safe to use, and highlight the fact that there are major differences now recognised. A number of agents have been withdrawn, such as trovafloxacin, and most recently, clinafloxacin. Of the agents remaining, levofloxacin appears to be extremely safe with a low rate of adverse events, and low rates of cardiac and hepatic problems. Professor Jacques F. Acar widened the discussion by comparing fluoroquinolones to other commonly used agents, such as macrolides and β -lactams. It is important to have a clear understanding of the side-effect profiles associated with all major classes of

antimicrobials, and Professor Acar provided this. He then discussed one of the newer issues gaining importance in the infectious disease community — namely, the ecological impact of antimicrobials upon the normal human gastrointestinal flora. Levofloxacin was shown to have less of an impact on gastrointestinal flora than many other compounds, and was not associated with overgrowth of resistant mutants, unlike other agents. Professor Hartmut Lode discussed pharmacological differences among the fluoroquinolones. He drew attention to the many advantages possessed by the newer fluoroquinolones, such as levofloxacin. They maintain excellent pharmacokinetic profiles, making them easy and effective to use.

Actual clinical data following extensive use of levofloxacin was then used to assess its safety profile and compare it to the clinical adverse-event profiles for the other agents. This was first assessed from a US perspective, with results reported by Dr. James B. Kahn, followed by Japanese and global data reported by Dr. Katsuro Yagawa. They showed that even following six years of levofloxacin use,

the incidence of adverse events with this drug remains extremely low, with 1.3% of adverse events reported in Japan.

Following the presentations, there was a lively discussion focusing upon some of the most clinically important differences. The discussion allowed faculty and participants to join in a relaxed and stimulating environment to discuss and answer questions and issues related to the safety of fluoroquinolones. Par-

ticipants attending from around the world took the opportunity to ask the distinguished panel their opinion on a range of important issues related to the safety of the fluoroquinolones. Throughout the roundtable discussion, it was apparent that levofloxacin is one of the most clinically effective fluoroquinolones, which is coupled with an extremely good safety profile.