Editorial

Leo P. ten Kate

I was recently invited to review secondary prevention of congenital and genetic disorders in The Netherlands. Martina Cornel, who edited the previous special issue of this journal on the prevention of birth defects, reviewed primary prevention. We decided that the concept of prevention can only be applied to living people or people yet to be born or conceived. For instance, calling termination of pregnancy after prenatal diagnosis 'prevention' is a perversion of terminology. I suggest that we should use the term 'reproductive choice' in such instances. By analogy with prevention, one might define different levels of reproductive choice [1]. Primary reproductive choice would then consist of actions to avoid conception of affected offspring, while secondary reproductive choice would bar implantation or birth of affected embryos and fetuses.

Secondary prevention of congenital and genetic disorders in The Netherlands includes neonatal screening (for phenylketonuria, congenital hypothyroidism and congenital adrenal hyperplasia), family testing for familial hypercholesterolemia (FH) and a host of other disorders. Implementation of a nation-wide program of cascade testing for FH is held up by uncertainty about the position of life insurers. The country's eight clinical genetics centers perform the great majority of DNA testing. They offer testing for over 230 disorders/genes, and about 20,000 tests are performed yearly (excluding FH). A quick and rough estimate reveals that about a quarter of the 230 disorders/genes would be amenable to secondary prevention and this applies to even 40% of the 20,000 tests. If one assumes that only 10% of these tests result in secondary prevention, the number of cases in which the development of disease was prevented by family testing, excluding FH, is about ten times higher than the number prevented through neonatal screening. Secondary prevention will soon affect more persons than secondary reproductive choice. This clearly shows the growing role of genetics in prevention.

Reference

- 1 Last JM: A Dictionary of Epidemiology, ed 4. Oxford, Oxford University Press, 2001.