Assignment¹ of maltase glucoamylase (MGAM) gene to bovine chromosome 4q34 by in situ hybridization and confirmation by radiation hybrid mapping

R. Roy,^a M. Gautier,^b H. Hayes,^b P. Laurent,^b P. Zaragoza,^a A. Eggen,^b and C. Rodellar^a

^aLaboratorio de Genética Bioquímica y Grupos Sanguíneos, Facultad de Veterinaria, Universidad de Zaragoza, Zaragoza (Spain); ^bLaboratoire de Génétique Biochimique et de Cytogénétique, INRA-CRJ, Jouy-en-Josas (France)

¹ To our knowledge this is the first time this gene has been mapped in cattle.

Rationale and significance

In mammals, hydrolysis of the nonreducing ends of oligosaccharide chains is carried out by an enzyme complex involving sucrase-isomaltase and maltase-glucoamylase which is anchored in the mucosa of the small intestine brush border (Semenza and Aurichino, 1989). It has been hypothesized that the human MGAM activity is an alternate pathway for starch digestion when luminal α -amylase activity is reduced because of immaturity and malnutrition and that MGAM plays a unique role in the digestion of malted oligosaccharides (Nichols et al., 1998). The MGAM gene has been cloned and sequenced in man (AF016833) and partially sequenced in mouse (BC025509.1). Human MGAM has been mapped to human (HSA) chromosome 7 (Nichols et al., 1998), but as yet, the bovine MGAM gene has not been mapped. We used the human MGAM gene sequence to design specific primers to partially amplify the bovine gene. We report the localization of the bovine MGAM gene on bovine chromosome BTA4q34 by fluorescence in situ hybridization and radiation hybrid mapping.

Materials and methods

Isolation of a specific bovine MGAM fragment

In order to obtain a specific bovine MGAM DNA fragment, PCR amplification was performed using the following set of primers:
MGAM.F: 5'-AGGGACAGGGGCCTGAGCAG-3'
MGAM.R: 5'-TCGTATGTGGGTCTGGTCTGG-3'

Primers were designed from the human MGAM sequence and amplified a 120-bp fragment. PCR amplification was performed using 20 ng of genomic DNA. Cycling conditions were 94 °C for 30 s, 55 °C for 30 s and 72 °C for 30 s for 35 cycles. PCR products were cloned and sequenced. Homology searches were performed with BLAST programs at the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/BLAST/>).

The isolated bovine sequence showed a high level of sequence homology with the human and mouse MGAM genes, 85 % and 80 %, respectively and it was used to design bovine-specific primers.

Screening of a bovine BAC library

Using the bovine-specific primers, a bovine BAC library (Eggen et al., 2001) was screened by PCR and a BAC clone containing the bovine MGAM gene was identified. Partial sequence analysis of the BAC clone (22G4) confirmed the presence of MGAM sequences (Accession number: AY188356).

Fluorescence in situ hybridization (FISH)

In situ hybridization was carried out on RBP-banded bovine chromosome preparations, as described previously by Hayes et al. (2000).

Probe name: 22G4

Probe type: genomic DNA

Vector: pBelo BAC11

Insert size: 150 kb

Proof of authenticity: DNA sequencing

Gene reference: AY188356

Radiation hybrid (RH) mapping

To confirm the cytogenetic localization obtained by FISH, a bovine $\times$ hamster radiation hybrid panel was analyzed. PCR reactions were performed on the 94 radiation hybrid cell lines that constitute the newly developed 3000 Rad bovine panel (Williams et al., 2002).

Primers and PCR conditions for radiation hybrid mapping were the same as those used for screening the BAC library.

The Carthagen software (Schiex et al., 2002) was used to perform two-point and multipoint analyses of the radiation hybrid data and to provide a map of the region where the MGAM gene is located. The distances between markers on the most likely map were calculated with the RHMAP3.0 software (Lange et al., 1995) under the equal retention probability model.

R.R. was supported by a doctoral grant from the M.E.C.

Received 16 December 2002; manuscript accepted 16 January 2003.

Corresponding author: Rosa Roy, Laboratorio de Genética Bioquímica
Facultad de veterinaria, c/ Miguel Servet, 50013 Zaragoza (Spain)
telephone: 34 97 676 1620; fax: 34 97 676 1612
email: 181228@vetez.unizar.es

FISH MGAM (4q34)

Fig. 1. FISH mapping of MGAM using a bovine BAC clone containing the MGAM gene. A specific signal is observed on chromosome 4q34.

Fig. 2. RH mapping of MGAM gene to bovine chromosome 4. Genetic distances in centiRay around MGAM are shown on the right.

Results

Mapping data

Most precise location: 4q34

Number of cells examined: 20

Number of cells with specific signal: 1 (0), 2 (2), 3 (6), 4 (12) chromatids per cell

Location of background signals: none observed

Radiation hybrid mapping

Vector result: 00000 00010 00010 00001 00101 00000 00000 00000 00000 00100 00001 00010 00000 10000 00000 10000 00000 10011 1000. Most significantly linked marker by two-point analysis: RM188 on BTA 4 (29.9 cRay; LOD = 9)

The localization of the MGAM gene on bovine chromosome 7 agrees totally with comparative mapping data between cattle and man since BTA4 corresponds to part of human chromosome 7.

References

- Eggen A, Gautier M, Billaut A, Petit E, Hayes H, Laurent P, Urban C, Pfister-Genskow M, Eilertsen K, Bishop MD: Construction and characterization of a bovine BAC library with four genome-equivalent coverage. *Genet Sel Evol* 33:543–548 (2001).
- Hayes H, Di Meo GP, Gautier M, Laurent P, Eggen A, Iannuzzi L: Localization by FISH of 31 Texas nomenclature type I markers to both Q and R banded bovine chromosomes. *Cytogenet Cell Genet* 90:315–320 (2000).
- Lange K, Boehnke M, Cox DR, Lunetta KL: Statistical methods for polyploid radiation hybrid mapping. *Genome Res* 5:136–150 (1995).
- Nichols BL, Eldering J, Hahn D, Quaroni A, Sterchi E: Human small intestinal maltase-glucoamylase cDNA cloning. Homology to sucrase-isomaltase. *J Biol Chem* 273:3076–3081 (1998).
- Semenza G, Aurichino S: The metabolic basis of inherited disease. II. Scriver CR, Beaudet AL, Sly WS, Valle D et al (eds): pp 2975–2997 (McGraw-Hill, New York 1989).
- Schiex T, Chabrier P, Bouchez M, Milan D: Boosting EM for radiation hybrid and genetic mapping. *Theor Appl Genet*, in press (2002).
- Williams JL, Eggen A, Ferreti L, Farr C, Gautier M, et al: A bovine whole genome radiation hybrid panel and outline map. *Mammal Genome* 13:469–474 (2002).