Editorial

Acute emergencies in gastroenterology are extraordinarily severe conditions with high morbidity and mortality. Particularly severe diseases include acute pancreatitis, a difficult course of non-specific intestinal inflammations manifested by toxic colon or acute intestinal obstruction, and even acutely developed intestinal pseudo-obstruction (Ogilvie's syndrome) or variceal and non-variceal bleeding into the gastrointestinal tract. Undoubtedly serious factors influencing the accuracy of diagnostics and effectiveness of therapy are the etiological multifactorial characteristics of changes that induce the acute state. Polymorbidity is also frequent among these patients and requires a complex diagnostic approach, often limiting the possibility of using an optimal therapeutic approach.

Effective diagnostics and therapy for acute conditions in gastroenterology requires a multidisciplinary team approach. In diagnostics, endoscopic examination enabling a simultaneous therapeutical solution is of fundamental importance in managing most diseases, which is valid for example in patients with acute bleeding into the alimentary tract, in acute pancreatitis, acute cholangitis or acute intestinal obstruction. However, endoscopy is an invasive method, and as many of these patients suffer from polymorbidity, the usage of endoscopic approaches is limited by the general clinical condition of patients, particularly with respect to cardiopulmonary risks. In such cases, the application of non-invasive diagnostic methods is suitable. These involve imaging methods such as ultrasound abdominal examination, computer tomography or nuclear magnetic resonance. Moreover, modifications of these methods, e.g. CT enteroclysis or CT colonography,

provide very precise and immediate results that allow the adoption of an optimal strategic course. Due to their increasing sensitivity and specificity, the above-mentioned methods may be expected to substitute, in future, endoscopic examinations, whose present efficiency remains of the highest value.

Optimal therapy for acute states in gastroenterology is unthinkable without the close cooperation of a number of disciplines, particularly gastroenterology and surgery. Correct timing in determining whether conservative therapy is an effective and safe treatment for a patient in a given situation or whether immediate surgery should be performed is the basic requirement for the disease outcome of a patient. Severe states in particular should be managed at centers that have sufficient experience with such problems, possess a complete range of diagnostic methods, carry out therapeutic endoscopy, and have available acute surgical care, i.e. provide complex diagnostic and therapeutical services.

Although acute conditions in gastroenterology and gastroenterological complications are undoubtedly extraordinarily severe states, systematically processed data about rational and correct diagnostics and therapy from the viewpoint of gastroenterologists and surgeons have not been sufficient and therefore they could not be generalized and utilized as recommendations for a rational approach in these states.

We believe that the topics published in this issue of *Digestive Diseases* will help, at least in part, fill this gap.

Petr Dítě