Cox GF, Bürger J, Lip V, Mau UA, Sperling K, Wu BL, Horsthemke B:

Intracytoplasmic sperm injection may increase the risk of imprinting defects.

Am J Hum Genet 2002;71:162–164.

Ørstavik KH, Eiklid K, Van der Hagen CB, Spetalen S, Kierulf K, Skjeldal O, Buiting K:

Another case of imprinting in a girl with Angelman syndrome who was conceived by intracytoplasmic sperm injection.

Am J Hum Genet 2003;72:218–219.

DeBaun MR, Niemitz EL, Feinberg AP:

Association of in vitro fertilization with Beckwith-Wiedemann syndrome and epigenetic alterations of LIT1 and H19.

Am J Hum Genet 2003;72:156–160.

Maher ER, Brueton LA, Bowdin SC, Luharia A, Cooper W, Cole TR, Macdonald F, Sampson JR, Barratt CL, Reik W, Hawkins MM:

Beckwith-Wiedemann syndrome and assisted reproduction technology (ART).

J Med Genet 2003;40:62–64.

Moll AC, Imhof SM, Cruysberg JRM, Schouten-van Meeteren AYN, Boers M, Van Leeuwen FE: Incidence of retinoblastoma in children born after in-vitro fertilisation.

Lancet 2003;361:309–311.

DOI: 10.1159/000073334

Increased Risk of Rare Diseases in Children Born after Assisted Reproduction Technology

Cox et al. (Germany) reported two children who were conceived by intracytoplasmic sperm injection (ICSI) and who developed Angelman syndrome. Molecular analysis revealed a sporadic im-

printing defect of the SNRPN locus on the maternal chromosome, a rare cause of Angelman syndrome (<5% of all cases). This result raised the possibility that ICSI might interfere with the establishment of the maternal imprint in the oocyte or pre-embryo and increase the risk of imprinting defects.

Ørstavik et al. (Norway) described another girl with Angelman syndrome due to a sporadic imprinting defect, born after ICSI and in vitro fertilisation (IVF). Molecular studies showed complete absence of a methylated maternal band at the SNRPN locus. This report supports the assumption that ICSI can lead to an increased risk for imprinting defects.

Since the introduction of the Washington University Beckwith-Wiedemann syndrome (BWS) Registry in 1994, several BWS patients born after assisted reproduction technology (ART) were noted. Therefore, DeBaun et al. (USA) designed a questionnaire in 2001 to assess the method of conception. A total of 7 children with BWS were identified as having been born after ART (all after IVF of whom 5 including ICSI). Three patients were identified during the questionnaire period (2001), resulting in a prevalence of ART of 4.6% (3 of 65). Compared to 0.76% live births resulting from ART in the United States in 1999, a sixfold increase in BWS in children born after ART was found. Molecular studies of 6 of the BWS patients showed imprinting mutations of the LIT1 and/or H19 genes in 5 children.

Maher et al. (UK) also investigated whether there is evidence of an association between BWS and ART. The records of 146 BWS patients were reviewed. A history of ART was recorded for 6 cases (4%) (3 were born after ICSI and 3 after IVF without ICSI) compared with 0.997% in the general population ($p = 0.018$) in the UK. Two cases were assessed for KvDMR1 methylation status and both cases showed loss of methylation on the maternal allele, which is the most common molecular abnormality in sporadic BWS.

Moll et al. (the Netherlands) described a possible association of an increased risk of retinoblastoma in children born after IVF. Between November 2000 and February 2002, 5 patients born after IVF were diagnosed with retinoblastoma (2 bilateral, 3 unilateral). None of the patients had a family history of retinoblastoma. DNA analysis revealed a pathogenic mutation of the retinoblastoma gene in only 1 of the 5 patients. A significantly increased risk ratio of retinoblastoma after IVF of 7.2 (95% CI 2.4–17.0) was calculated.

Comments

As the number of infants born after ART is steadily increasing in most Western countries, the above-mentioned articles suggesting an association between ART and rare diseases need further attention. The first report (Cox et al.) discussing the possibility of ART interfering with the establishment of the maternal imprint was not very

alarming as it described only 2 Angelman patients. However, additional reports describing an increased frequency of ART in Angelman or BWS children, strengthened this hypothesis. The apparent increased frequency of Angelman and BWS, both imprinting disorders, in children born after ART, does not seem to be a chance finding and underlines the need for further research. It is remarkable that in almost all BWS and Angelman patients born after ART studied, there is an imprinting defect of the maternal allele. The fact that the maternal methylation imprint is established after fertilization, suggests a postzygotic epigenetic defect.

The article of Moll et al. indicates an association between IVF and retinoblastoma. As this is the first article reporting such an association and only 5 retinoblastoma cases were described, more research is necessary to confirm this association. Furthermore, if this association is confirmed, the possible causal mechanism has to be explored. This disorder is not associated with an imprinting defect.

Fortunately, the diseases possibly associated with ART are all rare diseases. Therefore, a significant increase in birth prevalence, still results in only a small number of patients born with that disease after ART. Nevertheless, it underlines that procedures like ICSI and IVF were many of the normal steps involved in fertilization and activation of the oocyte are bypassed, should be carefully evaluated with a close follow-up of children conceived after these procedures till adulthood, as in postmarketing surveillance of new medication. Further research is required to determine: (1) the precise relationship between these rare diseases, especially human imprinting disorders, and ART (2) what specific step of ART is responsible for the association, and (3) what precautions can be taken to prevent the occurrence of these syndromes.

F.A.M. Poppelaars, Amsterdam

Maier AG, Duraisingh MT, Reeder JC, Patel SS, Kazura JW, Zimmerman PA, Cowman AF:

Plasmodium falciparum erythrocyte invasion through glycophorin C and selection for Gerbich negativity in human populations

Nat Med 2003;9:87–92

DOI: 10.1159/000073335

Gerbich Negativity in Melanesians through Negative Selection by Malaria

Previously, a strong geographical correlation was found in Melanesians between the occurrence of Gerbich (Ge) blood group system and the coastal areas of Papua New Guinea where malaria is hyperendemic. Ge negativity results from mutations in the glycophorin C gene. In the Melanesian population, an allele from which exon 3 is deleted reaches a frequency of 46.5% in those coastal regions. In this study, the authors provide evidence that the malaria parasite uses glycophorin C to enter human erythrocytes. The data also strongly suggest that a high prevalence of Ge negativity in Melanesian populations has occurred through natural selection by severe malaria.

Parasites that need to enter specific host cells for their life cycle, like malaria parasites, often make use of receptors on the cell surface of these host cells. Parasites express certain proteins (ligands) that specifically bind to these receptors. EBA175 is such a protein that the malaria parasite uses to invade human erythrocytes. The receptor to which EBA175 binds is glycophorin A. It has been postulated that the existence of a family of EBA proteins provides the parasite with a broader spectrum of binding sites at the cell surface of erythrocytes [1]. Maier et al. [2] used several strategies to determine the function of the EBA140 protein, a protein that is very similar to EBA175.

To identify the erythrocyte receptor to which EBA140 can bind, an EBA140-negative malaria strain was generated. The binding capacity of proteins derived from wild-type malaria and proteins derived from EBA140-negative malaria to receptors from erythrocytes was compared. A single erythrocyte-derived receptor was identified that did not bind to the EBA140-negative malaria strain. This protein appeared to be glycophorin C, indicating that EBA140 binds to glycophorin C. To show a direct binding of EBA140 with glycophorin C, erythrocytes from Ge-negative patients were compared with normal controls. Due to the exon 3 deletion in the Ge-negative patients, their erythrocytes express a smaller glycophorin C protein. Indeed, no binding of the EBA140 protein to the smaller glycophorin C was observed in the Ge-negative patients.

The authors also compared the invasion capacity of the EBA140-negative strain with wild-type malaria into human erythrocytes. No difference was found, suggesting that either EBA140 does not participate in the invasion process or the loss of EBA140 is compensated by other invasion pathways. However, parasites were found to be less efficient in invading Ge-negative erythrocytes (~ 61%) compared with normal erythrocytes. In addition, antibodies that bind to EBA140 also reduce the invasive capacity of the parasite because it inhibits the EBA140 function. This reduction in the invasive capacity is not observed when the antibody-treated parasites are used to infect Ge-negative erythrocytes because the EBA140-glycophorin C interaction is not functioning.

Taken together, these data show that the EBA140-glycophorin C interaction plays a role in the invasion process of malaria into human erythrocytes. This binding is disturbed in Ge-negative erythrocytes explaining the observed lower efficiency by which malaria parasites invade these erythrocytes. This defect may be a natural selection process underlying the high frequency of Ge negativity in coastal populations of Papua New Guinea where malaria is endemic.

Comments

There are interesting similarities as well as differences between Ge negativity and sickle-cell anemia. Both result from a genetic alteration and both conditions result in a protective effect against malaria infection. In the heterozygous state for sickle-cell anemia, there are no symptoms or mild anemia and the condition apparently confers resistance to malaria, while persons that are homozygous suffer from deformation of red blood cells causing severe anemia. In subjects with Ge negativity, however, homozygosity is beneficial in case of a malaria infection, whereas heterozygosity is not, and no adverse health effects have been described.

In conclusion, the results broaden our understanding of how the malaria parasite interacts with its host and may result in new targets for drug development.

Q. Waisfisz, Amsterdam

References

- 1 Miller LH, Baruch DI, Marsh K, Doumbo OK: The pathogenic basis of malaria. *Nature* 2002;415:673–679.
- 2 Maier AG, Duraisingh MT, Reeder JC, Patel SS, Kazura JW, Zimmerman PA, Cowman AF: *Plasmodium falciparum* erythrocyte invasion through glycophorin C and selection for Gerbich negativity in human populations. *Nat Med* 2003;9:87–92.

Kowalcsek I, Huber G, Lammers C, Brunk J, Bieniakiewicz I, Gembruch U:

Anxiety scores before and after prenatal testing for congenital anomalies

Arch Gynaecol Obstet 2003;267:126–129

DOI: 10.1159/000073336

Prenatal Examination Influences Anxiety Levels in Pregnant Women

Prenatal diagnosis is increasingly used and studies show that it may have psychological effects on pregnant women. The authors of this article studied the anxiety levels of pregnant women before and after prenatal examination. The research population consisted of a random consecutive sample of 332 German women who underwent prenatal diagnosis between the 12th and 20th week of pregnancy in a tertiary referral centre. The indications for prenatal diagnosis were: suspicious foetal anatomy ($n = 33$), exclusion of malformation ($n = 162$), maternal age (≥ 35 years; $n = 110$) or positive medical history ($n = 27$). The tests the participants underwent were not specified (amniocentesis, ultrasound, karyotyping). Anxiety was measured by the German version of the standardised State-Trait Anxiety Inventory (Spielberger). Trait anxiety was measured before prenatal examination, state anxiety was measured before and after examination.

In 37 women (11.1%), the outcome of the test was positive. No significant difference was found in trait anxiety before prenatal diagnosis in the tertiary referral centre between women who received negative or positive results ($p = 0.410$). There was a difference in state anxiety scores before prenatal diagnosis between both groups of women: women who received positive results scored higher than women who received negative results. The authors suspect that these higher state anxiety scores of women with positive outcomes before the test are due to the fact that in relatively many of these women (40.5%) the indication for prenatal diagnosis was suspected foetal malformations.

After prenatal diagnosis, state anxiety scores also differed significantly between women with positive and negative results. Women

with a positive result had higher scores on the items tension, anxiousness, nervousness, restlessness and fear. The state anxiety scores for women with a negative result were significantly lower after the examination when compared with the score before the examination ($p < 0.001$). The state anxiety scores of women with a positive result did not significantly differ between the pre- and post-examination situation ($p = 0.463$).

Based on the short questionnaire about the content of the anxiety, it seems that pregnant women are just as much afraid of a malformation being detected, as of it being missed.

Comments

The study was performed in a tertiary referral centre, sometimes after suspicion of a foetal anomaly had been raised in another clinic. All types of prenatal diagnosis seem to have been included, although this is not specified. This is a weakness of the study. The impact of 'increased maternal age' probably differs from 'suspicion of foetal anomaly'.

An interesting finding is the fact that women were just as much afraid of a malformation being detected, as of it being missed. One could wonder what women are gaining by doing such a test; if a favourable outcome is given, are they reassured or are they still left with worries? Weinans et al. [1] found that 13% of pregnant women continued to be anxious, even after the favourable result of the amniocentesis. This implicates that it is not only important to guide women who have been given an unfavourable outcome, but that some women who have heard good news might also need special guidance. Kemp et al. [2] found that after consultation by specialist staff, there was a significant reduction in state anxiety scores in parents.

Results in this article show that anxiety is lessened after a favourable test result. But has it returned to normal values? How high is anxiety compared with women who have chosen not to undergo prenatal examination or have not even thought about the possibility of testing? And what are the long-term effects of prenatal diagnosis? This study was only concerned with the immediate effects. The authors recognise this and refer to studies that indicated that even though the examination confirmed the baby's normality, worries can return later on in pregnancy [3]. To gain better insight into the effect of prenatal diagnosis on anxiety, long-term effects should be studied as well.

J.H. Kleinveld, Amsterdam

References

- 1 Weinans MJ, Huijssoon M, Tijmstra T, Gerrits MC, Beekhuis JR, Mantingh A: How women deal with the results of serum screening for Down syndrome in the second trimester of pregnancy. *Prenat Diagn* 2000;20:705–708.
- 2 Kemp J, Davenport M, Pernet A: Antenatally diagnosed surgical anomalies: The psychological effect of parental antenatal counseling. *J Pediatr Surg* 1998;33:1376–1379.
- 3 Hunter M, Tsoi MM, Pearce M, Chudleigh P, Campbell S: Ultrasound scanning in women with raised serum alpha fetoprotein: Long-term psychological effects. *J Psychosom Res* 1987;31:35–39.