Assignment¹ of DNA cytosine-5-methyltransferase 1 (DNMT1) gene to porcine chromosome 2q21→q22 by somatic cell and radiation hybrid panel mapping

N. Bosak,^a S. Fujisaki,^a S. Kiuchi,^a H. Hiraiwa,^b and H. Yasue^a

^a Genome Research Department, National Institute of Agrobiological Sciences, Tsukuba;

^b New Energy and Industrial Technology Developmental Organization, Toshima, Tokyo (Japan)

¹ To our knowledge this is the first time this gene has been mapped in swine.

Rationale and significance

DNA methylation plays an important role in animal development and gene regulation. In mammals, several genes encoding DNA cytosine methyltransferases have been identified. DNA cytosine-5-methyltransferase 1 has a role in the establishment and regulation of tissue-specific patterns of methylated cytosine residues. DNMT1 is constitutively expressed and is required for the maintenance of global methylation after DNA replication. Aberrant methylation patterns are associated with certain human tumors and developmental abnormalities. The human DNMT1 gene was localized on 19p13.3→p13.2 (el-Deiry et al., 1991; Yen et al., 1992). In this study using somatic cell hybrid cells and a radiation hybrid panel we have mapped the gene DNMT1 to porcine chromosome 2, thus confirming the expected localization of the gene in the porcine genome.

Materials and methods

Primer design

The porcine expressed sequences (ESTs), which were generated in our research projects for construction of full-length cDNA libraries from various tissues (in press), were clustered based on sequence similarities and aligned with human sequences assigned to the human physical map (<http://genopole.toulouse.inra.fr/lccare/>). Therefore, from the website, the porcine sequence

orthologous to human gene DNMT1 was selected to design a primer pair inside the putative exon, which was deduced from the exon/intron structure of the corresponding human gene. Primers were designed using Primer 3 software (<http://www.genome.wi.mit.edu/cgi-bin/primer/primer3.cgi>) (Rozen and Skaletsky, 2000).

PCR conditions/sequencing

The PCR conditions were optimized on 25 ng and 35 ng genomic DNAs from pig and Chinese hamster, respectively, in 15 µl of 10 mM Tris-HCl (pH 9.0), 1.5 mM MgCl₂, 50 mM KCl, 200 µM of each dNTP, 1.0 µM of primer pair, and 0.375 U AmpliTaq Gold (BE, Applied Biosystems, USA). The primers used for PCR analysis were 5'-GTATGGCGTGGCCAGAC-3' and 5'-AAACTTCTTATCATCGACCACGA-3'. The mixture was treated at 95 °C for 9 min, then with 35 cycles of denaturation at 95 °C for 30 sec, 65 °C for 30 s, and extension at 72 °C for 30 s, followed by a final elongation step at 72 °C for 5 min. All PCRs were carried out using the GeneAmpPCR System 9700 (Perkin-Elmer, CT, USA). The 142-bp porcine-specific PCR product was observed on agarose gel electrophoresis. The resultant PCR product was sequenced using ABI PRISM Big Dye Terminator Sequencing Ready Reaction Kit and a 3700 DNA Analyzer (Applied Biosystems, CA, USA) to confirm that the sequence was identical to the sequence used for primers design.

RH panel mapping

Using the primer pair thus prepared as well as the 90 selected IMPRH 7000-radiation hybrid cell panel DNAs, which were provided by INRA (France) and the University of Minnesota (USA) (Yerle et al., 1998), PCR was performed under the conditions described above. PCR products were separated in a 2.0% agarose gel, and scored for mapping as described previously (Kiuchi et al., 2002). Assignment of the gene to the IMPRH map was performed with the IMPRH mapping tool (Milan et al., 2000), which is accessible on the INRA web server (<http://imprh.toulouse.inra.fr>).

SCH mapping

The somatic cell hybrid panel DNAs (Yerle et al., 1996) provided by INRA (France) were used for mapping DNMT1. The PCR conditions for SCH mapping were the same as for RH mapping except that 20 ng of panel DNA was used for each PCR. The assignment was performed with software that calculates the likelihood of the localization of a marker in 115 individual regions of the porcine genome on the INRA web server (<http://www.toulouse.inra.fr/lgc/pig/pcr/pcr.htm>).

Supported by a grant from the Organized Research Combination (ORCS) program.

Received 24 April 2003; manuscript accepted 29 June 2003

Corresponding author: Hiroshi Yasue, Genome Research Department

National Institute of Agrobiological Sciences

Tsukuba, Ibaraki 305-0901 (Japan)

telephone: +81-298-38-8664; fax: +81-298-38-8674; e-mail: hyasue@affrc.go.jp

Results

Regional mapping results

Somatic porcine × rodent cell hybrid panel analysis of 27 hybrids resulted in the following vector: 01000 10000 00000 00000 10001 00. Statistical evaluation showed a correlation between the DNMT1 gene and chromosome 2 (1/2 q21) → (1/2 q22) with the highest probability and correlation values (69 and 85%, respectively; error risk < 1%). However, in repeated experiments, the positive signal was observed additionally in the hybrid N23 (01000 10000 00000 00000 10101 00), which resulted in the correlation between DNMT1 gene and chromosome 2p17→p14 with the highest probability and correlation values (79 and 87%, respectively; error risk < 0.5%). The IMpRH panel was employed to determine the precise location of DNMT1. The survey of the RH panel showed the following distribution within the 90 lines: 00010 00010 01101 11100 00101 11000 00000 00000 01000 00100 10100 01000 01000 01100 10111 01011 00000 11001, with the most significantly linked markers (LOD score 9.48 and 9.34) SWR1342 and SSC2F05 (Hawken et al., 1999) located on SSC2q21→q24. This supported the result of the former SCH mapping rather than the latter. Localization of the DNMT1 locus to SSC2q21→q22 using both SCH and IMpRH panels, confirmed the conservation of synteny between the SSC2 and HSA19.

References

- el-Deiry WS, Nelkin BD, Celano P, Yen RW, Falco JP, Hamilton SR, Baylin SB: High expression of the DNA methyltransferase gene characterizes human neoplastic cells and progression stages of colon cancer. *Proc natl Acad Sci, USA* 88:3470–3474 (1991).
- Hawken RJ, Murtaugh J, Flickinger GH, Yerle M, Robic A, Milan D, Gellin J, Beattie CW, Schook LB, Alexander LJ: A first-generation porcine whole-genome radiation hybrid map. *Mammal Genome* 10:824–830 (1999).
- Kiuchi S, Inage Y, Hiraiwa H, Uenishi H, Yasue H: Assignment of 280 swine genomic inserts including 31 microsatellites from BAC clones to the swine RH map (IMpRH). *Mammal Genome* 13:80–88 (2002).
- Milan D, Hawken R, Cabau C, Leroux S, Genet C, Lahbib Y, Tosser G, Robic A, Hatay F, Alexander L, Beattie C, Schook L, Yerle M, Gellin J: IMpRH server: an RH mapping server available on the Web. *Bioinformatics* 16:558–559 (2000).
- Rozen S, Skaletsky H: Primer 3 on the WWW for general users and for biologist programmers. *Meth molec Biol* 132:365–386 (2000).
- Yen RW, Vertino PM, Nelkin BD, Yu JJ, el-Deiry W, Cumaraswamy A, Lennon GG, Trask BJ, Celano P, Baylin SB: Isolation and characterization of the cDNA encoding human DNA methyltransferase. *Nucl Acids Res* 20:2287–2291 (1992).
- Yerle M, Echard G, Robic A, Mairal A, Dubut-Fontana C, Riquet J, Pinton P, Milan D, Lahbib-Mansais Y, Gellin J: A somatic cell hybrid panel for pig regional gene mapping characterized by molecular cytogenetics. *Cytogenet Cell Genet* 73:194–202 (1996).
- Yerle M, Pinton P, Robic A, Alfonso A, Palvadeau Y, Delcros C, Hawken R, Alexander L, Beattie C, Schook L, et al: Construction of a whole-genome radiation hybrid panel for high-resolution gene mapping in pigs. *Cytogenet Cell Genet* 82:182–188 (1998).