Preface

This assortment of papers on vertebrate sex determination is a small, but representative sample from the more than 100 papers presented at the Third International Symposium on Vertebrate Sex Determination, held in Kona, Hawaii, March 24–28, 2003. The emphasis of these symposia, held once every three years, has from the outset been on the evolution of sex determination among the vertebrates. This appears to be a topic of considerable interest, as the number of participants (from more than 15 countries) has increased at each successive symposium. We feel that it is not only the location that draws such a group, but the single session, single topic and small informal setting with intense and active discussions that makes these meetings so special.

The number of genes now shown to be involved in sex determination and gonad differentiation has increased dramatically since the last symposium. There were a large number of papers and posters presenting evidence for yet more sex-specific genes expressed in the developing gonads. Specific roles for many of these genes have yet to be assigned.

It is now more than 13 years since the discovery of the testis-determining gene on the Y chromosome of mouse and human (Koopman et al., 1990; Sinclair et al., 1990), a discovery that reinvigorated the relatively inactive field of sex determination and led to the discovery of whole families of genes important in vertebrate development. This gene known as SRY, human, *Sry*,

mouse (Sex determining region of the Y chromosome), however, continues to baffle researchers as to its mechanism of action. It is known that expression of this gene at a critical stage in development triggers the cascade of gene expression that leads to development of a testis, but despite an enormous amount of work, exactly how it initiates this extraordinary developmental wonder is not fully understood. Progress is being made, albeit slowly (see recent review by Harley et al. (2003) and papers by Lau in this issue).

After nine years and three symposia, patterns are beginning to emerge; a common suite of genes are found to be expressed in the developing gonads of teleost fish, amphibians, reptiles, birds and mammals. There are however, notable differences among the different classes of vertebrates in the timing and sequence of expression of some of these genes. Thus Sox-9 precedes and initiates expression of MIH (Mullerian Inhibiting Hormone, a peptide hormone secreted by the Sertoli cell) in mammals, but in birds and reptiles, MIH expression precedes that of Sox-9. Another gene in the sex-determining pathway that continues to confound is DAX1 (NR0B1). A number of papers were presented showing its essential role in normal testis development, yet when this gene is duplicated in a mouse an XY female develops. It also plays an important role in steroidogenesis in adrenal and gonad tissue.

Group photo of participants outside the King Kamehameha Hotel, Kona Hawaii at the close of the symposium.

One of the more fascinating stories to emerge from this evolutionary saga is that of the gene DMRT1. This gene, located on chromosome 9 of humans, has homologues in the sex-determining pathway of fruit flies and nematodes, (Raymond et al. 1998) and is expressed in a male specific pattern in all vertebrates studied. In birds DMRT1 is located on the Z sex chromosome and is absent from the W (Nanda et al., 1999). The medaka (*Oryzias latipes*), a teleost fish with an XX:XY chromosome mechanism of sex determination has two copies

of a DMRT1-like gene, *dmrt1a* and *dmrt1b*(Y). In this species the *dmrt1b*(Y) is located on the Y chromosome and has been identified as the sex-determining gene (Matsuda et al., 2002), but in other teleost species the gene is absent (Kondo et al., 2003).

We may expect more such intriguing conundrums before the next symposium.

Valentine A. Lance and Mark H. Bogart

References

- Harley VR, Clarkson MJ, Argentaro A: The molecular action and regulation of the testis determining factors, SRY (sex determining region on the Y chromosome) and SOX9 (Sry-related high-mobility group [HMG] box9). *Endocrin Rev* 24:466–487 (2003).
- Kondo M, Nanda I, Hornung U, Asakawa S, Shimizu N, Mitani H, Schmid M, Shima A, Shartl M: Absence of the candidate male sex-determining gene *dmrt1b*(Y) for medaka from other fish species. *Curr Biol* 13:416–420 (2003).
- Koopman P: Expression of a candidate sex-determining gene during mouse testis differentiation. *Nature* 348:450–452 (1990).
- Lau YFC, Lau HW, Kömüves LG: Expression pattern of a gonadoblastoma candidate gene suggests a role of the Y chromosome in prostate cancer. *Cytogenet Genome Res* 101:250–260 (2003).
- Li Y, Yue L, Taketo T, Lau YFC: Protein transduction as a strategy for evaluating important factors in mammalian sex determination and differentiation. *Cytogenet Genome Res* 101:237–241 (2003).
- Matsuda M, Nagahama Y, Shinomiya A, Sato T, Matsuda C, Kobayashi T, Morrey CE, Shibata N, Asakawa S, Shimizu N, Hori H, Hamaguchi S, Sakai-zumi M: DMY is a Y-specific DM-domain gene required for male development in the medaka fish. *Nature* 417:559–563 (2002).
- Nanda I, Shan Z, Scharl M, Burt DW, Koehler M, Nothwang H, Grutzner F, Paton IR, Windsor D, Dunn I, Engel W, Staeheli P, Mizuno S, Haaf T, Schmid M: 300 million years of conserved synteny between chicken Z and human chromosome nine. *Nature Genet* 21:258–259 (1999).
- Raymond CS, Shamu CE, Shen MM., Seifert KJ, Hirsch B, Hodgkin J, Zarkower D: Evidence for evolutionary conservation of sex-determining genes. *Nature* 391:691–695 (1998).
- Sinclair AH, Berta P, Palmer MS, Hawkins JR, Griffiths BL, Smith MJ, Foster JW, Firschauf AM, Lovell-Badge R, Goodfellow PN: A gene from the human sex determining region encodes a protein with a conserved DNA-binding motif. *Nature* 346:240–244 (1990).