Introduction

Ian G. McKeith

Department of Old Age Psychiatry, Institute for Ageing and Health, Wolfson Research Centre,
Newcastle General Hospital, Newcastle-upon-Tyne, UK

Lewy body disease is thought to be present in up to 20% of elderly patients with dementia, making dementia with Lewy bodies (DLB) the second most common form of degenerative dementia after Alzheimer's disease (AD). Cerebrovascular disease also may contribute to both disorders. Given the proportion of patients suffering from DLB, a group of international experts was convened to discuss the etiology, definition, and diagnosis of this condition, and to consider currently available treatment options.

Although the cognitive deterioration observed in AD is relatively stable over short periods, the cognitive impairments of DLB can fluctuate markedly, as do attention and alertness. Spontaneous motor/extrapyramidal features similar to those seen in Parkinson's disease develop in up to 75% of DLB patients, but the diagnosis can, and should, also be made in their absence. Recurrent visual hallucinations are another common core manifestation of DLB that may alert the clinician to the possible diagnosis and often necessitate pharmacologic treatment. Sensitivity to neuroleptics also can act as a signpost to a DLB diagnosis and is associated with significantly increased morbidity and mortality. Other features supportive of DLB are recurrent falls and syncope, transient losses of consciousness, delusions, depression, and REM sleep disorder.

Consensus clinical diagnostic criteria for DLB have been validated against neuropathologic autopsy findings. Mega and colleagues [1996] and McKeith and others

[2000] found sensitivity for a clinical diagnosis of DLB at 75 and 79% versus specificity at 83 and 95%, respectively – levels of diagnostic accuracy that are similar to those for AD and vascular dementia. False-positive diagnoses of DLB do not appear to be as frequent as false negatives, exemplified in the recent data of Lopez and coworkers [2002] reporting sensitivity and specificity of 30.7 and 100%, respectively.

Neuroimaging was reviewed as a potential diagnostic method to differentiate AD from DLB. Although structural MRI shows relative preservation of medial temporal lobe structures in DLB compared with AD, this is of limited specificity in individual case diagnosis. The use of positron emission tomography (PET) and single-photon emission [computed] tomography (SPET, SPECT) to identify occipital hypoperfusion can help differentiate DLB from AD. SPECT visualization of the striatal dopamine transporter with β -CIT or related ligands may provide a more powerful discriminator, but confirmatory studies are needed.

Severe cholinergic deficits present in DLB are the focus of pharmacologic treatment goals. Acetylcholinesterase inhibitors (AChEIs) reduce the degradation of acetylcholine and preserve neuronal function in the temporal, parietal, and midfrontal cortex, the targets of more extensive cholinergic depletion in DLB compared with AD. Safety and efficacy of the AChEIs currently available were presented and discussed at the meeting, and the following articles reflect the deliberations of those present.

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2004 S. Karger AG, Basel
1420–8008/04/0175–0001\$21.00/0

Accessible online at:
www.karger.com/dem

Ian G. McKeith MD
Department of Old Age Psychiatry, Institute for Ageing and Health
Wolfson Research Centre, Newcastle General Hospital, Westgate Road
Newcastle-upon-Tyne, NE4 6B3 (UK)
Tel. +44 191 256 3018, Fax +44 191 219 5071, E-Mail maureen.middlemist@ncl.ac.uk

Suggested Reading

- Chui HC, Victoroff JI, Margolin D, Jagust W, Shankle R, Katzman R: Criteria for the diagnosis of ischemic vascular dementia proposed by the State of California Alzheimer's Disease Diagnostic and Treatment Centers. *Neurology* 1992;42:473–480.
- Doody RS, Stevens JC, Beck C, et al: Practice parameter: management of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001;56:1154–1166.
- Ishii K, Imamura T, Sasaki M, et al: Regional cerebral glucose metabolism in dementia with Lewy bodies and Alzheimer's disease. *Neurology* 1998;51:125–130.
- Ishii K, Yamaji S, Kitagaki H, Imamura T, Hirono N, Mori E: Regional cerebral blood flow difference between dementia with Lewy bodies and AD. *Neurology* 1999;53:413–416.
- McKeith I, Fairbairn A, Perry R, Thompson P, Perry E: Neuroleptic sensitivity in patients with senile dementia of Lewy body type. *BMJ* 1992;305:673–678.
- McKeith IG, Ballard CG, Perry RH, et al: Prospective validation of consensus criteria for the diagnosis of dementia with Lewy bodies. *Neurology* 2000;54:1050–1058.
- McKeith IG, Galasko D, Kosaka K, et al: Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB): report of the consortium on DLB international workshop. *Neurology* 1996;47:1113–1124.
- McKeith IG, O'Brien JT, Ballard C: Diagnosing dementia with Lewy bodies. *Lancet* 1999;354:1227–1228.
- McKeith IG, Perry EK, Perry RH: Report of the second dementia with Lewy body international workshop: diagnosis and treatment. *Neurology* 1999;53:902–905.
- McKeith IG: Dementia with Lewy bodies. *Br J Psychiatry* 2002;180:144–147.
- McKeith IG: Lewy body disease. *Rev Neurol (Paris)* 2001;157(suppl 10):28.
- Mega MS, Cummings JL, Fiorello T, Gornbein J: The spectrum of behavioral changes in Alzheimer's disease. *Neurology* 1996;46:130–135.
- Okamura N, Arai H, Higuchi M, et al: [¹⁸F]FDG-PET study in dementia with Lewy bodies and Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 2001;25:447–456.