- Struhsaker TT (1981). Census methods for estimating densities. In *Techniques for the Study of Primate Population Ecology* (Subcommittee on Conservation of Natural Populations, National Research Council, ed.), pp 36-80. Washington, National Academy Press.
- Thorington RW, Ruiz, JC, Eisenberg JF (1984). A study of a black howling monkey (*Alouatta caraya*) population in Northern Argentina. *American Journal of Primatology* 6: 357–366.
- Zunino GE, Bravo S, Ferreira FM, Reisenman C (1996). Characteristics of two types of habitat and the status of the howler monkey (*Alouatta caraya*) in northern Argentina. *Neotropical Primates* 4: 48–50.
- Zunino GE, González V, Kowalewski MM, Bravo SP (2001). *Alouatta caraya*: Relations among habitat, density and social organization. *Primate Report* 61: 37–46.

Erratum

In the article ‘Seed Dispersal by Sympatric Tamarins, *Saguinus mystax* and *Saguinus fuscicollis*: Diversity and Characteristics of Plant Species’, by Christoph Knogge and Eckhard W. Heymann, published in *Folia Primatologica* 2003;74:33–47, figure 3 was not published correctly. The solid curve should have been labelled ‘*Saguinus fuscicollis*’ and the dotted one ‘*Saguinus mystax*’. Corrected figure see below.

Fig. 3. Rarefaction curves for the relationship between the number of seed dispersal events and the number of seed species encountered in faecal samples.