Introduction

The importance of chromosome aberrations for evolution and their association with human health have been recognized for almost a century. However, the mechanisms involved in the formation of chromosome aberrations are still not well understood. Ionizing radiation is most efficient in inducing chromosome aberrations and the basic principles of aberration formation were laid out in the early 1930's, even though the molecular structure of the eukaryotic chromosome was not known at that time. Techniques to prepare and stain chromosomes have improved gradually in the last decades, parallel with increased ability to identify and quantify chromosome aberrations, thus leading to a better understanding of their origin. Use of colchicine, squash technique for plant chromosomes, hypotonic shock for mammalian chromosomes, various banding techniques in the 1960's, sister chromatid differentiation in the early 1970's and fluorescence in situ hybridization (FISH) in the late 1980's have been main break-throughs enabling us to study chromosome aberrations in great detail. Especially the increasing availability of whole chromosome-specific, arm-specific, region-specific, centromere- and telomere-specific DNA probes has revolutionized the ways to look at chromosomes and their aberrations.

The articles in this volume deal with chromosome structure, mechanisms of aberration formation and applied aspects of aberration analyses. Papers which were presented at the 6th International Symposium on Chromosome Aberrations held at the University of Duisburg-Essen, Germany (10th to 13th September, 2003) and a collection of invited papers in this field are included.

The first group of 14 papers deal with **Basic Aspects** of chromosome aberrations, such as DNA damage processing in relation to aberration formation, the importance of non-homologous end joining, homologous recombination and the nature of aberrations.

The role of telomeres and telomerase in maintaining the integrity of chromosomes is an important area of research and five papers under **Telomeres** deal with this area.

Chromosome aberrations induced by ionizing radiation have been studied in great detail over the years and data generated in the 1930's and 1940's were used as a quantitative basis for formulating theories to explain the biological action of radiation. Knowledge of types and frequencies of lesions induced by ionizing radiation, especially DNA double strand breaks, as well as the quantification of aberrations have increased in the last decade and it has been possible to generate biophysical models for the origin of radiation-induced chromosome aberrations. Three papers are presented under **Modelling**.

Papers dealing with ionizing radiation-induced chromosome aberrations are presented in two groups, namely **Low-LET-Radiation** (eight papers) and **High-LET-Radiation** (nine papers). There are qualitative and quantitative differences between chromosome aberrations induced by low LET (such as X-rays and γ -rays) and high LET radiation (such as neutrons, heavy ions and radon). High LET radiation induces high frequencies of complex chromosome exchanges and FISH studies have shed light on the origin of these types of aberrations.

A fraction of chromosome aberrations induced in germ cells are transmitted to the next generation and this is of great concern from the point of view of genetic risk. Four papers included under **Heritable Effects** deal with this aspect.

Numerical and structural aberrations are important both in congenital abnormalities and tumors. Studying micronuclei using centromere-specific probes can easily make assessment of aneuploidy as well as chromosome breaks. Four papers are presented in the section **Aneuploidy and Micronuclei**.

Sister chromatid exchanges (SCEs) are intrachromosomal exchanges probably reflecting a homologous recombination process. Three papers dealing with mechanisms of formation of SCEs are presented under **Sister Chromatid Exchanges**.

Cells derived from patients suffering from recessive diseases (such as ataxia telangiectasia, Fanconi anemia, Bloom's syndrome) are chromosomally unstable. The genetic basis of these diseases is understood now, thanks to the molecular biological techniques, which have become available. Specific chromosome translocations or deletions characterize many human cancers. Chromosome aberrations in lymphocytes of individuals can be used as a biomarker of cancer risk. Twelve papers in this area of research are presented under the heading **Applied Aspects/Cancer**.

The papers in this special issue show the importance of basic and applied aspects of studying chromosome aberrations. It becomes clear that this field of research is of great interest and by far not out of the scope of modern science. Molecular cytogenetics is an especially rapidly expanding field and has contributed to our knowledge of chromosome structure and interphase nuclei, as well as the formation of chromosome aberrations. In the area of clinical genetics and oncology, these techniques have been of great value in diagnosis and prevention of outcome of diseases. We strongly believe that the future for this area is very bright.

We would like to thank Mrs. Rita Nadorf for her unfailing and devoted assistance in organizing the symposium as well as in editing of this volume.

G. Obe

A.T. Natarajan