Foreword

Paul Enck

Department of General Surgery, University Hospitals Tübingen, Tübingen, Germany

Incontinence, both urinary and fecal, is a devastating condition for all affected: it is *not* a symptom of only the elderly but is found secondary to a variety of systemic medical diseases and clinical conditions in women and men, is seen with pelvic floor trauma, after childbirth as well as following pelvic floor surgery, in association with local inflammatory disorders of the lower gastrointestinal and urinary tract, and in many cases where a specific mechanism for symptom occurrence cannot be identified. It significantly decreases the quality of life in all [1]. Despite its high prevalence [2], the pathophysiology of incontinence is still incompletely understood, and part of the reason for this is that within the medical system, incontinence is frequently ignored or neglected by doctors [3], and patients still tend to hide their incontinence symptoms due to embarrassment and shame [4].

The European Community's 5th Framework Program (1998–2002) 'Quality of life and management of living resources' identified 'The Aging Population' as a key action, and asked for research proposal on 'Age-related illnesses and health problems'; among these 'incontinence' was specifically nominated in a call in 2001.

At this time the coordinator of the OASIS project (P.E.) had developed a specific research hypothesis of the role of asymmetric sphincter innervation in fecal incontinence, which was supported by a national German grant (DFG En 50/21) – the evidence for this is discussed and illustrated in one of the chapters of this issue [5]. Based on

these data, and with the help of another European Community sponsored project (SENIAM) which had developed a new surface EMG technology [6], it was possible to apply a 16-channel multi-electrode array technique to record motor unit action potentials from muscle fibers of the external anal sphincter for the first time – the starting point of the OASIS project.

It marked the beginning of a fruitful collaboration between four medical subspecialties (gastroenterology, surgery, gynecology, and urology), a medical engineering department, and two small companies involved in development and marketing of technology for incontinence diagnosis and therapy, which form the OASIS consortium. The submitted research proposal was approved and received European funding of approx. EUR 1.7 million for a period of 3 years, the project started January 2002.

The main objective of the project is to gather knowledge about pelvic floor asymmetry and its role in development, severity, diagnosis, treatment, and prevention of incontinence in the elderly. This goal will be achieved by the following steps: (a) quantification of asymmetry in healthy subjects and in major patient subgroups with incontinence; (b) to correlate asymmetry (by type and degree) to age and age-related changes in pelvic floor functions as assessed by functional diagnostic testing; (c) to identify risk groups predisposed to become incontinent at higher age; (d) to provide a scientific basis to treat inconti-

KARGER

Fax + 41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2004 S. Karger AG, Basel
0012-2823/04/0692-0084\$21.00/0

Accessible online at:
www.karger.com/dig

Prof. Dr. Paul Enck, Coordinator
Department of General Surgery, University Hospitals Tübingen
Waldhörnlestrasse 22
DE-72072 Tübingen (Germany)
E-Mail paul.enck@uni-tuebingen.de

nence by applying S-EMG to routine diagnosis and therapy systems; (e) to develop a clinical strategy to deal with incontinence especially in the elderly.

The project consists of 14 work packages, of which 2 are administrative, 6 are clinical, and 6 technical in nature. It proposed to develop prototype hardware and software tools, especially anal and vaginal/urethral EMG probes, which allow non-invasive recording of individual motor units from each side of the pelvic floor muscles and to identify individuals with significant asymmetric sphincter muscle innervation. Subsequently, these tools will be applied to measure pelvic floor innervation and its symmetry/asymmetry in subgroups of patients with urinary and/or fecal incontinence, as identified in major medical subspecialties (urology, gynecology, gastroenterology, surgery) in different countries in Europe. Furthermore, probe and system validation will be performed in patients in whom selective stimulation of left and right dorsal sacral roots is possible after sacral root stimulator implantation. Finally, in women undergoing childbirth, the risk of developing incontinence after pelvic floor trauma may be assessed and related to the existence and degree of asymmetric sphincter innervation.

The papers in this issue of *Digestion* mark the 'half time' of the project; consequently, they reflect the scientific 'roots' of the project – the paper by Vodusek [7] on 'Neurocontrol of the pelvic floor', the paper on current concepts of understanding urinary incontinence by Heesakkers and Gerretsen [8], and the paper on collected evidence for asymmetric sphincter innervation in the pathogenesis of fecal incontinence by Enck et al. [5] – as well as first results of OASIS: the application of multi-electrode array surface EMG to recordings from the anal and urethral sphincter in theory [9] and in practice [10].

We cannot yet answer the question whether and what role asymmetry of innervation plays in the pathogenesis of incontinence, but recordings in patients with our new technology are in progress, and by the end of 2004 a collective database from more than 1,000 patients will hopefully allow a sufficient answer. Already, however, the OASIS project has changed our way of thinking of neural control of the pelvic floor, as patterns of innervation in health and disease, between gender, and across the age range urge us to overcome traditional thinking: new technology allows assessment of sphincter functions and their relationship to clinical findings in much greater detail than ever before. Future investigations will ask for the effects of short- and long-term exercise, of muscle fatigue, but also of surgery, and of drug treatment on these sphincter functions – with or without asymmetry.

OASIS Participants

Prof. Dr. Paul Enck – Coordinator
University Hospitals Tübingen
Department of General Surgery
Waldhörnlestrasse 22
DE-72072 Tübingen (Germany)
E-Mail paul.enck@uni-tuebingen.de

Dr. John Heesakkers
Department of Urology, UMC Nijmegen
PO Box 9101
NL-6500 HB Nijmegen (The Netherlands)
E-Mail j.heesakkers@uro.azn.nl

Priv.-Doz. Dr. Heiko Franz
Department of Gynaecology
Vivantes Hospitals Berlin
Mariendorf Weg 28
DE-12051 Berlin (Germany)
E-Mail heiko.franz@vivantes.de

Dr. Fernando Azpiroz
Digestive System Research Unit
University Hospital Vall d'Hebron
ES-08035 Barcelona (Spain)
E-Mail fernando.azpiroz@wol.es

Prof. Dr. Roberto Merletti
COREP-Politecnico di Torino
Department of Electronics
Corso Duca degli Abruzzi, 24
IT-10129 Torino (Italy)
E-Mail merletti@polito.it

Andreas Schönfeld
Standard Instruments GmbH
Im Schleht 26
DE-76187 Karlsruhe (Germany)
E-Mail standard-instruments@t-online.de

Hans Hofman
Fornix Medical Systems b.v.
Schoenaker 10e, PO Box 18
NL-6640 AA Beuningen (The Netherlands)
E-Mail Hans.Hofman@Fornixmedicalsyste.ms.com

References

- 1 Miner PB: Economic and personal impact of fecal and urinary incontinence. *Gastroenterology* 2004;126:S8–S13.
- 2 Nelson RL: Epidemiology of fecal incontinence. *Gastroenterology* 2004;126:S3–S7.
- 3 Enck P, Gabor S, v. Ferber L, Rathmann W, Erckenbrecht JF: Prevalence of fecal incontinence and degree of information in family doctors and health maintenance organisation. *Z Gastroenterol* 1991;29:538–540.
- 4 Leigh RJ, Turnberg LA: Faecal incontinence: The unvoiced symptom. *Lancet* 1982;i:1349–1351.
- 5 Enck P, Hinrichhofen H, Wietek B, Becker HD: Functional asymmetry of sphincter innervation and its role in the pathogenesis of fecal incontinence. *Digestion*, in press.
- 6 Hermens HJ, Freriks B: The state of the art on sensors and sensor placement procedures for surface electromyography: A proposal for sensor placement procedures. Deliverable of the SENIAM Project 1997, Roessingh Research and Development (Pub).
- 7 Vodusek DB: Anatomy and neurocontrol of the pelvic floor. *Digestion*, in press.
- 8 Heesakkers JPFA, Gerretsen RR: Urinary incontinence. Sphincter functioning from a urologic perspective. *Digestion*, in press.
- 9 Merletti R, Bottin A, Cescon C, Farina D, Gazuzoni M, Martina S, Mesin L, Pozzo M, Rainoldi A, Enck P: Multi-channel surface EMG for the non-invasive assessment of the anal sphincter muscle. *Digestion*, in press.
- 10 Enck P, Franz H, Azpiroz F, Farga J, Hinrichhofen H, Kaske-Bretag K, Bottin A, Martina S, Merletti R: Innervation zones of the external anal sphincter in healthy male and female subjects: Preliminary results. *Digestion*, in press.