Patrinos GP, Giardine B, Riemer C, Miller W, Chui DHK, Anagnou NP, Wajcman H, Hardison RC:

Improvements in the *HbVar* database of human hemoglobin variants and thalassemia mutations for population and sequence variation studies.

Nucleic Acids Res 2004;32:D537–D541.

Tadmouri GO, Gulen RI:

Deniz: The electronic database for β -thalassemia mutations in the Arab world.

Saudi Med J 2003;24:1192–1198.

DOI: 10.1159/000078509

The Use of Genetic Databases in Population Screening Programs for Hemoglobinopathies

HbVar (<http://globin.cse.psu.edu/globin/hbvar/>) is a relational database developed by multicenter academic effort to provide information on the genomic sequence changes leading to hemoglobin variants and all types of hemoglobinopathies. The database stores data from previous studies [1, 2], but also allows regular updates and corrections following from new discoveries. Published information on pathology, hematology, clinical presentation, laboratory findings, biochemical data and techniques to identify variants are included in the database. All these data are publicly available, and are of particular interest to the research community and physicians as an aid in diagnosis. Recently, the possibilities of this 4-year-old database have been expanded with two significant advances:

(1) On request of the current users, the frequencies of β -thalassemia mutations in different countries/populations have been added. The data are based on the frequency spectrum in 48 ethnic subgroups for 121 β -thalassemia mutations reported in the published literature. This results in the possibility to query the most frequently occurring mutations in a specific population.

(2) The second improvement is the linkage of *HbVar* to the Genome Alignment and Annotation database (GALA; available at <http://globin.cse.psu.edu/gala/>). GALA is a database which provides information about features of genomic DNA, such as gene structures, repetitive elements and many others. An example of this new capacity is the ability to generate details about the position and size of a mutation on the gene. Recent developments in the assembly of the human reference sequence will lead to much more information on human sequence variants; thus, improving the database is an ongoing process.

Many scientific papers deal with the distribution of β -thalassemia mutations in Arab populations. As a consequence, knowledge of this subject has been widely scattered, and access to the information is beyond the capacity of the individual investigator. Thus, Deniz (<http://biobase.fatih.edu.tr/>), an integrating electronic database, was established to prevent references to outdated data and to condense all information into clear trends and facts. The database currently includes data on β -thalassemia mutations from 14 Arab countries, but the design and structure allow the addition of future data. According to the Deniz database, six mutations are responsible for 66.2% of the Arab β -thalassemia chromosomes. Other applications are the prediction of diagnostic strategies that may be offered to natives of unstudied countries and provision of insights into the origins of β -thalassemia in the Arab world. Integration of Deniz with other databases is currently in process.

Comments

Both the *HbVar* and Deniz databases try to make (integrated) genetic information available for health care workers. *HbVar* recently expanded with frequency distribution data, and Deniz focuses on this kind of data by definition. As mentioned by Patrinos et al., such information significantly simplifies mutation analysis. Tadmouri and Gulen also mention the application of Deniz in the prediction of diagnostic strategies. Another potential field of use would be in effectuating genetic screening programs for hemoglobinopathies. Hemoglobinopathies are recessive inherited disorders. Screening programs aim to detect carrier couples of a mutation in the hemoglobin genes to inform them about the risk of having an affected child. Such programs are currently utilized in some countries with a high prevalence of hemoglobinopathies (for example Italy, Greece and Cyprus) [3]. To assess the properties of a database for use in clinical and population screening applications, it is important to evaluate database design and content, sources of data and accessibility [4].

Both databases deal with frequency distribution data of β -thalassemia mutations. The available information may be useful in a screening program, because it provides the ability to start genetic screening of most frequently occurring mutations in the relevant ethnic background. This is particularly valuable in countries with many different ethnic groups at risk. However, screening of β -thalassemia mutations is not enough to inform an individual about the risk of an affected newborn. Other mutations in the hemoglobin genes are also able to induce a major β -thalassemia phenotype. An example is the compound heterozygote combination of HbE and a β -thalassemia mutation, which is very common in Southeast Asia. Consequently, a screening program must include many variants of hemoglobinopathies, and therefore a database with distribution frequency data for all types of hemoglobinopathy mutations would be more helpful.

Other useful data would be the phenotype which belongs to a specific genotype. This information is needed to fully inform the individual undergoing screening about the severity of the disease. *HbVar* tries to include these data (unfortunately, knowledge about the geno-

type-phenotype correlation is not always that clear). Tadmouri and Gulen do not mention this feature of the Deniz database explicitly.

The World Wide Web has made it possible to provide both databases to a widespread community, so accessibility is not a problem. However, with the existence of databases containing more or less the same information, the user does not always know *which one* to access. The user can compare databases according to his/her own needs, and it may be possible that competition between databases could lead to optimization of their quality. On the other hand, it takes a lot of effort for the potential user to assess the databases in the correct way. Besides, the quality of a database may change over time, depending on the frequency of updating. All similar databases have to be updated with the same new information. This means, therefore, a lot of construction work. In conclusion, the existence of two more or less similar databases is not efficient. However, with the existence of only one database, the user must totally rely on that specific one. To guarantee quality of 'The hemoglobinopathy database', an option is to ask for financial contributions, which can be used to increase quality control. Both publications discussed herein mention intentions to work on greater integration of databases, so perhaps setting up one comprehensive database is possible?

R. Achterbergh, Amsterdam

References

- 1 Huisman THJ, Carver MFH, Efremov GD: A syllabus of human hemoglobin variants, ed 2. Augusta, The Sickle Cell Anemia Foundation, 1998.
- 2 Huisman THJ, Carver MFH, Baysal E: A syllabus of thalassemia mutations. Augusta, The Sickle Cell Anemia Foundation, 1997.
- 3 Modell B: Delivering genetic screening to the community. *Ann Med* 1997; 29:591-599.
- 4 Birch P, Friedman JM: Utility and limitations of genetic disease databases in clinical genetics research: A neurofibromatosis 1 database example. *Am J Med Genet* 2004;125C:42-49.

Michie S, Smith JA, Senior V, Marteau TM:

Understanding why negative genetic test results sometimes fail to reassure.

Am J Med Genet 2003;119A:340-347.

DOI: 10.1159/000078510

Understanding Why Negative Genetic Test Results Sometimes Fail to Reassure

Predictive genetic testing enables people at risk for inherited conditions to learn about their chance of developing the condition. Where early intervention is effective, clinical surveillance or treatment can be implemented. Some of the first studies, however, suggested that there might be problems in compliance with surveillance amongst those individuals with positive test results, whereas others

reported problems of the desire for unwarranted clinical surveillance amongst those individuals receiving negative test results. In the paper of Michie et al., it is discussed why some people apparently are not reassured by receiving a negative genetic test result.

In an earlier quantitative study of 127 people receiving negative test results following familial adenomatous polyposis testing, it was found that 42% expected to attend future bowel cancer screening [1]. The main predictor of wanting to continue to attend for bowel screening was doubt about the accuracy of the genetic test. There is, however, little research investigating how people think about screening and diagnostic tests and what psychological impact they may have. The aim of the study of Michie et al. was to gain greater understanding of how people represent the genetic test and the genetic basis of the condition in order to explain why some desire to continue with clinically unnecessary screening. The study employed a quantitative method, 'interpretative phenomenological analyses', which has been used to study a wide range of topics concerning psychological aspects of new genetics. Nine patients who received negative results following predictive testing for familial adenomatous polyposis within the previous year at a London hospital were interviewed. Five of them felt no need for future bowel or fecal screening; four expressed a desire for future screening.

The results of the study show that the reasons for not feeling reassured by the negative genetic test results could be divided into two main categories: (1) *perceptions of the genetic basis of the condition*: the dominantly inherited condition could be caused by a new spontaneously arising mutation, or caused by other factors in addition to genes; (2) *perceptions of the genetic test*: a blood test was either not represented as genetic or was seen as insignificant since it was not related to the part of the body in which the potential problem would occur.

The authors suggest that eliciting and, when appropriate, changing people's representations prior to testing may enable those receiving negative results to be more reassured about their residual risk.

Comments

This paper shows the unfamiliarity of lay people with genetic testing. Genetic tests might therefore be perceived as less reliable and/or less reassuring than they should be based on a medical perspective. The findings in this paper are therefore of great interest to all who work in the field of (clinical) genetics or in the field of informing the population on genetics. Besides changing people's representations prior to testing, more general information about genetic testing provided to the population at large is needed in order to ensure that genetic test results will be perceived as reliable and become as reassuring as they should be.

However, there might also be other reasons why some patients are not reassured by negative genetic test results and perceive a continuing risk for themselves and their children. In a study of hereditary non-polyposis colorectal cancer, McAllister [2] found that family members who were offered a predictive test for hereditary non-polyposis colorectal cancer had very fixed ideas about what the result(s) of the test(s) would be, and that these beliefs might affect adjustment to predictive test results. Accordingly, some patients might still feel at risk for the disease, despite a negative test result. Moreover, in that study [2], patients were told during the pretest consultation that a negative test result is a guarantee that they will not develop polyposis, but that they still had the same risk as the general population of developing bowel cancer. Since these patients are familiar with this specific kind of cancer, they might be more afraid of it than the gener-

al population. It is known that individuals with a family history of cancer commonly overestimate their risk of developing cancer [3], even if they understand that the risk is small. This can be explained in terms of the availability heuristic, i.e. the tendency for an event (e.g. disease) to be judged more probable to the extent that it is more easily pictured or recalled. Peay and Peay [4] found that in particular for symptoms that might be indicative of cancer, patients perceived treatment as being more urgently necessary than did general practitioners. Annual bowel screening might therefore be very reassuring for this particular group of patients who were initially indicated as being at high risk for bowel cancer because of their family history, even though the genetic test results proved differently afterwards, and even if the genetic test results were understood correctly. Awareness of these processes might help genetic counselors to better support their clients.

A.M.C. Plass, Amsterdam

References

- 1 Michie S, Collins V, Halliday J, Marteau TM; FAP Collaborative Research Group: Likelihood of attending bowel screening after a negative genetic test result: The possible influence of health professionals. *Genet Test* 2002;6: 307–311.
- 2 McAllister M: Predictive genetic testing and beyond: A theory of engagement. *J Health Psychol* 2002;7:491–508.
- 3 Croyle RT, Lerman C: Risk communication in genetic testing for cancer susceptibility. *J Natl Cancer Inst Monogr* 1999;25:59–66.
- 4 Peay MY, Paey ER: The evaluation of medical symptoms by patients and doctors. *J Behav Med* 1998;21:57–81.

Mulvey S, Zachariah R, McIlwaine K, Wallace EM:

Do women prefer to have screening tests for Down syndrome that have the lowest screen-positive rate or the highest detection rate?

Prenat Diagn 2003;23:828–832.

DOI: 10.1159/000078508

Preferences for Down Syndrome Screening

The authors explored women's preferences for prenatal screening test performances. Pregnant women ($n = 120$) who wished to have a Down syndrome screening performed were asked to choose one of six hypothetical screening tests for Down syndrome. The study was performed at the Monash Medical Centre, Victoria, Australia. The tests to choose from had different detection rates ranging from 80 to 99% and the corresponding screening positive rates varied from 1 to 10%. Other information such as demographic characteristics – age, ethnicity and level of education – and regarding the performance of prenatal tests in earlier pregnancies was also requested.

The results of the questionnaire showed that most of the women (67%) prefer a screening test with a low screen-positive rate rather than a high detection rate. However, women of 37 years and older,

the age at which diagnostic tests are offered in Australia, preferred tests with a high detection rate and therefore a high screen-positive rate. Results of previous screening tests did not appear to affect the preferences of the women.

According to these results the authors conclude that age, rather than previous screening experiences, is the most significant factor determining the preferences of women for one of the possible screening tests. In the light of these findings, the authors suggest that different tests with different screen-positive and detection rates must be made available by the service providers. This would allow every individual woman to decide which screening test she wants to be performed, according to her personal evaluation of the outcomes of the screening test and the following diagnostic procedure as well as the possible birth of a Down child or the risk of miscarrying a normal child.

Comments

The authors conclude that expanding the same screening test standards for high-risk pregnant women to low-risk women may have negative effects due to the related effect of false-positive results such as unnecessary amniocentesis and avoidable psychological effects. However, the suggestion that individual women must be allowed to decide which specificity and sensitivity of the screening test performance they wished may cause more harm than good.

The two main aims of prenatal screening procedures are to allow the parents-to-be to make informed reproductive choices and to reduce the number of invasive procedures needed compared with age-based screening. Personal autonomy and individual choice are important aspects in the legitimization of prenatal genetic screening. Giving each woman the choice among different available prenatal screening possibilities allows her to choose according to her personal views and values. In spite of this, an increase in choices offered may also negatively influence the autonomy of the persons involved. Determining what kind of test is the best for each individual person is a complex issue that requires enough knowledge about the outcomes and consequences of the screening procedures. Since autonomy is assumed to be the capacity to reflect critically about their preferences, desires and values and to develop views on them, women have to be given adequate information about each test.

Screening tests introduce uncertainty, and some suggest that they may introduce more private (moral) dilemmas for pregnant women causing emotional reactions that can affect the biological development of the fetus [1]. More attention needs to focus on the technology implementation as opposed to the technology development, including experiences of women who have to decide on these offers.

Another aspect is that the number of high-risk women included in the study – only 19 in a total of 120 – is too small to conclude that older women usually prefer high-detection screening tests. As the authors suggest, a more detailed examination of the apparent age effect described must be performed using age-specific screening parameters, rather than the population-based scenarios used in their study. Additionally more information about what contributes to women's decision making and how it could be facilitated would be significant information in order to establish which screening program should be applied.

E. Garcia, Amsterdam

Reference

- 1 Getz L, Kirkengen AL: Ultrasound screening in pregnancy: Advancing technology, soft markers for fetal chromosomal aberrations, and unacknowledged ethical dilemmas. *Soc Sci Med* 2003;56:2045–2057.