Dystonia

J.M.S. Pearce

Emeritus Consultant Neurologist, Department of Neurology, Hull Royal Infirmary, Hull, UK

Dystonia, comes from modern Latin through German *Dystonie* (Hermann Oppenheim, Neurol Centralbl 1911; 30:1090–1107), from *dys-* + Greek *-tonos*. It is defined as: A state of disordered tonicity, especially of muscle tissue.

The best known and early example is dystonia musculorum deformans (torsion spasm), a rare hereditary neurological disease which causes (usually generalized) spasm and contortion of muscles and commonly develops during childhood.

After Oppenheim, another early example is found in 1916 (J Am Med, 11 Nov. 1437/1):

‘The paradoxical phenomenon of dystonia, when present, I believe to be a pathognomonic symptom of this disease and an important differential sign’.

The word tone itself had musical connotations. It derives from 13th century old French – *ton*, of the voice. Latin *tonus*, meant a stretching, quality of sound, tone, or accent; this derived from the Greek *tonos* similarly translated as stretching, tension, raising of voice, pitch. It is also applied to language and modes of expression: special or characteristic style or tendency of thought, feeling, behaviour or the spirit, character, tenor. In art, tone is the effect of the combination of light and shade, showing the degree of luminosity of a colour.

In modern usage, dystonic is applied to abnormal tension resulting in abnormal postures present in many disorders. Gross examples are seen in the athetoid form of cerebral palsy, and in the acquired dystonias – focal, segmental or generalised (table 1).

Torsion dystonia is characterised by torsion spasms of muscle contraction, which distort the limbs and trunk into dystonic postures. Dystonic spasms occur during voluntary movement (action dystonia); they may be repetitive and rhythmic or sustained in a fixed dystonic posture.

Table 1. Simplified classification of dystonia

<i>Primary, idiopathic</i>
Hereditary
Autosomal-dominant DYT1 – gene map locus 9q34
Autosomal-recessive DYT2
X-linked. DYT3 – gene map locus Xq13.1
<i>Symptomatic</i>
Acquired
Perinatal or traumatic brain damage
Encephalitis
Parkinsonism
Tumour
Deep cerebral infarct/haemorrhage
Drug- or toxin-induced (e.g. dopamine, neuroleptics)
Hereditary syndrome
Huntington’s disease
Wilson’s disease
Dystonia, DYT5; dopa-responsive; autosomal-dominant; gene map locus 14q22.1-q22.2
Neuroacanthocytosis
Factitious dystonia

Many sufferers of dystonia show both abnormal postures and involuntary movements. The latter are either called dystonic movements (torsion spasms) particularly when proximal (axial dystonia) or athetoid movements if they affect more distal segments. They can affect the whole body (generalized dystonia) or adjacent parts such as an arm and the neck (segmental dystonia), or it may be confined to one part (focal dystonia) as in spasmodic torticollis, blepharospasm, dystonic writer's cramp and oromandibular dystonia.

Early Descriptions

Standard references suggest the first description of 'A peculiar tonic cramp' was that of Marcus Walter Schwalbe from Ziehen's clinic in 1907 [1], but so-called hysteri-

cal symptoms complicate precise interpretation. In 1911, Hermann Oppenheim [2] and Georg Theodor Ziehen [3] published almost simultaneous reports. Oppenheim had described four new cases in Jewish children and deserves credit for attempts to formulate a new nosologic variety under the name: 'dysbasia lordotica progressiva dystonia: musculorum deformans', and for describing its 'dromedary gait'.

But the grandson of the Spanish neurologist Lluís Barraquer i Roviralta (1855–1928) has made a credible case [4] for his prior claim [5]. In 1897, he related the characteristic signs of dystonia musculorum deformans; however, he described them as athetosis not dystonia: probably an issue of semantics.

References

- 1 Schwalbe MW: Eine eigentümliche tonische Krampfform mit hysterischen Symptomen; inaugural dissertation, Berlin, 1907.
- 2 Oppenheim H: Über eine eigenartige Krampfkrankheit des kindlichen und jugendlichen Alters (Dysbasia lordotica progressiva, Dystonia musculorum deformans). *Neurol Centralbl* 1911;30:1090–1107.
- 3 Ziehen GT: Ein Fall von tonischer Torsionsneurose. Demonstrationen im Psychiatrischen Verein zu Berlin. *Neurol Zentralbl* 1911; 30: 109–110.
- 4 Barraquer Bordan L: Sobre l'Escola de Neurologia de l'Hospital de la Sta. Creu i St. Pau amb especial atenció a la personalitat i a l'obra del seu fundador: Lluís Barraquer i Roviralta. *Sant Pau* 1980;1:85–93.
- 5 Pearce JMS: Lluís Barraquer i Roviralta and the origins of torsion dystonia. *J Neurol Neurosurg Psychiatry* 2004;75:582.