Cephalic

J.M.S. Pearce

Emeritus Consultant Neurologist, Department of Neurology, Hull Royal Infirmary, Hull, UK

Medical terms related to the head are many. Those based on the Greek *kefhal* (cephal-), hence the classical Latin *cephalus*, are the commonest. In 81 AD, Aretaeus of Cappadocia used cephele and cephalgia to describe patterns of headache. The modern use appeared much later in English in the mid-19th century, and seems to be applied to indicate the head, including skull or brain.

The Greek prefix *cephal-* is found in cephalad – towards the head, cephalhaematoma – a bloody swelling beneath the pericranium, cephalin or kephalin – a phospholipid accelerating the coagulation of blood and cephalgia – head-ache. Enkephalos (encephalon) referred to the brain as a whole. The word cerebrum (Latin *cerebrum*) is more specific – pertaining to the brain proper. An early quotation dates to 1615 in Crooke's *Body of Man* [p. 475]:

'It filleth almost the whole Scull; and this is properly called Cerebrum or the Braine.'

The rhinencephalon is literally nose-brain; mesencephalon, the mid-brain – recorded in the French *mésocéphale* and mesencephalon in English was used by Robert Bentley Todd in *The cyclopaedia of Anatomy and Physiology* [1835–1859, vol. 3, 684/2] and in 1853 in the fourth edition of W.B. Carpenter's *Principles of Human Physiology* [p. 734]:

'The part of the encephalon known as the Tuber Annulare to which the name of Mesocephale has been given.'

We have anatomical applications in cephalic artery, cephalic ganglia, cephalic index, cephalic vein, cephalocele. In therapeutics, we have cephalosporins, the antibiotics developed from *Cephalosporium*, a genus of fungi. When investigating the head, we have echoencephalography and electroencephalography.

These combination words were developed mainly in the second half of the 19th century. The early uses of adjectival forms derived from this root are: microcephalic (micro – small), hydrocephalic (hydro – water), brachycephalic, (brachy – short), oxycephalic, porencephalic and so forth. Dolichocephalic (dolicho – long) is found in Darwin's *Descent of Man* [1871, vol. 1, ch. 4, p. 148]:

'... finds that short men incline more to brachycephaly, and tall men to dolichocephaly.'

The French *microcéphale* appears in E.F.M. Bosquillon 1787, in G. Cullen *Éléments de Médecine Pratique* [vol. 2, p. 508]. The part of the brain between the mesencephalon and prosencephalon (fore-brain) became known as diencephalic, pertaining to the diencephalon (modern Latin *di-*, from Greek *dia-* through + *encephalon*, brain).

There are numerous other examples where *cephal-* is incorporated into medical words – 'too many to get your head round' – to use a modern colloquialism. Examples are: anencephaly, encephalitis, encephalopathy, acrocephaly, bucephalus (ox-head, also the name of Alexander the Great's celebrated horse).

The word is not confined to Medicine. *Cephalaspsis* is a genus of fossil ganoid fishes found in the Old Red Sandstone, having a large buckler-shaped plate attached to the head. *Cephalopods* are a class of Mollusca, with a distinct head with 'arms' or tentacles attached, comprising cuttlefishes, the nautilus and numerous fossil species.

The older word cranium (medieval Latin from Greek *kranion*) is found in English in Bartholomew Traheron's *The Most Excellent Workes of Chirurgerye Made by J. Vigon* tr. 1543:

'The bone named Cranium, or the fyrst table of bones of the heade.'

Cranium is distinct from cephalon, referring specifically to the bones of the whole head collectively – the skull; hence the descriptors: intracranial and extracranial.

Many great art works show Cephalus and Aurora. She fell in love with Cephalus and tried to seduce him. Nicolas Poussin and Rubens both created paintings based on interpretations by the poet Ovid.