Citron Kinase Is Required for Postnatal Neurogenesis in the Hippocampus

James B. Ackman^a Raddy L. Ramos^a Matthew R. Sarkisian^b Joseph J. LoTurco^a^aDepartment of Physiology and Neurobiology, University of Connecticut, Storrs, Conn., and^bDepartment of Neurobiology, Yale University School of Medicine, New Haven, Conn., USA

Key Words

Neural stem cells • Neurogenesis • Dentate gyrus • Epilepsy

Abstract

The dentate gyrus is a site of continual neurogenesis in the postnatal mammalian brain. Here we investigated postnatal neurogenesis in the citron kinase (citron-K) null-mutant rat (*flathead*). The *flathead* rat has substantial deficits in embryonic neurogenesis that are due to failed cytokinesis and cell death. We report here the loss of citron-K function has an even severer effect on postnatal neurogenesis in the dentate gyrus. Analysis of phosphorylated histone H3 expression in postnatal neurogenic regions of the *flathead* mutant revealed a complete lack of mitotic cells in the dentate gyrus and a large reduction in the number of dividing cells in the *flathead* subventricular zone. Examination of 5-bromodeoxyuridine incorporation in the *flathead* rat revealed that the *flathead* rat had a 99% reduction in the number of newly generated cells in the dentate gyrus at postnatal day 10. In addition, doublecortin-positive cells were essentially absent from the postnatal *flathead* dentate gyrus which also lacked the vimentin- and nestin-positive radial glia scaffold that defines the neurogenic niche in the postnatal subgranular zone. Together these results indicate that postnatal neurogenesis in the dentate gyrus is eliminated by loss of citron-K function, and suggests that a citron-K-dependent progenitor lineage forms the postnatal neuronal progenitor population in the dentate gyrus.

Copyright © 2007 S. Karger AG, Basel

Introduction

Neurogenesis is a continual process in the dentate gyrus of the hippocampus. Generation of dentate granule neurons has been shown to occur in the adult brain of rodents [Altman and Das, 1965; Cameron et al., 1993; Kaplan and Hinds, 1977], primates [Gould et al., 1999; Kornack and Rakic, 1999], and humans [Eriksson et al., 1998]. These newly born dentate granule neurons originate from a pool of neural stem cells present within the subgranular zone (SGZ) of the dentate gyrus [Gage et al., 1995; Palmer et al., 1997; Seri et al., 2001]. Neural stem cells present in neurogenic regions of the adult mammalian brain, including the SGZ and the subventricular zone (SVZ), have been identified as a constituent of the radial glia-astrocyte lineage [Doetsch et al., 1999a, b; Laywell et al., 2000; Imura et al., 2003; Merkle et al., 2004; Garcia et al., 2004; Seri et al., 2001].

Though neurogenesis occurs in the dentate gyrus throughout life, the peak of dentate granule cell production occurs during the first 2 weeks of postnatal development in rodents [Bayer, 1980; Schlessinger et al., 1975]. This period of peak neurogenesis occurs following a number of discrete developmental steps that take place during embryonic and postnatal development [Altman and Bayer, 1990a, b; Li and Pleasure, 2005]. Among the initial steps in dentate gyrus development is the migration of the earliest-born granule neurons from their place of birth in the neuroepithelium to the dentate gyrus anlage to form the primordial granule cell layer [Altman

and Bayer, 1990a, b]. These pioneer granule cells migrate along the radial glia scaffold that extends from the ventricular zone (VZ) to the dentate primordium together with a secondary population of neurogenic progenitors [Altman and Bayer, 1990a, b; Rickmann et al., 1987]. This secondary progenitor population is established in the hilar region of the developing dentate formation during late embryonic development and undergoes a radial transformation to form the SGZ neuroepithelium from which postnatal and adult neurogenesis will originate [Altman and Bayer, 1990a, b; Eckenhoﬀ and Rakic, 1984; Guéneau et al., 1982; Alvarez-Buylla and Lim, 2004].

The *flathead* mutant rat exhibits a central nervous system-specific phenotype that includes severe micrencephaly and frequent seizures [Sarkisian et al., 1999]. The numbers of neurons in the cerebellum, dentate gyrus, olfactory bulb, and upper neocortical layers are dramatically reduced in the *flathead* rat and the brain is 50% smaller than that of the wild type at birth. Although migration in and lamination of the *flathead* neocortex are intact, up to 50% of pyramidal and nonpyramidal neurons are binucleate, indicating a cytokinesis failure during embryonic development [Sarkisian et al., 2001]. In addition, loss of citron kinase (citron-K) function causes deficits in mitosis [LoTurco et al., 2003] and an increase in apoptosis [Di Cunto et al., 2000; Roberts et al., 2000; Sarkisian et al., 2001]. The *flathead* mutation has been identified as a premature stop codon in the first exon of the citron-K gene [Sarkisian et al., 2002]. Citron-K is expressed in dividing cells during embryonic development where it localizes to the cleavage furrow of cells undergoing cytokinesis but is absent in the *flathead* mutant [Sarkisian et al., 2002; Di Cunto et al., 2000]. It is hypothesized that citron-K is essential for cytokinesis in neurogenic divisions, since citron-K is expressed in the developing cerebral cortex during the peak period for generation of late-born cortical neurons, and because many neurons in the *flathead* mutant are binucleate whereas glia are not binucleate [LoTurco et al., 2003].

Here, we show that there is a dramatic reduction in the number of neurons in the dentate gyrus of the *flathead* rat, greater than the decrease in the brain at large. The decrease is associated with a near loss of dividing cells and new neurons in the postnatal dentate germinal zone (DGZ). In comparison, there is a more modest but significant reduction in the numbers of dividing cells and neuroblasts in the postnatal SVZ and rostral migratory stream (RMS) of the *flathead* rat. These results demonstrate that postnatal neurogenesis is limited in the SVZ/RMS and virtually absent in the dentate gyrus after loss of citron-K function.

Materials and Methods

Animals and 5-Bromodeoxyuridine Injections

All animal procedures were performed according to protocols approved by the University of Connecticut Animal Care and Use Committee. *Flathead* mutant rats were obtained from bred colonies at the University of Connecticut. For all experiments, normal littermates from *flathead* litters were used as controls and were designated as wild type. The normal littermates included both heterozygous and homozygous wild-type animals. Brains from *flathead* and littermate rats were examined at embryonic day 14 (E14), postnatal day 0 (P0), P7, P14, P16, P17, P19, P21, P22, and P24.

As for 5-bromodeoxyuridine (BrdU) injections, pregnant dams received 50 mg/kg BrdU intraperitoneally at 14 and 15 days of gestation. Postnatal *flathead* and control pups received 40 mg/kg BrdU intraperitoneally. Embryonic rat brains were collected into cold HBSS and fixed overnight in 4% paraformaldehyde (PFA). Postnatal rat pups were euthanized with wet ice (less than P7) or isoflurane (greater than P7) and perfused intracardially with phosphate-buffered saline (PBS) followed by 4% PFA. Brains were dissected out and postfixed overnight at 4°C in 4% PFA followed by embedding in 2% agarose. They were cut at 40 μ m on a vibratome (Leica VT-1000S). Free-floating sections were stored at -20°C in a cryoprotectant solution (PVP-40, 30% sucrose, in ethylene glycol and 0.2 M phosphate buffer, pH 7.4) until use.

Immunocytochemistry, Histochemistry, and Confocal Microscopy

BrdU immunocytochemistry was performed by rinsing free-floating brain sections in PBS followed by denaturation in 2 N HCl for 30 min at room temperature. Sections were neutralized in 0.1% sodium borate and rinsed in PBS before continuation with the standard immunocytochemistry protocol. The standard immunocytochemistry protocol included blocking sections in 5% donkey serum (Sigma, No. D9663) + 0.3% Triton X-100 for 1 h at room temperature followed by incubation in primary antibodies overnight at 4°C. After rinsing for 1 h, sections were incubated in secondary antibodies for 1 h at room temperature followed by a final rinse in PBS, mounted on glass slides, and coverslipped in Prolong Antifade (Molecular Probes). The primary antibodies used were: rat anti-BrdU (1:100, Accurate, No. OBT0030), goat antiodoublecortin (1:100, Santa Cruz, No. sc-8066), mouse anti-NeuN (1:1,000, Chemicon, No. MAB377), mouse anticalbindin D28K (1:1,000, Sigma, No. C8666), rabbit antiphospho-histone H3 (1:500, Chemicon, No. 06-570), mouse antivimentin (1:200, Chemicon, No. MAB3400), mouse antinestin (1:200, Chemicon, No. MAB353), and mouse anti-GFAP (1:200, Sigma, No. G3893). The secondary antibodies used were: donkey antirabbit Alexa 488 (1:200, Molecular Probes), donkey antimouse Alexa 488 (1:200, Molecular Probes), donkey antirat Cy3 (1:200, Jackson ImmunoResearch), donkey antimouse Alexa 568 (1:200, Molecular Probes), and donkey antimouse Cy5 (1:200, Jackson ImmunoResearch). Some sections were counterstained with the nuclear label TO-PRO-3 (Molecular Probes).

For use with nonfluorescent histochemistry, sections were rinsed several times with 2.5% normal goat serum in PBS and then incubated in biotinylated secondary antibodies (biotinylated goat antimouse; 1:200, Vector Labs, Burlingame, Calif., USA) for 2 h at room temperature. Sections were rinsed several times with

PBS and incubated for 1 h in an avidin-horseradish peroxidase mixture. Then they were rinsed in PBS and reacted with 0.05% diaminobenzidine in the presence of 0.0015% H₂O₂. Sections were collected onto gelatin-coated slides, dried for several hours, and coverslipped with Cytoseal.

For Timm histochemistry, rats were perfused with a sodium sulfide medium (4.76 g NaH₂PO₄, 4.68 g Na₂S in 400 ml dH₂O) followed by 4% PFA. Brains were cryoprotected in 30% sucrose and frozen over liquid N₂. Sections (20 μ m) were developed in the dark for 30–60 min in a solution containing gum arabic (120 ml of a 20% stock), citrate buffer (20 ml of a 2 M stock), hydroquinone (60 ml of a 5.7% stock), and 0.17 g silver nitrate. Sections were dehydrated in ethanol, cleared in xylene and coverslipped with DPX.

Slides were imaged with either a Nikon E400 microscope and Spot camera or with a laser scanning confocal microscope (Leica TCS-SP2; laser lines at 488, 543, 633). Confocal images were acquired at 0.3–1.0 μ m per z-section.

Quantification

For quantification of mitotic cells, the number of phosphorylated histone H3 (phosH3)-positive cells per sagittal section was counted for the dentate gyrus and the SVZ including the first 300 μ m of the RMS from the *flathead* rats (ages P10, P16, and P22) and the wild-type rats (ages P16 and P22). For BrdU quantification, animals aged P17 or P22 that had received BrdU injections at P10 or P19, respectively, were used. In the dentate gyrus, the number of BrdU-positive cells per sagittal section was counted. For the SVZ, confocal images (8-bit indexed color) from sagittal sections were thresholded in Image J with a tolerance of 21 and the polygon selection tool was used to outline the SVZ and the first 300 μ m of the RMS. The analyze particles tool with a pixel radius of 3 was used to automatically count the number of BrdU-positive cells in the SVZ. Several sections of the *flathead* and littermate SVZ were hand counted to confirm the validity of the automatic counting procedure. For both phosH3 and BrdU quantification in the dentate gyrus and SVZ, the mean and SEM for the number of positive cells per section were calculated for the *flathead* rat and wild type. For all conditions, quantification was performed on an average of 13 sagittal sections from the medial extent (range of 0.5–2.5 mm from midline). Differences between compared means were determined with Student's t test, with $p < 0.05$ considered significant (for all experiments: N = number of animals, n = number of stained sections).

Results

Granule Layer Dysgenesis in the *flathead* Dentate Gyrus

The brain of the citron-K null mutant (*flathead*) is approximately one-half the size of that of the wild type at birth. Areas of the central nervous system that undergo significant amounts of late embryonic or postnatal neurogenesis, such as the retina, cerebellum, olfactory bulb, and dentate gyrus, are the regions most affected by a lack of citron-K function [Di Cunto et al., 2000; Roberts et al.,

2000]. The dentate gyrus is markedly reduced in size in the *flathead* rat (fig. 1a–c). Nissl-stained sagittal and coronal sections of P21 wild-type and *flathead* brains reveal that the inferior blade of the dentate gyrus is missing in the *flathead* rat (fig. 1b, c). The superior blade of the dentate gyrus that is present in the *flathead* rat is severely reduced in the dorsoventral and mediolateral dimensions. In addition, the *flathead* dentate gyrus is reduced in the rostrocaudal extent. In contrast to the dentate gyrus, the CA3–CA1 pyramidal layers of Ammon's horn are more moderately reduced in size in the *flathead* rat and have no apparent gross disruption in organization. To confirm that the cells remaining in the disrupted granule layer of the *flathead* dentate gyrus were granule neurons, we performed calbindin D28K immunocytochemistry on horizontal sections of P21 *flathead* and wild-type hippocampi (fig. 1e). We found that calbindin was expressed by cells in the granule cell layer of the *flathead* rat. Furthermore, we performed Timm's silver sulfide staining on horizontal sections of P14 wild-type and *flathead* hippocampi (fig. 1d). We found that Timm staining intensely labeled granule cell mossy fiber axons in the hilar region and CA3 pyramidal layer of the *flathead* hippocampus in a pattern appropriate for mature dentate granule neurons.

Loss of Progenitor Cells in the Postnatal *flathead* Dentate Gyrus

Next we determined whether proliferating cells were present in the SGZ of the *flathead* dentate gyrus using phosH3 and BrdU immunocytochemistry (fig. 2). There was a lack of phosH3-positive dividing cells in the *flathead* dentate gyrus at P16 and P22 (fig. 2a, b, f). The number of phosH3-positive cells per section in the *flathead* rat was reduced by 97% with respect to the wild type (0.3 ± 0.1 , N = 4, n = 19 for the *flathead* rat; 8.8 ± 2.9 , N = 2, n = 5 for the wild type; $p < 0.05$). Consistent with a lack of proliferating cells in the dentate gyrus, the number of cells in the P17 *flathead* dentate gyrus that incorporated BrdU at P10 was reduced by 99% (fig. 2c, d, g; 3.5 ± 0.9 cells/section, N = 7, n = 11 for the *flathead* rat; 364.3 ± 32.0 cells/section, N = 7, n = 18 for the wild type; $p < 0.001$).

Next we tested if dentate granule neurons in the *flathead* rat were generated during embryonic development. Consistent with an embryonic birth date, we found that granule neurons in the P19 *flathead* dentate gyrus were labelled by BrdU administered at E14 and E15 (fig. 2e; N = 1). Note the presence of several binucleate NeuN-positive cells near the internal limb of the dentate gyrus (arrowheads, fig. 2e).

Fig. 1. Dysgenesis of the granule cell layer in the *flathead* dentate gyrus. **a** Nissl-stained sagittal sections of wild-type and *flathead* brains at P21. **b** Higher-magnification comparison of Nissl staining in sagittal sections of the wild-type and *flathead* hippocampus. **c** Higher-magnification comparison of Nissl staining in coronal sections of the P21 wild-type and *flathead* hippocampus. **d** Timm staining of granule cell mossy fiber axons (arrows) in horizontal sections of the P14 wild-type and *flathead* hippocampus. **e** Expression of calbindin in horizontal sections of the P21 wild-type and *flathead* dentate gyrus. In the images, medial is down and rostral is to the right. Scale bars: 1 mm (**a**); 250 μm (**b-e**).

Decreased Proliferation in the SVZ and RMS of the flathead Rat

Since the SVZ/RMS is a postnatal neurogenic region, we tested for expression of the immature neuron marker, doublecortin, in the SVZ/RMS of the *flathead* rat. Doublecortin expression was reduced in the SVZ and RMS of the *flathead* rat at P22 (fig. 3a). Next we determined whether there was a decrease in the numbers of mitotic progenitors in the SVZ/RMS of the *flathead* rat. Analysis of phosH3 immunocytochemistry on sagittal sections of the P22 *flathead* brain revealed that the number of phosH3-expressing cells per section was reduced by 92% in the *flathead* SVZ/RMS (fig. 3b, d; 11.8 ± 1.7 cells/section, $N = 4$, $n = 19$ for the *flathead* rat; 146.8 ± 26.9 cells/section, $N = 2$, $n = 4$ for the wild type; $p < 0.05$). BrdU immunocytochemistry on sections from P17 animals that received BrdU at P10 confirmed a reduction in newly generated cells in the *flathead* SVZ/RMS (fig. 3c, e). The decrease in BrdU-labeled cells per section was 83% in the *flathead* SVZ/RMS (179.4 ± 39.8 cells/section, $N = 5$, $n = 15$ for the *flathead* rat; $1,059.7 \pm 113.2$ cells/section, $N = 4$, $n = 9$ for the wild type; $p < 0.001$).

Altered Neurogenic Niche in the flathead Dentate Gyrus

Next we assessed whether doublecortin-expressing cells were present in the postnatal *flathead* dentate gyrus. In contrast to the wild type where there was doublecortin immunoreactivity throughout the dentate gyrus at P17 and P22, most immunostained sections from the *flathead* rat contained no doublecortin-positive cells or fibers (fig. 4a, b; 0.39 ± 0.29 cells/section, $N = 8$, $n = 23$ for the *flathead* rat; $N = 3$, $n = 8$ for the wild type). In the few sections of the *flathead* hippocampus where isolated doublecortin immunoreactivity was present, the positive cells were located in the hilar region between the dentate

Fig. 2. Proliferating cells are absent in the postnatal *flathead* dentate gyrus. **a** Lack of phosH3-positive nuclei (white) in the *flathead* SGZ at P16. The section is counterstained with the nuclear label TO-PRO-3 (blue). **b** Lack of phosH3-positive nuclei in the *flathead* SGZ at P22. Dashed lines outline the granule cell layer. **c** Decreased numbers of BrdU-positive cells in the postnatal *flathead* dentate gyrus. The images are from P17 wild-type and *flathead* animals that received BrdU injections at P10. The few BrdU-positive objects next to the dentate gyrus in this image are blood vessels. **d** Lack of newly generated granule neurons in the *flathead* dentate gyrus. BrdU is green and NeuN is red. **e** Dentate granule neurons in the *flathead* rat incorporate BrdU at E14–E15 (arrows). Image is from a P19 *flathead* rat that received BrdU on E14 and E15. Binucleate neurons are indicated by arrowheads. **f** Quantifi-

cation of phosH3-positive nuclei in the SGZ. Numbers are expressed as the mean number of phosH3-positive cells in the SGZ per section ($p < 0.05$). **g** Quantification of the number of BrdU-positive cells in the dentate gyrus. Numbers are expressed as the mean number of BrdU-positive nuclei per section ($p < 0.001$). Scale bars: 40 μ m (**a–e**).

Fig. 3. Decreased proliferation in the postnatal SVZ of the *flathead* rat. **a** Expression of doublecortin (DCX) and phosH3 in the SVZ of the wild-type and *flathead* rat at P22. **b** Reduced numbers of phosH3-positive nuclei in the *flathead* SVZ at P22. The images are from the same sections shown in **a**. **c** Decreased numbers of BrdU-positive cells in the postnatal *flathead* SVZ. The images are from P17 wild-type and *flathead* animals that received BrdU injections at P10. **d** Quantification of phosH3-positive nuclei in the

horn and CA3 (fig. 4b), whereas doublecortin-positive cells were never found outside the granule cell layer or SGZ in the wild-type dentate gyrus. In sagittal stained sections, the dendrites of hilar doublecortin-positive cells in the *flathead* rat were generally oriented in the caudal or ventral directions towards the dentate molecular layer.

The postnatal SGZ of the dentate gyrus contains radial-type astrocytes that originate from radial glia during development [Eckenhoff and Rakic, 1984; Rickmann et al., 1987]. Radial-type astrocytes generate granule neurons in the dentate gyrus [Seri et al., 2004] and express the intermediate filament proteins vimentin, nestin, and GFAP [Eckenhoff and Rakic, 1984; Rickmann et al., 1987; Lendahl et al., 1990; Seri et al., 2004]. To determine whether radial-type astrocytes were present in the postnatal *flathead* SGZ, we performed vimentin, nestin, and GFAP immunocytochemistry on sagittal sections of the wild-type and *flathead* dentate gyrus at P21–P24. The postnatal *flathead* dentate gyrus lacked vimentin- and nestin-positive radial fibers (fig. 4c, d) and contained few GFAP-positive radial fibers (arrows in fig. 4e). In contrast to the reduction in radial-type astrocytes, multipolar astrocytes that expressed GFAP were abundant in the *flathead* SGZ, as well as in the hilus and in the molecular layer. In other postnatal neurogenic regions, such as the SVZ/RMS, both radial glia and multipolar-type astrocytes that had intense staining for vimentin, nestin, and GFAP were present in the *flathead* rat (data not shown).

Reduced Migration of Dentate Precursors during Development of the *flathead* Dentate Gyrus

Next we examined the cytoarchitecture of germinative matrices associated with the development of the dentate gyrus. The hippocampal neuroepithelium is established along the medial wall of the lateral ventricle during embryonic development (fig. 5a). An indentation of the hippocampal neuroepithelium, the dentate notch, is the location of the primary dentate neuroepithelium from which the earliest-generated granule cells originate as well as migrating progenitors that establish the secondary dentate

SVZ. Numbers are expressed as the mean number of phosH3-positive cells in the SVZ per section ($p < 0.05$). **e** Quantification of the number of newly generated cells in the SVZ. Numbers are expressed as the mean number of BrdU-positive nuclei in the SVZ ($p < 0.001$). Scale bars: 40 μm (**a–c**).

matrix. We analyzed Nissl-stained coronal sections of the hippocampal neuroepithelium at E14 to see if any gross cytoarchitectural abnormalities were present in the primary dentate neuroepithelium of the *flathead* rat before formation of the secondary dentate matrix. No distinct alterations were present in the primary dentate neuroepithelium of the *flathead* rat at E14 (fig. 5b). The secondary dentate matrix is a germinative matrix located adjacent to the VZ of the dentate neuroepithelium and becomes evident around E18 in the rat [Altman and Bayer, 1990a]. The secondary dentate matrix consists of proliferative cells and migratory cells and extends tangentially towards the dentate gyrus as a subpial migration. This dentate migration is robust during late embryonic and early postnatal development (E20–P5 in the rat) [Altman and Bayer, 1990a, b] and contains neuroblasts that migrate to the granule cell layer and progenitors that migrate to establish the proliferative tertiary dentate matrix in the hilus (fig. 5c). We examined the dentate migration in the perinatal hippocampus, a time during which there are large numbers of cells migrating towards the dentate gyrus. Nissl-stained horizontal sections of the P0 *flathead* and wild-type hippocampus revealed that the dentate migration in the *flathead* rat is distinctly reduced (fig. 5d). In addition, there appeared to be a lack of cells aggregating in the region of the internal dentate blade in the *flathead* rat, as well as an areal reduction in the region of the tertiary dentate matrix. In contrast to the dentate migration, which appeared thin between the pial surface and CA3 (boxed area in fig. 5d), the SVZ, Ammon's horn, stratum radiatum, and the fimbria did not appear particularly reduced in the *flathead* rat at P0. Additionally, we observed an increased number of pyknotic nuclei in Nissl-stained sections at P0 in the secondary dentate matrix, the dentate migration, and the tertiary matrix in the *flathead* rat (data not shown). Increased cell death in the dentate germinative matrices is consistent with previous studies indicating that a lack of citron-K function leads to failed cytoki-

Fig. 4. Altered neurogenic niche in the *flathead* dentate gyrus. **a** Doublecortin (DCX) expression in the wild-type and *flathead* dentate gyrus at P17. Notice the dramatic reduction in doublecortin-positive cells and processes within the dentate granule cell layer of the *flathead* rat. **b** Doublecortin expression in the dentate gyrus of the wild-type and *flathead* rat at P22. **c** Vimentin expression in the dentate gyrus of the wild-type and *flathead* rat at P22. Notice the high density of vimentin-positive radial fibers in the wild-type dentate gyrus and the lack of radial fibers in the *flathead* rat. **d** Nestin expression in the dentate gyrus of the wild-type and *flathead* rat at P24. Nestin expression was located in endothe-

lial cells, multipolar astrocytes, and radial-type astrocytes in the wild-type dentate gyrus, whereas in the *flathead* dentate gyrus only nestin-positive endothelial cells and multipolar astrocytes were found. Arrows indicate examples of nestin-positive radial fibers in the wild-type dentate gyrus. **e** GFAP immunohistochemistry in the dentate gyrus of the wild-type and *flathead* rat at P21. Notice the lack of GFAP-positive radial fibers in the *flathead* dentate gyrus. Arrows indicate GFAP-positive radial fibers. Scale bars: 40 μ m (**a–d**); 25 μ m (**e**).

Fig. 5. Diminished dentate migratory stream during development of the *flathead* dentate gyrus. **a** Location of the hippocampal neuroepithelium (hi) in the developing rat brain. Dentate notch is indicated by an asterisk. chp = Choroid plexus. **b** Nissl-stained coronal sections of the wild-type (wt) and *flathead* (fh) hippocampal neuroepithelium at E14. Note the similarity in appearance between the wild-type and *flathead* ammonic neuroepithelium (ah) and primary dentate neuroepithelium (dgp). chp = Choroid plexus. **c** Overview of dentate gyrus development during the late embryonic and early postnatal periods. The dentate migratory

stream (dgm) consists of neuroblasts and progenitors which migrate to form the internal blade of the dentate gyrus (DGi) and the proliferative tertiary dentate matrix (dgt) in the hilus. AH = Ammon's horn; DGe = external blade of the dentate gyrus; fi = fimbria; dgs = secondary dentate matrix; ah = ammonic neuroepithelium; dgp = primary dentate neuroepithelium. **d** Nissl-stained horizontal sections of the wild-type (wt) and *flathead* (fh) hippocampus at P0. Note the decreased number of cells in the *flathead* dentate migratory stream (dgm) compared with the wild-type dentate migratory stream (arrows). The panel of high-magnifica-

nesis and massive apoptosis in germinal zones such as the ganglionic eminence, neocortical VZ/SVZ, and the cerebellar granule layer [Roberts et al., 2000; Sarkisian et al., 2001; Di Cunto et al., 2000]. By P7, the density of cells in the dentate migration has decreased in both the wild-type and *flathead* rats (fig. 5e). Notice that by P7 the dentate gyrus has greatly increased in size and the internal blade of the dentate gyrus has formed in the wild type, whereas in the *flathead* rat, the dentate gyrus has not undergone a similar increase in size and the internal blade is mostly absent.

Discussion

Previous studies have demonstrated severe deficits in embryonic neurogenesis and increased apoptosis in citron-K null mutants [Di Cunto et al., 2000; Roberts et al., 2000; Sarkisian et al., 2002]. Here, we show that postnatal neurogenesis is depleted in the citron-K null-mutant rat (*flathead*). The depletion of postnatal neurogenesis in the *flathead* rat can most likely be attributed to a loss of mitotic progenitors in postnatal neurogenic regions as proliferation was decreased in the SVZ/RMS and nearly completely absent in the dentate gyrus. These results indicate that citron-K function is required for development of the neurogenic progenitor pool in the postnatal dentate gyrus.

Among the developmental steps required for formation of the dentate gyrus is the establishment of a tertiary progenitor population in the SGZ during late embryonic and early postnatal development (also termed the 'tertiary germinative matrix'; see Altman and Bayer [1990a, b]). It seems likely from our results that the tertiary precursor pool fails to be established in the dentate gyrus of the citron-K null mutant (*flathead*) as there was a lack of mitotic progenitors in the SGZ, as well as a lack of newly generated neurons in the granule cell layer of the postnatal *flathead* hippocampus. In addition, radial-type astrocytes were absent in the postnatal *flathead* SGZ. Since radial-type astrocytes serve as progenitors of new granule neurons in the adult dentate gyrus [Seri et al., 2004], these results indicate that formation of the neurogenic niche is disrupted in the *flathead* SGZ.

tion images of the dentate migratory stream below were taken from the region outlined by the boxes in the upper panel. **e** Nissl-stained horizontal sections of the wild-type (wt) and *flathead* (*fh*) hippocampus at P7. Scale bars: 50 μ m (**b**); 100 μ m (**d**, upper panel; **e**); 25 μ m (**d**, lower panel).

Though there was a lack of neurogenesis in the postnatal *flathead* dentate gyrus, granule cells were present in the rudimentary granule layer of the *flathead* rat. These *flathead* granule cells possessed key differentiated granule neuron characteristics such as calbindin expression and Timm staining of mossy fiber axons. In addition, granule cells that were present in the *flathead* dentate gyrus were found to incorporate BrdU administered at E14–E15, a time during which only the primordial granule neuron population is being generated. The presence of a rudimentary granule cell layer consisting of early-born granule neurons in the *flathead* rat indicates that the disruption in dentate gyrus development is fundamentally different from other rodent models involving deficits in SGZ precursor pool establishment such as Wnt signaling and CXCR4 mutants where the dentate gyrus completely fails to form [Galceran et al., 2000; Lee et al., 2000; Zhou et al., 2004; Bagri et al., 2002; Lu et al., 2002]. These results indicate that early in development generation of the pioneer granule neurons and migration into the dentate anlage can occur in the absence of citron-K function but that the subsequent progenitor-progenitor or progenitor-neuron divisions from which the majority of granule neurons originates require citron-K function.

Exactly how citron-K function is involved in the development of the SGZ progenitor population is not known. During embryonic development, citron-K is expressed in a portion of dividing progenitors along the VZ surface resulting in about half of cortical neurons being multinucleate due to cytokinesis failure [Sarkisian et al., 2002]. The study of the postnatal brain with in situ hybridization has revealed that the citron-K message is expressed in the external granule layer of the cerebellum as well as in the SVZ and RMS of the forebrain [Di Cunto et al., 2000]. Although citron-K expression has not been reported in the postnatal dentate gyrus, it is possible that it is present at low levels and that a lack of citron-K function leads to progenitor cell depletion in the SGZ due to cytokinesis failure and apoptosis. Alternatively, a lack of citron-K function could cause progenitor cell depletion in the hippocampal neuroepithelium, before the tertiary precursor population migrates into the hilus and SGZ. Our results showing a diminished dentate migration from the secondary germinative matrix to the dentate formation indicate that a lack of citron-K function may deplete progenitors and neuroblasts before their migration into the hilus and the granule cell layer.

Citron-K mutants exhibit frequent seizures that begin near the end of the first postnatal week [Sarkisian et al., 1999; Di Cunto et al., 2000]. Temporal lobe epilepsy is

often associated with disorganization or loss of cells in the granule cell layer of the dentate gyrus [Houser, 1990, 1992]. It seems possible that frequent seizure activity may have a detrimental effect on development of the *flathead* dentate gyrus. However, it seems unlikely that the frequent seizures in the *flathead* rat would cause the loss of the SGZ progenitor population. First, seizures in the *flathead* rat start at the end of the first postnatal week, after the secondary progenitor population has migrated to and become established in the SGZ/hilus [Sarkisian et al., 1999]. Second, seizure activity has been reported to increase granule neuron neurogenesis in the dentate gyrus [Parent et al., 1997]. Although the results presented here did not include analysis of BrdU incorporation in the postnatal *flathead* brain earlier than the first postnatal week, we have examined BrdU incorporation at P2 in *flathead* rats and found that the lack of dentate granule neuron generation was present at this early postnatal age

as well (data not shown). We have not assessed proliferation at this early postnatal age with mitotic markers or short survival BrdU assays, but examination of the neonatal *flathead* hippocampus with Nissl staining indicates that the morphological abnormalities in dentate gyrus development are present in the *flathead* rat within the first postnatal week. In conclusion, postnatal neurogenesis in the dentate gyrus is dependent upon citron-K function, and as there is an apparent loss of dividing cells in postnatal DGZ, we further suggest that citron-K function is required for the production of the SGZ progenitor population from earlier, perhaps fetal, progenitors.

Acknowledgement

This work was supported by NIH grant MH056524 to J.J.L.

References

- Altman J, Bayer SA (1990a): Mosaic organization of the hippocampal neuroepithelium and the multiple germinal sources of dentate granule cells. *J Comp Neurol* 301:325–342.
- Altman J, Bayer SA (1990b): Migration and distribution of two populations of hippocampal granule cell precursors during the perinatal and postnatal periods. *J Comp Neurol* 301:365–381.
- Altman J, Das GD (1965): Autoradiographic and histological evidence of postnatal hippocampal neurogenesis in rats. *J Comp Neurol* 124:319–335.
- Alvarez-Buylla A, Lim DA (2004): For the long run: maintaining germinal niches in the adult brain. *Neuron* 41:683–686.
- Bagri A, Gurney T, He X, Zou YR, Littman DR, Tessier-Lavigne M, Pleasure SJ (2002): The chemokine SDF1 regulates migration of dentate granule cells. *Development* 129:4249–4260.
- Bayer SA (1980): Development of the hippocampal region in the rat. I. Neurogenesis examined with ³H-thymidine autoradiography. *J Comp Neurol* 190:87–114.
- Cameron HA, Woolley CS, McEwen BS, Gould E (1993): Differentiation of newly born neurons and glia in the dentate gyrus of the adult rat. *Neuroscience* 56:337–344.
- Di Cunto F, Imarisio S, Hirsch E, Broccoli V, Bulfone A, Migheli A, Atzori C, Turco E, Triolo R, Dotto GP, Silengo L, Altruda F (2000): Defective neurogenesis in citron kinase knockout mice by altered cytokinesis and massive apoptosis. *Neuron* 28:115–127.
- Doetsch F, Caille I, Lim DA, Garcia-Verdugo JM, Alvarez-Buylla A (1999a): Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 97:703–716.
- Doetsch F, Garcia-Verdugo JM, Alvarez-Buylla A (1999b): Regeneration of a germinal layer in the adult mammalian brain. *Proc Natl Acad Sci USA* 96:11619–11624.
- Eckenhoff MF, Rakic P (1984): Radial organization of the hippocampal dentate gyrus: a Golgi, ultrastructural, and immunocytochemical analysis in the developing rhesus monkey. *J Comp Neurol* 223:1–21.
- Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA, Gage FH (1998): Neurogenesis in the adult human hippocampus. *Nat Med* 4:1313–1317.
- Gage FH, Coates PW, Palmer TD, Kuhn HG, Fisher LJ, Suhonen JO, Peterson DA, Suhr ST, Ray J (1995): Survival and differentiation of adult neuronal progenitor cells transplanted to the adult brain. *Proc Natl Acad Sci USA* 92:11879–11873.
- Galceran J, Miyashita-Lin EM, Devaney E, Rubenstein JL, Grosschedl R (2000): Hippocampus development and generation of dentate gyrus granule cells is regulated by LEF1. *Development* 127:469–482.
- Garcia AD, Doan NB, Imura T, Bush TG, Sofroniew MV (2004): GFAP-expressing progenitors are the principal source of constitutive neurogenesis in adult mouse forebrain. *Nat Neurosci* 7:1233–1241.
- Gould E, Reeves AJ, Fallah M, Tanapat P, Gross CG, Fuchs E (1999): Hippocampal neurogenesis in adult Old World primates. *Proc Natl Acad Sci USA* 96:5263–5267.
- Guéneau G, Privat A, Drouet J, Court L (1982): Subgranular zone of the dentate gyrus of young rabbits as a secondary matrix. A high-resolution autoradiographic study. *Dev Neurosci* 5:345–358.
- Houser CR (1990): Granule cell dispersion in the dentate gyrus of humans with temporal lobe epilepsy. *Brain Res* 535:195–204.
- Houser CR (1992): Morphological changes in the dentate gyrus in human temporal lobe epilepsy. *Epilepsy Res Suppl* 7:223–234.
- Imura T, Kornblum HI, Sofroniew MV (2003): The predominant neural stem cell isolated from postnatal and adult forebrain but not early embryonic forebrain expresses GFAP. *J Neurosci* 23:2824–2832.
- Kaplan MS, Hinds JW (1977): Neurogenesis in the adult rat: electron microscopic analysis of light radioautographs. *Science* 197:1092–1094.
- Kornack DR, Rakic P (1999): Continuation of neurogenesis in the hippocampus of the adult macaque monkey. *Proc Natl Acad Sci USA* 96:5768–5763.
- Laywell ED, Rakic P, Kukekov VG, Holland EC, Steindler DA (2000): Identification of a multipotent astrocytic stem cell in the immature and adult mouse brain. *Proc Natl Acad Sci USA* 97:13883–13888.

- Lee SM, Tole S, Grove E, McMahon AP (2000): A local Wnt-3a signal is required for development of the mammalian hippocampus. *Development* 127:457–467.
- Lendahl U, Zimmerman LB, McKay RD (1990): CNS stem cells express a new class of intermediate filament protein. *Cell* 60:585–595.
- Li G, Pleasure SJ (2005): Morphogenesis of the dentate gyrus: what we are learning from mouse mutants. *Dev Neurosci* 27:93–99.
- LoTurco JJ, Sarkisian MR, Cosker L, Bai J (2003): Citron kinase is a regulator of mitosis and neurogenic cytokinesis in the neocortical ventricular zone. *Cereb Cortex* 13:588–591.
- Lu M, Grove EA, Miller RJ (2002): Abnormal development of the hippocampal dentate gyrus in mice lacking the CXCR4 chemokine receptor. *Proc Natl Acad Sci USA* 99:7090–7095.
- Merkle FT, Tramontin AD, García-Verdugo JM, Alvarez-Buylla A (2004): Radial glia give rise to adult neural stem cells in the subventricular zone. *Proc Natl Acad Sci USA* 101:17528–17532.
- Palmer TD, Takahashi J, Gage FH (1997): The adult rat hippocampus contains primordial neural stem cells. *Mol Cell Neurosci* 8:389–404.
- Parent JM, Yu TW, Leibowitz RT, Geschwind DH, Sloviter RS, Lowenstein DH (1997): Dentate granule cell neurogenesis is increased by seizures and contributes to aberrant network reorganization in the adult rat hippocampus. *J Neurosci* 17:3727–3738.
- Rickmann M, Amaral DG, Cowan WM (1987): Organization of radial glial cells during the development of the rat dentate gyrus. *J Comp Neurol* 264:449–479.
- Roberts MR, Bittman K, Li WW, French R, Mitchell B, LoTurco JJ, D'Mello SR (2000): The flathead mutation causes CNS-specific developmental abnormalities and apoptosis. *J Neurosci* 20:2295–2306.
- Sarkisian MR, Frenkel M, Li W, Oborski JA, LoTurco JJ (2001): Altered interneuron development in the cerebral cortex of the flathead mutant. *Cereb Cortex* 11:734–743.
- Sarkisian MR, Li W, Di Cunto F, D'Mello SR, LoTurco JJ (2002): Citron-kinase, a protein essential to cytokinesis in neuronal progenitors, is deleted in the flathead mutant rat. *J Neurosci* 22:RC217.
- Sarkisian M, Rattan S, D'Mello S, LoTurco J (1999): Characterization of seizures in the flathead rat: a new genetic model of epilepsy in early postnatal development. *Epilepsia* 40:394–400.
- Schlessinger AR, Cowan WM, Gottlieb DI (1975): An autoradiographic study of the time of origin and the pattern of granule cell migration in the dentate gyrus of the rat. *J Comp Neurol* 159:149–175.
- Seri B, García-Verdugo JM, Collado-Morente L, McEwen BS, Alvarez-Buylla A (2004): Cell types, lineage, and architecture of the germinal zone in the adult dentate gyrus. *J Comp Neurol* 478:359–378.
- Seri B, Garcia-Verdugo JM, McEwen BS, Alvarez-Buylla A (2001): Astrocytes give rise to new neurons in the adult mammalian hippocampus. *J Neurosci* 21:7153–7160.
- Zhou CJ, Zhao C, Pleasure SJ (2004): Wnt signaling mutants have decreased dentate granule cell production and radial glial scaffolding abnormalities. *J Neurosci* 24:121–126.