References

- Grebe S, Ichida F, Grabitz R, Bultmann B, Heideman S, von Kaisenberg CS: Reversed pulmonary artery flow in isolated noncompaction of the ventricular myocardium. *Fetal Diagn Ther* 2006;22:29–32.
- Finsterer J, Stöllberger C, Krugluger W: Positive troponin-T in noncompaction is associated with neuromuscular disorders and poor outcome. *Clin Res Cardiol* 2006;in press.
- Stöllberger C, Finsterer J, Blazek G: Left ventricular hypertrabeculation/noncompaction and association with additional cardiac abnormalities and neuromuscular disorders. *Am J Cardiol* 2002;90:899–902.
- Finsterer J, Stöllberger C, Blazek G: Neuromuscular implications in left ventricular hypertrabeculation/noncompaction. *Int J Cardiol* 2006;110:288–300.
- Johnson MT, Zhang S, Gilkeson R, Ameduri R, Siwik E, Patel CR, Chebotarev O, Kenton AB, Bowles KR, Towbin JA, Robin NH, Brozovich F, Hoit BD: Intrafamilial variability of noncompaction of the ventricular myocardium. *Am Heart J* 2006;151:1012, e7–e14.
- Finsterer J, Stöllberger C, Schubert B: Acquired left ventricular hypertrabeculation/noncompaction in mitochondriopathy. *Cardiology* 2004;102:228–230.

Reply

C.S. von Kaisenberg

Department of Obstetrics and Gynecology, University of Schleswig-Holstein, Campus Kiel, Kiel, Germany

The comments made by Finsterer and Stöllberger raise the question if they have been made to actually contribute critical, but helpful and supportive comments, or whether they have been made in order to make the point that actually Finsterer published most of the work on LVHT and quoted his own references. Even reviewers, although anonymously, usually give much more positive comments.

Finsterer is questioning the diagnosis and suggesting to include first-degree family members because this may reduce costs and facilitate genetic investigations by narrowing the broad range of possible differential diagnosis.

The diagnosis was made by the Institute of Pathology, University of Schleswig-Holstein, Campus Kiel, Germany, and was confirmed by a reference Institute of Pathology at the University of Tübingen, Germany. Because the family history was negative, genetic investigations in healthy first-degree family members would be indicated only when a mutation in the index case, the affected fetus, has been found. Otherwise, we would not know what to look for and it would increase costs dramatically.

Finsterer on *candidate genes*: The authors suspected mutations in the taffazine, dystrobrevin, desmin, FKBP12, syntrophin, cypher/zasp genes in the fetus with LVHT and carried out molecular genetic investigations for appropriate mutations. However, mutations in the desmin, FKBP12, and syntrophin genes have not

been reported to be associated with human LVHT.

Reply Ichida: *Desmin*: defects produce the myofibrillar or desmin-related myopathy, it was therefore considered a candidate gene for the present investigation (source: OMIM). *FKBP12*: Deletion of the FK506-binding protein-1A (FKBP12) gene results in INVM associated with congenital heart disease in mice, although mutations have not yet been reported in patients. It was therefore considered a candidate gene for the present investigation (source: OMIM). *Syntrophin*: Interest in syntrophin came first from its location at the neuromuscular junction and later from the demonstration that it is directly associated with dystrophin. It was therefore considered a candidate gene for the present investigation (source: OMIM).

All the investigated candidate genes using mutation analysis are important genes to investigate left ventricular noncompaction (LVNC) patients for disease-causing mutations using the 'final common pathway' hypothesis [Towbin JA: Etiologies of cardiomyopathy and heart failure. *Nature Med* 1999].

Finsterer on *further candidate genes*: It is advisable to investigate for further mutations in the DMPK (myotonic dystrophy 1), AMPD1 (myoadenylate-deaminase deficiency), dystrophin (dystrophinopathies), mitochondrial (mitochondrial disorders), or PMP22 (Charcot-Marie-Tooth disease) because LVHT is associated with neuromuscular disorders in up to 80%.

Reply von Kaisenberg: Yes, if you have the funding and the fetal DNA.

Finsterer on *skeletal muscle and blood components*: Skeletal muscle was not histologically investigated for myopathic lesions; neutropenia, a non-neuromuscular feature of this disorder, was not investigated in this postmortem fetus; creatinekinase or troponin-T, frequently elevated in LVHT, was not determined.

Reply von Kaisenberg: True, but this was a 24-weeks fetus when the diagnosis of INVM was made and no blood was available to investigate neutropenia, CK or troponin-T; further, DNA was extracted from liver, lung and kidney tissue, but the total amount was limited, therefore investigations of further candidate genes, such as the large dystrophin gene, was difficult macroscopically; nothing suggested muscular dystrophy.

Finsterer on *electron microscopy*: Were there any abnormalities in any of the cellular compartments, in particular the nucleus, the mitochondria, the contractile apparatus, or the cell membranes? Was there abnormal material within the cytosol or any of the cellular compartments? Left ventricle LVHT rarely affects the interventricular septum, where was it localized?

Reply Bültmann: Electron microscopy detected no abnormalities in any of the cellular compartments of endothelial cells and cardiac myocytes, e.g. in the nucleus, mitochondria, the cytoskeleton, the cell membrane and the sarcoplasm. LVHT

was localized in the free wall of the left ventricle. At autopsy, neither abnormalities of the coronary arteries nor associated defects were noted.

Finsterer on *pathophysiology of hemodynamics*: Explain the pathophysiology of LVHT and reversed pulmonary artery flow.

Reply Grabitz: In cases of very poor right ventricular or global pump function, there may be functional pulmonary atresia with pulmonary regurgitation. Many cases of right heart disease are associated with reverse flow in the ductus venosus, also found in the present case.

Finsterer on the *family pedigree*: Since LVHT occurs familiarly in the majority of

the cases, it would be necessary to look if any of the members of the fetus' family also had LVHT. Sequencing of first-degree family members may reduce costs and facilitate genetic investigations by narrowing the broad range of possible differentials.

Reply Heidemann: The family history was negative, there was no further index patient; therefore, sequencing of unaffected first-degree family members would not have made sense.

Finsterer on the *terminology* LVHT, LVNC, INVM: Finsterer disagrees that the defect was isolated (left ventricular hypertrabeculation, isolated (LVHT); left ventricular noncompaction, isolated (LVNC);

isolated noncompaction of the ventricular myocardium (INVM)).

Reply von Kaisenberg: Ichida et al. 2001 called this isolated left ventricular noncompaction, although it was associated with a dilated, mildly hypertrophic heart with poor systolic function on echocardiogram and clinical heart failure.

All of the above terms may be used to describe the identical pathology and in cardiomyopathy there may be deviations from normal pathology, but ventricular hypertrabeculation may be isolated.