

Original Paper

Comparison between Cystatin C- and Creatinine-Estimated Glomerular Filtration Rate in Cardiology Patients

Axel Åkerblom^{a–c} Johanna Helmersson-Karlqvist^a Mats Flodin^a
Anders Larsson^a

^aDepartment of Medical Sciences, University Hospital, Uppsala University, and
^bUppsala Clinical Research Center, Uppsala, Sweden; ^cDuke Clinical Research
Institute, Durham, N.C., USA

Key Words

Cystatin C · Creatinine · Glomerular filtration rate · Heart · Kidney · CKD-EPI · Sweden

Abstract

Objective: Estimation of the glomerular filtration rate (GFR) is essential for identification, evaluation and risk prediction in patients with kidney disease. Estimated GFR (eGFR) is also needed for the correct dosing of drugs eliminated by the kidneys and to identify high-risk individuals in whom coronary angiography or other procedures may lead to kidney failure. Both cystatin C and creatinine are used for the determination of GFR, and we aimed to investigate if eGFR by the two methods differ in cardiology patients. **Methods:** We compared cystatin C and creatinine (CKD-EPI) eGFR calculated from the same request from a cardiology outpatient unit (n = 2,716), a cardiology ward (n = 980), a coronary care unit (n = 1,464), and an advanced coronary care unit (n = 518) in an observational, cross-sectional study. **Results:** The median creatinine eGFR results are approximately 10 ml/min/1.73 m² higher than the median cystatin C eGFR that is up to 90 ml/min/1.73 m², irrespective of the level of care. Creatinine eGFR resulted in a less advanced eGFR category in the majority of patients with a cystatin C eGFR <60 ml/min/1.73 m². **Conclusions:** Our study demonstrates a difference between creatinine and cystatin C eGFR in cardiology patients. It is important to be aware of which marker is used for the reported eGFR to minimize erroneous interpretations of the test results, as this could lead to under- or overmedication. Further studies are needed to determine the best method of estimating the GFR in cardiology units.

© 2015 S. Karger AG, Basel

Anders Larsson
Department of Medical Sciences, University Hospital, Uppsala University
Entrance 61, 2nd floor, Akademiska Sjukhuset
SE-751 85 Uppsala (Sweden)
E-Mail anders.larsson@akademiska.se

Introduction

Decreased renal function often accompanies congestive heart failure, and this interdependent relationship has become known as the cardiorenal syndrome [1–3]. Studies have documented a strong association between chronic kidney disease (CKD) and cardiovascular disease (CVD) morbidity and mortality [4, 5]. Cardiovascular mortality is 10–20 times more frequent in renal failure patients [6], and over 40% of the deaths in patients with end-stage renal disease are due to CVD [7]. A meta-analysis showed that patients with end-stage renal disease are more likely to die from CVD [8]. Even a mildly to moderately decreased renal function leads to a significantly increased CVD morbidity and mortality [9–13].

Estimated glomerular filtration rate (eGFR) is used for the diagnosis, staging, and prognosis of kidney disease, but is also an important part of drug dosing and risk stratification for clinical procedures and future outcomes. Assessment of kidney damage and dysfunction is thus an integral component of clinical medicine, endorsed by international guidelines [14, 15]. eGFR is usually calculated from plasma creatinine values using Modification of Diet in Renal Disease [16] or CKD-EPI [17] equations. Cystatin C has been shown to be a valuable eGFR marker, and cystatin C has also been reported to be a better marker than creatinine for mortality predictions [18].

A problem with both cystatin C and creatinine equations is that they are developed based on the measurement of GFR using exogenous markers in patients with known or suspected kidney disorders. An equation based on one patient population does not necessarily have to be accurate in other populations [19]. Considering the role of creatinine and cystatin C as GFR markers, the aim was to compare eGFR based on the two markers in different patient populations with the common feature of cardiac disease. We thus compared the creatinine-based eGFR utilizing the CKD-EPI equation with the cystatin C-based eGFR using the new CAPA (Caucasian, Asian, pediatric, adult) equation in patient populations treated at an outpatient cardiology clinic, a general cardiology ward, a cardiology care unit (CCU), and an acute coronary care unit (ACCU). Only samples with simultaneous requests for both creatinine and cystatin C were included in the study. The eGFR equations were originally constructed using patients younger than many of those seeking health care. We thus also wanted to study possible differences between creatinine and cystatin C eGFR in relation to patient age in different groups of cardiology patients.

Methods

Study Population

The comparison was performed with consecutive routine requests for creatinine and cystatin C at the Department of Clinical Chemistry, Uppsala University Hospital, Uppsala, Sweden, between 2011 and 2012. Only samples with requests for both analytes were extracted from the database, and only the first result for each patient was included in the study. The comparisons were limited to samples from a general cardiology outpatient unit, a general cardiology ward with mainly arrhythmia and heart failure patients, a CCU (predominantly patients with acute coronary syndromes), and an ACCU with mainly patients requiring postoperative and intensive care.

Creatinine and Cystatin C Assays

Plasma creatinine was measured by an enzymatic method on an Architect Ci8200 analyzer (Abbott Laboratories, Abbott Park, Ill., USA) and reported using SI units ($\mu\text{mol/l}$). The method is Isotope Dilution Mass Spectrometry (IDMS) calibrated in collaboration with the Swedish external quality assurance organization (Equalis, Uppsala, Sweden). The total analytical imprecision of the creatinine method was 0.7% at 75 $\mu\text{mol/l}$ and 0.9% at 346 $\mu\text{mol/l}$. $\text{eGFR}_{\text{CKD-EPI}}$ was calculated from creatinine using the CKD-EPI formula [17].

Table 1. The difference between eGFR_{CKD-EPI} and eGFR_{CystC} in the GFR intervals <30, 30–59, 60–89, and ≥90 ml/min/1.73 m²

	<30	30–59	60–89	≥90
<i>Outpatient cohort</i>				
Median eGFR _{CKD-EPI}	34	60	82	95
Median eGFR _{CystC}	23	48	73	100
eGFR difference	11	12	9	-5
n (%)	190 (7.0)	886 (32.7)	1,188 (43.9)	443 (16.4)
<i>Cardiology ward</i>				
Median eGFR _{CKD-EPI}	35	60	84	102
Median eGFR _{CystC}	22	44	74	101
eGFR difference	13	16	10	1
n (%)	146 (15.0)	329 (33.9)	329 (33.9)	167 (17.2)
<i>CCU</i>				
Median eGFR _{CKD-EPI}	30	59	85	96
Median eGFR _{CystC}	21	47	75	102
eGFR difference	9	12	10	-6
n (%)	146 (10.0)	388 (26.7)	565 (38.8)	356 (24.5)
<i>ACCU</i>				
Median eGFR _{CKD-EPI}	26	52	84	103
Median eGFR _{CystC}	21	41	72	103
eGFR difference	5	11	12	0
n (%)	146 (28.2)	201 (38.9)	108 (20.9)	62 (12.0)

The patients are divided into the eGFR groups according to cystatin C eGFR.

Serum cystatin C measurements were performed on an Architect Ci8200 with reagents from Gentian (Moss, Norway). The total analytical imprecision of the cystatin C method was 1.7% at 0.77 mg/l and 1.1% at 1.25 mg/l. The cystatin C method used was calibrated according to the new certified reference material ERM-DA471/IFCC [20].

The CAPA equation was used for calculating cystatin C eGFR: $eGFR_{CystC} = 130 \times \text{cystatin C}^{-1.069} \times \text{age}^{-0.117} - 7$ [21].

Statistical Calculations

Descriptive statistics, linear regression analysis, and figures were made utilizing Excel 2000 (Microsoft Corporation, Seattle, Wash., USA). The GFR partitioning was based on the cystatin C eGFR. Samples with cystatin C eGFR values >150 ml/min/1.73 m² were excluded from the statistical analysis.

Results

The creatinine and cystatin C samples from the cardiology outpatient unit [n = 2,716; 905 females and 1,807 males; median age 67 years (range 18–95)], the cardiology ward [n = 980; 363 females and 617 males; median age 67 years (range 18–94)], the CCU [n = 1,464; 469 females and 995 males; median age 67 years (range 18–99)], and the ACCU [n = 518; 158 females and 360 males; median age 66 years (range 18–89)] were analyzed between 2011 and 2012.

The differences between eGFR by eGFR_{CKD-EPI} and eGFR_{CystC} per subgroups of eGFR_{CystC} are reported in table 1. eGFR_{CKD-EPI} values were approximately 10 ml/min/1.73 m² higher

Fig. 1. Plot showing patient age (x-axis) versus the difference between $eGFR_{CKD-EPI}$ and $eGFR_{CystC}$ (y-axis) in individual patient samples from the cardiology outpatient clinic.





Fig. 2. Plot showing patient age (x-axis) versus the difference between $eGFR_{CKD-EPI}$ and $eGFR_{CystC}$ (y-axis) in individual patient samples from the cardiology ward.



than $eGFR_{CystC}$, which were up to 90 ml/min/1.73 m², with small differences in the median $eGFR$ over 90 ml/min/1.73 m². Age variation had limited effects on the difference between $eGFR_{CKD-EPI}$ and $eGFR_{CystC}$ in the different subgroups (fig. 1–4).

Comparison of CKD Staging Based on $eGFR_{CKD-EPI}$ and $eGFR_{CystC}$

In the patient groups with $eGFR_{CystC} < 30$ ml/min/1.73 m², 69.5% (cardiology outpatient unit), 63.7% (cardiology ward), 50.0% (CCU), and 36.3% (ACCU) of the patients were reclassified to a less advanced GFR category when using $eGFR_{CKD-EPI}$ (table 2). Out of the patients with an $eGFR_{CystC}$ of 30–44 ml/min/1.73 m², 71.3% were reclassified to a less advanced $eGFR$ category, and 1.5% were reclassified to a lower group by $eGFR_{CKD-EPI}$ in the outpatient unit. The corresponding figures for the cardiology ward were 78.9 and 1.8%, for the CCU 62.1 and 1.8%, and for the ACCU 48.4 and 7.1%. The percentages of reclassified patients with $eGFR_{CystC}$ 45–59 ml/min/1.73 m² were 70.1 and 3.5% in the outpatient unit, 75.3 and 1.9% in the cardiac ward, 66.2 and 5.9% in the CCU, and 60.0 and 9.3% in the ACCU.

Discussion

CKD is a growing global health problem. The incidence of CKD is expected to continue to rise in the future, partly due to an increased number of patients with diabetic nephropathy as a consequence of increased age and obesity [22]. Improvements in the detection and risk

Fig. 3. Plot showing patient age (x-axis) versus the difference between $eGFR_{CKD-EPI}$ and $eGFR_{CystC}$ (y-axis) in individual patient samples from the CCU.





Fig. 4. Plot showing patient age (x-axis) versus the difference between $eGFR_{CKD-EPI}$ and $eGFR_{CystC}$ (y-axis) in individual patient samples from the ACCU.



stratification of CKD patients are crucial, and we expect the classification of CKD by eGFR to remain important. It is seldom possible to measure GFR in real-life patient settings with an exogenous GFR marker. Thus, we will continue to rely on inexpensive endogenous GFR markers such as creatinine and cystatin C. We usually classify patients according to eGFR and do not distinguish between the molecules used for the measurements. Basically, we assume all GFR markers to give the same results, and this would be highly desirable. However, as the eGFR equations are calibrated against measured GFR, they are highly influenced by the population that they are derived from. When we use the same equations in other patient groups with other features, consequently the results may differ. Ethnic differences in creatinine eGFR are often addressed, and there are several ethnicity-specific equations or factors [16, 17, 23, 24]. In contrast, there is limited discussion on the differences between patient groups within a hospital.

Considering the importance of eGFR, we compared cystatin C and creatinine eGFR in different groups of cardiology patients. We used cystatin C eGFR to divide the patients into different GFR strata in accordance with the CKD classification [25]. There are no GFR measurements with iothexol, iothalamate or chromium-51-EDTA in this study, so we do not have the true GFR for the patients. Nevertheless, our study demonstrates a clear difference between creatinine and cystatin C eGFR in cardiology patients. The difference between the two measurements is fairly consistent in patients treated at the different cardiology units. The median creatinine eGFR results were approximately 10 ml/min/1.73 m² higher than the median cystatin C eGFR, which was up to 90 ml/min/1.73 m². This bias is not negligible. If

Table 2. Reclassification of patients when using eGFR_{CKD-EPI} instead of eGFR_{CystC}

eGFR _{CystC}	eGFR _{CKD-EPI} , n			
	<30	30–44	45–59	≥60
<i>Outpatient unit</i>				
<30 (190)	58	93	34	5
30–<45 (341)	5	93	165	78
45–<60 (545)	0	19	144	382
>60 (1,628)	0	2	69	1,557
<i>Cardiology ward</i>				
<30 (146)	53	59	26	8
30–<45 (171)	3	33	89	46
45–<60 (158)	0	3	36	119
>60 (496)	0	0	18	478
<i>CCU</i>				
<30 (146)	73	48	21	4
30–<45 (169)	3	61	67	38
45–<60 (219)	1	12	61	145
>60 (921)	0	2	16	903
<i>ACCU</i>				
<30 (146)	93	40	10	3
30–<45 (126)	9	56	39	22
45–<60 (75)	0	7	23	45
>60 (170)	1	1	11	157

The number of the patients classified in GFR strata by cystatin C and the CKD-EPI classification of the same patients.

eGFR values are ml/min/1.73 m².

we use important clinical decision limits such as 15, 30, 45, and 60 ml/min/1.73 m², a large number of patients will be reclassified depending on which GFR marker is used.

This finding is important and needs to be stressed, as treatment strategies cannot be accepted to be dependent on the marker used for defining the eGFR of the individual patient. We thus recommend that different decision limits for the cystatin C- and creatinine-based eGFR for CVD risk estimation should be considered. The different cardiology units showed similar differences, indicating that the same decision limits could be used in the different units.

Limitations

Our results are based on a large, single-center, tertiary care hospital, and hence they may not be reproducible in other regions and countries due to biochemical or analytical differences. The new CAPA equation is based on methods calibrated with the recently developed ERM-DA471/IFCC reference material. CAPA is an assay-independent cystatin C-based estimating equation for GFR using 7 different cystatin C assays. This equation is validated in a North European/Asian community setting and may not be applicable in other regions.

The patients admitted to a cardiology outpatient clinic or cardiology ward may not always end up with a cardiology diagnosis. We did not scrutinize patient diagnoses, but rather decided to include all patients with a cardiology affiliation who had dually requested creatinine and cystatin C measurements.

Creatinine and cystatin C may differ in regard to the time of sampling. Cystatin C reacts 1–2 days earlier to acute renal failure, a condition that may occur in hospitalized patients. Thus, the difference observed in this study could partly be due to the early response of cystatin C to acute renal failure [26].

Conclusion

Our study demonstrates a difference between creatinine and cystatin C eGFR in cardiology patients. It is important to be aware of which marker is used for the reported eGFR to minimize erroneous interpretations of test results, as this could lead to under- or overmedication. Further studies are needed to determine the best method of estimating GFR in cardiology units.

Acknowledgement

This study was supported by the Uppsala University Hospital Research Fund.

Disclosure Statement

The study was investigator initiated and investigator driven. A.Å. received institutional grants from AstraZeneca. The other authors declare that they have no conflicts of interest.

References

- 1 Ronco C, Di LL: Cardiorenal syndrome. *Heart Fail Clin* 2014;10:251–280.
- 2 Virzi G, Day S, de Cal M, et al: Heart-kidney crosstalk and role of humoral signaling in critical illness. *Crit Care* 2014;18:201.
- 3 Braam B, Joles JA, Danishwar AH, Gaillard CA: Cardiorenal syndrome – current understanding and future perspectives. *Nat Rev Nephrol* 2014;10:48–55.
- 4 Weiner DE, Tighiouart H, Amin MG, et al: Chronic kidney disease as a risk factor for cardiovascular disease and all-cause mortality: a pooled analysis of community-based studies. *J Am Soc Nephrol* 2004;15:1307–1315.
- 5 McCullough PA, Verrill TA: Cardiorenal interaction: appropriate treatment of cardiovascular risk factors to improve outcomes in chronic kidney disease. *Postgrad Med* 2010;122:25–34.
- 6 Raine AE, Margreiter R, Brunner FP, et al: Report on management of renal failure in Europe, XXII, 1991. *Nephrol Dial Transplant* 1992;7(suppl 2):7–35.
- 7 Moody WE, Edwards NC, Chue CD, et al: Arterial disease in chronic kidney disease. *Heart* 2013;99:365–372.
- 8 Sarnak MJ, Levey AS, Schoolwerth AC, et al: Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Circulation* 2003;108:2154–2169.
- 9 Jernberg T, Lindahl B, James S, et al: Cystatin C: a novel predictor of outcome in suspected or confirmed non-ST-elevation acute coronary syndrome. *Circulation* 2004;110:2342–2348.
- 10 Nerpin E, Ingelsson E, Riserus U, et al: The combined contribution of albuminuria and glomerular filtration rate to the prediction of cardiovascular mortality in elderly men. *Nephrol Dial Transplant* 2011;26:2820–2827.
- 11 Nitsch D, Grams M, Sang Y, et al: Associations of estimated glomerular filtration rate and albuminuria with mortality and renal failure by sex: a meta-analysis. *BMJ* 2013;346:f324.
- 12 Hallan SI, Matsushita K, Sang Y, et al: Age and association of kidney measures with mortality and end-stage renal disease. *JAMA* 2012;308:2349–2360.
- 13 Mahmoodi BK, Matsushita K, Woodward M, et al: Associations of kidney disease measures with mortality and end-stage renal disease in individuals with and without hypertension: a meta-analysis. *Lancet* 2012;380:1649–1661.

- 14 Fox KA, Goodman SG, Anderson FA Jr, et al: From guidelines to clinical practice: the impact of hospital and geographical characteristics on temporal trends in the management of acute coronary syndromes. The Global Registry of Acute Coronary Events (GRACE). *Eur Heart J* 2003;24:1414–1424.
- 15 Anderson JL, Adams CD, Antman EM, et al: 2012 ACCF/AHA focused update incorporated into the ACCF/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;61:e179–e347.
- 16 Levey AS, Coresh J, Greene T, et al: Expressing the Modification of Diet in Renal Disease Study equation for estimating glomerular filtration rate with standardized serum creatinine values. *Clin Chem* 2007;53:766–772.
- 17 Levey AS, Stevens LA, Schmid CH, et al: A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604–612.
- 18 Shlipak MG, Matsushita K, Arnlov J, et al: Cystatin C versus creatinine in determining risk based on kidney function. *N Engl J Med* 2013;369:932–943.
- 19 Larsson A, Flodin M, Hansson LO, Carlsson L: Patient selection has a strong impact on cystatin C and Modification of Diet in Renal Disease (MDRD) estimated glomerular filtration rate. *Clin Biochem* 2008;41:1355–1361.
- 20 Grubb A, Blirup-Jensen S, Lindstrom V, et al: First certified reference material for cystatin C in human serum ERM-DA471/IFCC. *Clin Chem Lab Med* 2010;48:1619–1621.
- 21 Grubb A, Horio M, Hansson LO, Bjork J, Nyman U, Flodin M, et al: Generation of a new cystatin C-based estimating equation for glomerular filtration rate by use of 7 assays standardized to the international calibrator. *Clin Chem* 2014;60:974–986.
- 22 National Kidney Foundation: KDOQI clinical practice guideline for diabetes and CKD: 2012 update. *Am J Kidney Dis* 2012;60:850–886.
- 23 Matsuo S, Imai E, Horio M, et al: Revised equations for estimated GFR from serum creatinine in Japan. *Am J Kidney Dis* 2009;53:982–992.
- 24 Björk J, Grubb A, Sterner G, Nyman U: Revised equations for estimating glomerular filtration rate based on the Lund-Malmö Study cohort. *Scand J Clin Lab Invest* 2011;71:232–239.
- 25 National Kidney Foundation: KDOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002;39:S1–S266.
- 26 Herget-Rosenthal S, Marggraf G, Hüsing J, Göring F, Pietruck F, Janssen O, et al: Early detection of acute renal failure by serum cystatin C. *Kidney Int* 2004;66:1115–1122.